

the measured values. As shown in the Fig. 4(inset), this difference can be attributed to an experimental increase in  $r_0 \sim \omega_x^{-2/3}$  due to the measured decrease of  $\omega_x$  with  $\alpha^2$ . This implies that the average particle charge  $q$  is effectively constant, as previously observed for constant neutral pressure [8, 19]. This increase in  $r_0$  with  $\alpha^2$  is not as apparent for  $y_{rms}$  since  $y_{rms} \lesssim 0.2x_{rms}$  and  $y_{rms} \rightarrow 0$  as  $\alpha^2$  increases.

In summary, we have provided direct evidence that the width of a Yukawa cluster exhibits power law behavior for the 1D to 2D zigzag transition caused by decreasing the confining well anisotropy parameter  $\alpha^2$ , confirming a previous prediction [16]. Experiments were performed using a dusty plasma with  $n = 6$  particles confined in the biharmonic well above a rectangular depression. The width  $d$  of the rectangular depression was increased while the plasma remained on to decrease  $\alpha^2$  while the Debye shielding parameter  $\kappa$  remained essentially constant. The dependence of  $\alpha^2$  on  $d$  was accurately determined by measuring the c.m. frequencies of the dusty plasma. A transition from the zigzag configuration to an elliptical configuration was also observed. The cluster width was found to be in excellent agreement with the predictions of a model which assumes identical particles confined in a 2D biharmonic well and interacting through a Yukawa potential. From the fit to the model we found the Debye length is comparable to the inter-particle distance, so that Debye shielding significantly effects the physics of these clusters.

## Acknowledgments

Portions of this paper are taken from A. L. M.'s undergraduate physics thesis.

- 
- [1] F. F. Chen, Introduction to Plasma Physics (Plenum, New York, 1974), p. 3.
  - [2] M. Lampe, G. Joyce, G. Ganguli and V. Gavrilchaka, Phys. Plasmas **7**, 3851 (2000).
  - [3] A. Homann, A. Melzer, S. Peters and A. Piel, Phys. Rev. E **56**, 7138 (1997).
  - [4] T. Misawa, N. Ohno, K. Asano, M. Sawai, S. Takamura and P. K. Kaw, Phys. Rev. Lett. **86**, 1219 (2001).
  - [5] B. Liu and J. Goree, Phys. Rev. E **71**, 046410 (2005).
  - [6] W.-T. Juan, Z.-H. Huang, J.-W. Hsu, Y.-J. Lai and L. I, Phys. Rev. E **58**, R6947 (1998).
  - [7] T. E. Sheridan, J. Phys. D: Appl. Phys **39**, 693 (2006).