

- [16] J. A. Leegwater, The Journal of Physical Chemistry **100**, 14403 (1996).
- [17] A. G. Redfield, IBM Journal of Research and Development **1**, 19 (1957).
- [18] W. T. Pollard and R. A. Friesner, The Journal of Chemical Physics **100**, 5054 (1994).
- [19] A. Nitzan, *Chemical Dynamics in Condensed Phases: Relaxation, Transfer, and Reactions in Condensed Molecular Systems* (Oxford University Press, USA, 2006).
- [20] M. Richter, K. J. Ahn, A. Knorr, A. Schliwa, D. Bimberg, M. E. Madjet, and T. Renger, physica status solidi (b) **243**, 2302 (2006).
- [21] T. Renger and R. A. Marcus, The Journal of Chemical Physics **116**, 999710019 (2002).
- [22] T. Calarco, A. Datta, P. Fedichev, E. Pazy, and P. Zoller, Physical Review A **68**, 012310 (2003).
- [23] We consider a typical GaAs QD with deformation potentials $D_e = -14.6$ eV, $D_h = -4.8$ eV, mass density of $\rho = 5.4$ g/cm³, speed of sound $u = 5000$ cm/s, and radius of $l = 5$ nm.
- [24] O. Madelung, ed., *Semiconductors: data handbook* (Springer, Berlin, 2004), 3rd ed.
- [25] G. D. Scholes, Annual Review of Physical Chemistry **54**, 57 (2003), ISSN 0066-426X.
- [26] A. L. Rogach, T. A. Klar, J. M. Lupton, A. Meijerink, and J. Feldmann, Journal of Materials Chemistry **19**, 1208 (2009).
- [27] I. L. Medintz and H. Mattoussi, Physical Chemistry Chemical Physics **11**, 17 (2009).
- [28] D. Kim, S. Okahara, M. Nakayama, and Y. Shim, Physical Review B **78**, 153301 (2008).
- [29] V. I. Klimov, Annual Review of Physical Chemistry **58**, 635 (2007).
- [30] A. Nazir, B. W. Lovett, S. D. Barrett, J. H. Reina, and G. A. D. Briggs, Physical Review B **71**, 045334 (2005).