

Computación

Científica

Marcos Iturbe

Ejercicio 1

$p=8$

$$f(x) = x(1 - \cos x)$$

$$\alpha=0 \quad f'(x) = (1 - \cos x) + x(\sin x)$$

$$f(\alpha) = 0 \quad f'(\alpha) = 0$$

$$f''(x) = \sin x + \sin x + x \cos x$$

$$f''(\alpha) = 0$$

$$x_n \rightarrow x_{n-1} - \frac{f(x_{n-1})}{f'(x_{n-1})}$$

$$x_0 \rightarrow 1$$

$$x_1 \rightarrow 1 - \frac{0,45964769}{1,30116868} = 0,646704$$

$$x_2 \rightarrow 0,646704 - \frac{0,43058625}{0,59177669} = 0,42603588$$

$$x_3 \rightarrow 0,42603588 - \frac{0,38082866 \times 10^{-1}}{0,26545428} = 0,28257289$$

$$x_4 \rightarrow 0,28257289 - \frac{0,11206495 \times 10^{-1}}{0,11844784} = 0,18798607$$

$$x_5 \rightarrow 0,18798607 - \frac{3,3178273 \times 10^{-3}}{0,52748300 \times 10^{-1}} = 0,12520059$$

$$x_6 \rightarrow 0,12520059 - \frac{9,7997024 \times 10^{-4}}{2,3461629 \times 10^{-2}} = 8,3430675 \times 10^{-2}$$

$$x_7 \rightarrow 8,3430675 \times 10^{-2} - \frac{2,9019862 \times 10^{-4}}{10,430926 \times 10^{-2}} = 5,5609692 \times 10^{-2}$$

$$\epsilon_6/\epsilon_5 = \frac{-4,1769915 \times 10^{-2}}{-6,278548 \times 10^{-2}} = 0,66432422$$

$$\epsilon_7/\epsilon_6 = \frac{-2,7820983 \times 10^{-2}}{-4,1709915 \times 10^{-2}} = 0,667011261$$

$$cte \approx 0,66$$

$$\frac{\epsilon_{k+1}}{\epsilon_k} \approx \frac{m-1}{m} \approx 0,66 \Rightarrow m=3$$

* Newton modificado:

$$x_{n+1} = x_n - m \frac{f(x)}{f'(x)}$$

$$x_0 \leftarrow 1$$

$$x_1 \leftarrow 1 - 3 \frac{0,45967769}{1,30116868} = -0,059888$$

$$x_2 \leftarrow -0,05988 - 3 \frac{-1,0736423 \times 10^{-4}}{5,3757436 \times 10^{-3}} = 3,5932374 \times 10^{-5}$$

Ejercicio 2

Distancia recorrida
entre el $s=11$ y
 $s=16$

$$\int_{11}^{16} v(t) dt$$

$t(s)$	$v(m/s)$
0	0
10	227,04
15	362,78
20	517,35
22,5	602,97
30	901,67

$$v(t) \approx P_2(t) = 12,05 + 17,733t + 0,3766t^2$$

$$v(t) \approx P_3(t) = -4,381 + 21,289t + 0,13065t^2 + 0,005466t^3$$

$$\int_{11}^{16} P_2(t) dt = 12,05t + \frac{17,733t^2}{2} + \frac{0,3766t^3}{3} \Big|_{11}^{16} =$$

$$1604,327167 \text{ m}$$

$$\int_{11}^{16} P_3(t) dt = -4,381t + \frac{21,289t^2}{2} + \frac{0,13065t^3}{3} + \frac{0,005466t^4}{4} \Big|_{11}^{16} =$$

$$1605,066268 \text{ m}$$

aceleración en $t=16$

$$a = \frac{dv}{dt}$$

$$v(t) \approx f_{2,2}(t)$$

$$f_2'(t) = 17,733 + (0,3766t \times 2)$$

$$f_2'(16) = 29,7842 \text{ m/s}^2$$

$$f_3'(t) = 21,289 + 2 \times 0,13065t + 3 \times 0,005466t^2$$

$$f_3'(16) = 29,667688 \text{ m/s}^2$$

Ejercicio 3

$$\sum_{i=1}^n i^3 =$$

$$\varphi(n) \equiv 1 + 7(n-1) + 6(n-1)(n-2) + (n-1)(n-2)(n-3)$$

n	$\varphi(n)$				
1	1				
2	8	7			
3	27	19	6		
4	64	37	9	1	
5	125	61	12	1	0
6	216	91	15	1	0

Ejercicio 4

$$I(f) = \int_0^1 \sqrt{1+2t} \, dt$$

$$h = \frac{1}{2}$$

$$S_1 = \frac{h}{3} \left(f(0) + f(1) + 4f\left(\frac{1}{2}\right) \right) = \frac{h}{3} \left[1 + \sqrt{3} + 4\sqrt{\frac{3}{2}} \right] = 1,271838382$$

$$h = \frac{1}{4}$$

$$S_2 = \frac{h}{3} \left(f(0) + f(1) + 4 \left[f\left(\frac{1}{4}\right) + f\left(\frac{3}{4}\right) \right] + 2f\left(\frac{1}{2}\right) \right) = \frac{h}{3} \left[\sqrt{3} + 1 + 4 \left(\frac{3\sqrt{2}}{4} + \frac{\sqrt{34}}{4} \right) + 2\sqrt{\frac{3}{2}} \right] = 1,271261094$$

$$h = \frac{1}{8}$$

$$S_4 = \frac{h}{3} \left[f(0) + f(1) + 4 \left(f\left(\frac{1}{8}\right) + f\left(\frac{3}{8}\right) + f\left(\frac{5}{8}\right) + f\left(\frac{7}{8}\right) \right) + 2 \left(f\left(\frac{1}{4}\right) + f\left(\frac{3}{4}\right) \right) \right] = \frac{h}{3} \left[1 + \sqrt{3} + 4 \left(\frac{\sqrt{66}}{8} + \frac{\sqrt{82}}{8} + \frac{\sqrt{114}}{8} + \frac{9\sqrt{2}}{8} \right) + 2 \left(\sqrt{\frac{3}{2}} + \frac{3\sqrt{2}}{4} + \frac{\sqrt{34}}{4} \right) \right] = 1,271272865$$

0	$\frac{1}{8}$	$\frac{1}{4}$	$\frac{3}{8}$	$\frac{1}{2}$	$\frac{5}{8}$	$\frac{3}{4}$	$\frac{7}{8}$	1
---	---------------	---------------	---------------	---------------	---------------	---------------	---------------	---

$$\approx -3,84 \times 10^{-5}$$

$$S_1 = 1,271838382$$

$$S_2 = 1,271261094$$

$$S_4 = 1,271272865$$

$$S_{n1} = S_2 + \frac{S_2 - S_1}{4^2 - 1} = 1,271222608$$

$$S_{n2} = 1,27127364$$

$$7,8411 \times 10^{-7}$$

$$S_{n3} = S_{n2} + \frac{S_{n2} - S_{n1}}{4^3 - 1} =$$

$$\epsilon \approx 8,1 \times 10^{-7}$$

$$1,27127445$$

Ejercicio 5

$$\int_1^3 \frac{e^x}{x} dx \simeq 8,038733067$$

$$h = 1$$

$$S_1 = \frac{h}{3} [f(1) + f(3) + 4f(2)] = \frac{1}{3} [2,718281828 + 6,695178174 + 4 \times (3,694528649)] = 8,063857667$$

$$S_2 = 8,040946349$$

$$S_n = S_2 + \frac{S_2 - S_1}{4^2 - 1} = 8,039418928$$

$$\downarrow$$
$$-1,5274 \times 10^{-3} \text{ (5)} \rightarrow \text{tendra 4 cifras significativas}$$

Ejercicio 6

$$I(f) = \int_0^2 (x-1) e^{-x} dx = -\int_0^1 (x-1) e^{-x} dx + \int_1^2 (x-1) e^{-x} dx$$

$$h = \frac{1}{2}$$

$$I(f) \simeq -\frac{1}{6} \left(-1 + 0 + 4 \times (-0,5 e^{-0,5}) \right) + \frac{1}{6} \left(0 + e^{-2} + 4(0,5 e^{-1}) \right)$$

$$= 0,4657761538$$