

SOLITARY-WAVE FAMILIES OF THE MICRO-STRUCTURE PDE

S. ROY CHOUDHURY, JONATHAN LETO

ABSTRACT. The Micro-Structure PDE is a model for ... Since solitary-wave solutions often play a central role in the long-time evolution of an initial disturbance, we consider such solutions here (via normal form approach) within the framework of reversible systems theory. On isolated curves in the relevant parameter region, the delocalized waves reduce to genuine embedded solitons. Directions for future work are also described.

1. INTRODUCTION

The nonlinear dispersive equation [2]

$$(1.1) \quad v_{tt} - bv_{xx} - \frac{\mu}{2} (v^2)_{xx} - \delta (\beta v_{tt} - \gamma v_{xx})_{xx} = 0$$

is known as the Micro-Structure PDE.

BACKGROUND

We use the theory of reversible systems and the method of normal forms to categorize the possible solitary waves of (1.1). As we shall see, several families of solitary waves exist in various regions of parameter space. Our main focus here will be on delineating the possible occurrence and multiplicity of solitary waves in different parameter regimes. Certain delicate questions relating to specific waves or wave families will form the basis of future work.

The remainder of this paper is organized as follows. In Section 2, we delineate the possible families of solitary waves in various parameter domains and on certain important curves using the theory of reversible systems. In Section 3 and 4, we next focus on the various transition curves and derive normal forms in their vicinity to confirm the existence of families of regular or delocalized solitary-wave solutions in their vicinity.

2. SOLITARY WAVES; LOCAL BIFURCATIONS

Solitary waves of (1.1) of the form $v(x, t) = \phi(x - ct) = \phi(z)$ satisfy the fourth-order travelling wave ODE

$$(2.1) \quad \phi_{zzzz} - q\phi_{zz} + p\phi = \mathcal{N}[\phi]$$

where

$$(2.2) \quad \mathcal{N}[\phi] = -\Delta_1 \phi_z^2 - b\Delta_1 \phi \phi_{zz}$$

$$(2.3a) \quad \Delta_1 = \frac{\mu}{\delta(\beta c^2 - \gamma)}$$

$$(2.3b) \quad p = 0$$

Equation (2.1) is invariant under the transformation $z \mapsto -z$ and is thus a reversible system. In this section we shall use the theory of reversible systems to characterize the homoclinic orbits to the fixed point of (2.1), which correspond to pulses or solitary waves of the Micro-Structure PDE in various regions of the (p, q) plane.

The linearized system corresponding to (2.1)

$$(2.4) \quad \phi_{zzzz} - q\phi_{zz} + p\phi = 0$$

has a fixed point

$$(2.5) \quad \phi = \phi_z = \phi_{zz} = \phi_{zzz} = 0$$

Solutions $\phi = ke^{\lambda x}$ satisfy the characteristic equation $\lambda^4 - q\lambda^2 + p = 0$ from which one may deduce that the structure of the eigenvalues is distinct in two regions of (p, q) -space. Since $p = 0$ we have only two possible regions of eigenvalues. We denote C_0 as the positive q axis and C_1 the negative q -axis. First we shall consider the bounding curves C_0 and C_1 and their neighborhoods, then we shall discuss the possible occurrence and multiplicities of homoclinic orbits to (2.5), corresponding to pulse solitary waves of (1.1), in each region:

Near C_0 : The eigenvalues have the structure $\lambda_{1-4} = 0, 0, \pm\lambda$, ($\lambda \in \mathbb{R}$) and the fixed point (2.5) is a saddle-focus.

Near C_1 : Here the eigenvalues have the structure $\lambda_{1-4} = 0, 0, \pm i\omega$, ($\omega \in \mathbb{R}$). We will show by analysis of a four-dimensional normal form in Section 4 that there exists a *sech*² homoclinic orbit near C_1 .

Having outlined the possible families of orbits homoclinic to the fixed point (2.5) of (2.4), corresponding to pulse solitary waves of (1.1), we now derive normal forms near the transition curves C_0 and C_1 to confirm the existence of regular or delocalized solitary waves in the corresponding regions of (p, q) parameter space.

3. NORMAL FORM NEAR C_0 : SOLITARY-WAVE SOLUTIONS

Using (2.4), the curve C_0 , corresponding to $\lambda = 0, 0, \pm\tilde{\lambda}$, is given by

$$(3.1) \quad C_0 : p = 0, q > 0$$

Using (2.3a) this implies

$$(3.2) \quad \frac{c^2 - b}{\delta(\beta c^2 - \gamma)} > 0$$

Denoting ϕ by y_1 , (2.1) may be written as the system

$$(3.3a) \quad \frac{dy_1}{dz} = y_2$$

$$(3.3b) \quad \frac{dy_2}{dz} = y_3$$

$$(3.3c) \quad \frac{dy_3}{dz} = y_4$$

$$(3.3d) \quad \frac{dy_4}{dz} = qy_3 - py_1 - (\Delta_1 y_2^2 + b\Delta_1 y_1 y_3)^T$$

We wish to rewrite this as a first order reversible system in order to invoke the relevant theory [1]. To that end, defining $Y = \langle y_1, y_2, y_3, y_4 \rangle^T$ (3.3) may be written

$$(3.4) \quad \frac{dY}{dz} = L_{pq}Y - F_2(Y, Y)$$

where

$$(3.5) \quad L_{pq} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ q/3 & 0 & 1 & 0 \\ 0 & q/3 & 0 & 1 \\ q^2 - p & 0 & q/3 & 0 \end{pmatrix}$$

Since $p = 0$ for (1.1), we have

$$(3.6) \quad \frac{dY}{dz} = L_{0q}Y - F_2(Y, Y)$$

where

$$(3.7) \quad F_2(Y, Y) = \langle 0, 0, 0, \Delta_1 y_2^2 + b\Delta_1 y_1 y_3 \rangle^T$$

Next we calculate the normal form of (3.6) near C_0 . The procedure is closely modeled on [1] and many intermediate steps may be found there.

3.1. **Near C_0 .** Near C_0 the dynamics reduce to a 2-D Center Manifold

$$(3.8) \quad Y = A\zeta_0 + B\zeta_1 + \Psi(\epsilon, A, B)$$

and the corresponding normal form is

$$(3.9a) \quad \frac{dA}{dz} = B$$

$$(3.9b) \quad \frac{dB}{dz} = b\epsilon A + \tilde{c}A^2$$

Here,

$$(3.10) \quad \epsilon = \left(\frac{q^2}{9} - p \right) - \left(\frac{q}{3} \right)^2 = -p$$

measures the perturbation around C_0 , and

$$(3.11a) \quad \zeta_0 = \langle 1, 0, -q/3, 0 \rangle^T$$

$$(3.11b) \quad \zeta_1 = \langle 0, 1, 0, -2q/3 \rangle^T$$

The linear eigenvalue of (3.9a) satisfies

$$(3.12) \quad \lambda^2 = b\epsilon$$

The characteristic equation of the linear part of (3.6) is

$$(3.13) \quad \lambda^4 - q\lambda^2 - \epsilon = 0$$

Hence, the eigenvalues near zero (the Center Manifold) satisfy $\lambda^4 \ll \lambda^2$ and hence

$$(3.14) \quad \lambda^2 \sim -\frac{\epsilon}{q}$$

Matching (3.12) and (3.14)

$$(3.15) \quad b = -\frac{1}{q}$$

and only the nonlinear coefficient \tilde{c} remains to be determined in the normal form (3.9a).

In order to determine \tilde{c} (the coefficient of A^2 in (3.9a) we calculate $\frac{dY}{dz}$ in two ways and match the $\mathcal{O}(A^2)$ terms.

To this end, using the standard 'suspension' trick of treating the perturbation parameter ϵ as a variable, we expand the function Ψ in (3.8) as

$$(3.16) \quad \Psi(\epsilon, A, B) = \epsilon A \Psi_{10}^1 + \epsilon B \Psi_{01}^1 + A^2 \Psi_{20}^0 + AB \Psi_{11}^0 + B^2 \Psi_{02}^0 + \dots$$

where the subscripts denote powers of A and B , respectively, and the superscript denotes the power of ϵ .

In the first way of computing dY/dz , we take the z derivative of (3.8) (using (3.9a) and (4.4)). The coefficient of A^2 in the resulting expression is $\tilde{c}\zeta_1$. In the second way of computing dY/dz , use (3.8) and (4.4) in (3.4). The coefficient of A^2 in the resulting expression is $L_{0,q}\Psi_{20}^0 - F_2(\zeta_0, \zeta_0)$. Hence

$$(3.17) \quad \tilde{c}\zeta_1 = L_{0,q}\Psi_{20}^0 - F_2(\zeta_0, \zeta_0)$$

Using (3.11a) and (3.7) and denoting $\Psi_{20}^0 = \langle x_1, x_2, x_3, x_4 \rangle$ in (3.17) yields the equations

FIX

$$(3.18a) \quad 0 = x_2$$

$$(3.18b) \quad c = \frac{q}{3}x_1 + x_3$$

$$(3.18c) \quad 0 = \frac{q}{3}x_2 + x_4 \implies x_4 = 0$$

and

$$(3.19) \quad -\frac{2q}{3}\tilde{c} = \frac{q}{3} \left(\frac{q}{3}x_1 + x_3 \right) + \frac{2q}{3} = \frac{q}{3}\tilde{c} + \frac{b\Delta_1}{3}$$

using (3.18b). Hence we obtain

$$(3.20) \quad \tilde{c} = -\frac{b\Delta_1}{3}$$

DETAIL Therefore, the normal form for (1.1) near C_0 is

$$(3.21a) \quad \frac{dA}{dz} = B$$

$$(3.21b) \quad \frac{dB}{dz} = -\frac{\epsilon}{q}A - \frac{b\Delta_1}{3}A^2$$

4. NORMAL FORM NEAR C_1 : POSSIBLE SOLITARY-WAVE SOLUTIONS

Near C_1 the dynamics reduce to a 4-D Center Manifold [1]

$$(4.1) \quad Y = A\zeta_0 + B\zeta_0 + C\zeta_+ + \bar{C}\zeta_- + \Psi(\epsilon, A, B, C, \bar{C})$$

with

$$(4.2a) \quad \frac{dA}{dz} = B$$

$$(4.2b) \quad \frac{dB}{dz} = \bar{\nu}A + b_*A^2 + c_*|C|^2$$

$$(4.2c) \quad \frac{dC}{dz} = id_0C + i\bar{\nu}d_1C + id_2AC$$

where

$$(4.3a) \quad \zeta_{\pm} = \left\langle 1, \lambda_{\pm}, 2q/3, \frac{\lambda_{\pm}}{3}q \right\rangle^T$$

$$(4.3b) \quad \bar{\nu} = -\frac{\epsilon}{q}$$

$$(4.3c) \quad b_* = -\frac{b\Delta_1}{3}$$

$$(4.3d) \quad \lambda_{\pm} = \pm i\sqrt{-q}$$

Comparing to the linearized equations gives $d_0 = \sqrt{-q}$.

If we do a dominant balance argument after the change of variable $\tilde{\epsilon} = \sqrt{-3\alpha}$ on the characteristic equation as $\lambda \rightarrow 0$ then we find $d_1 = \frac{\sqrt{-3\alpha}}{18\alpha^2}$. Using $\alpha = q/3$ we find $d_1 = \frac{\sqrt{-q}}{2q^2}$

To determine b_*, c_* and d_2 we expand

$$(4.4) \quad \Psi(\epsilon, A, B, C, \bar{C}) = \epsilon A \Psi_{1000}^1 + \epsilon B \Psi_{0100}^1 + A^2 \Psi_{2000}^0 + AB \Psi_{1100}^0 + AC \Psi_{1010}^0 + \epsilon C \Psi_{0010}^1 + \dots$$

Now we use (4.2b), (4.2c), (4.2c) in (4.4) when we calculate $\frac{dY}{dz}$. Matching coefficients yields

$$(4.5a) \quad \mathcal{O}(A^2) : \quad b_* \zeta_1 = L_{0q} \Psi_{2000}^0 - F_2(\zeta_0, \zeta_0)$$

$$(4.5b) \quad \mathcal{O}(|C|^2) : \quad c_* \zeta_1 = L_{0q} \Psi_{0011}^0 - 2F_2(\zeta_+, \zeta_-)$$

$$(4.5c) \quad \mathcal{O}(\epsilon C) : \quad -\frac{i}{q}(d_1 \zeta_+ + d_0 \Psi_{0010}^1) = L_{0q} \Psi_{0010}^1 - F_2(\Psi_{0010}^1, \Psi_{0010}^1)$$

$$(4.5d) \quad \mathcal{O}(AC) : \quad id_2 \zeta_+ + id_0 \Psi_{1010}^0 = L_{0q} \Psi_{1010}^0 - 2F_2(\zeta_0, \zeta_+)$$

$$(4.5e)$$

where we have used the fact that F_2 is a symmetric bilinear form. Equation (4.5b) is decoupled and yields $c_* = 2\Delta_1 \left(\frac{1}{q} + \frac{2b}{3} \right)$ The only coefficient left to determine is d_2 which we shall compute now.

Using $\Psi_{1010}^0 = \langle x_1, x_2, x_3, x_4 \rangle^T$ in (4.5e) implies

$$(4.6a) \quad id_2 + id_0x_1 = x_2$$

$$(4.6b) \quad -d_0d_2 + id_0x_2 = \frac{q}{3}x_1 + x_3$$

$$(4.6c) \quad \frac{2iq}{3}d_2 + id_0x_3 = \frac{q}{3}x_2 + x_4$$

$$(4.6d) \quad -\frac{q}{3}d_0d_2 + id_0x_4 = \frac{q}{3}\left(\frac{q}{3}x_1 + x_3\right) - \frac{2bq\Delta_1}{3}$$

Using (4.6a) in (4.6b) , (4.6b) in (4.6d) and using these in (4.6c) yields $d_2 = \frac{b\Delta_1}{3\sqrt{-q}}$.

Therefore the normal form for (1.1) near C_1 is

$$(4.7a) \quad \frac{dA}{dz} = B$$

$$(4.7b) \quad \frac{dB}{dz} = -\frac{\epsilon}{q}A - \frac{b\Delta_1}{3}A^2 + 2\Delta_1\left(\frac{1}{q} + \frac{2b}{3}\right)|C|^2$$

$$(4.7c) \quad \frac{dC}{dz} = i\sqrt{-q}C - i\frac{\epsilon\sqrt{-q}}{q^3}C + i\frac{b\Delta_1}{3\sqrt{-q}}AC$$

ACKNOWLEDGEMENTS

The authors are extremely grateful to Prof. Gerard Iooss

REFERENCES

- [1] IOOSS G., ADELMAYER, M., *Topics in Bifurcation Theory and Applications*, World Scientific, Singapore, 1998.
- [2] S. ROY CHOUDHURY, *Solitary-wave families of the Ostrovsky equation: An approach via reversible systems theory and normal forms* , Elsevier, (2007), pp. 1468–1479.