



Anomalous Behavior (bad habits) (vices)

By Dr. Hebatallah Elshafaei

Bad habits due to lack of work

Kicking

biting

crib-
biting

Wind
sucking

Weaving

Self-
mutilatio
n



Kicking


Causes:

1. **Lack of work.**
2. Keeping the horse in dark stable
3. Any inflammation in the hoof
4. Mare in estrous also kicking.

Kicking

Means of cure:

1. Remove the cause of kicking if know
2. The following methods may be tried:
 - Anti-kicking boards
 - Bags of sand
 - Anti-kicking log
 - One of the hind legs may be fixed of the stable.
3. It is advisable to keep a kicker in a loose box to avoid disturbance and possible injury.



Training techniques

Desensitization: Gradually introduce "scary" objects (like flags, ropes, or umbrellas) to the horse's hindquarters in a controlled environment. This helps them realize that unusual movements behind them are not threats.

Move the Feet: When a horse shows aggression (like pinning ears), immediately put them to "work" by making them move their feet in a circle or back up. This redirects their focus and associates the bad attitude with effort.

Establish Leadership: Consistent groundwork (like the **Seven Games**) helps a horse see you as a leader, reducing their need to use dominance or defensive kicking to feel safe.

Biting

Causes:

1. Stallion during breeding season and mare in estrous.
2. Lack of work.

Signs:

3. Lying back of the ear.
4. A small sunken eye "Pig eye".
5. Perhaps showing the white of the eye.



Biting:

Means of cure:

- a. Remove the cause
 - b. The following methods
 - side-stick
 - muzzle
 - cradle.
- Tying the horse short to side ring in the stable.




Crib-Biting

The horse grasps a surface with its teeth, arches its neck, and gulps air. It leads to tympany of the stomach and intestines causing indigestion.

Causes : lack of work

Signs:

1. Shows the ribs.
2. Has a staring coat (rough).
3. Pot belly (large abdomen).
4. Worn down incisor teeth.



Crib-Biting:

Prevention:

1. Crib-bitter should be **isolated**.
2. Tying the horse up when there is **no fixed object**
3. **muzzle** when not feeding.
4. **crib biting strap**, which hurts him when he arches his neck.
5. Keeping the horse in a loose-box with
6. By use of **flute-bit (whistling bit)**.

Wind-Sucking:

Wind sucking may be described as crib-biting without the aid of a fixed object.

Remedy:

1. Use the flute bit (whistling bit)
2. Apply muzzle and remove at time of feeding.

Weaving:

A side-to-side swaying of the head and neck, sometimes shifting weight on the feet. It is a **nervous habit when the horse separated from the herd or confined in stall for long time**

Causes: Lack of work

Means of cure:

1. **Tying** the animal up.
2. Giving long hours of easy **regular work**, sufficient but not too much food and a good bed.
3. **Head chains** are changed to head ropes to prevent their continuous noise.
4. a weaver should be kept separated

Self-mutilation

Self-injury through vigorous body friction or flank-biting .

bite at their sides or **rub** their neck crests damaging the coat, the mane, the skin and occasionally causing wounds.

Control:

1. Affected animals can be given **freedom of grazing** since the condition typically occurs in circumstances of confinement and isolation.
2. **Tranquilization** may be necessary to terminate an episode.
3. **Regular exercise** may be given due to lack of work
4. Treatment **of skin disease** .

Bad habits due to nutritional deficiency

01

Eating dung
“ Coprophagia
”

02

Eating soil or
sand
(Geophagia)

03

Chewing and
eating of
wood
(Lignophagia)

04

Eating
bedding

05

Licking the
walls

Eating dung “ Coprophagia”:

Eating dung generally occurs in the **starved animals** or in those suffering from **indigestion** of which may be a symptom. **Foals** sometimes learn the habit when kept in through bad weather.

Remedy:

1. An underfed horse stops the vice if given a **sufficient hay** ration and foals will discontinue it when let out to grass or when they begin to eat solid food.
2. A **muzzle** should be applied.
3. Treat the **indigestion**.
4. **Dung** must be removed three or four times daily or immediately after it is passed.

Eating soil or sand (Geophagia)

Causes:

1. This condition has been thought to be the result of mineral-deficient diets (phosphorus and iron).
2. Close confinement and lack of exercise appear to be the most common causal circumstances.

Dangerous:

Animals with this practice are susceptible to alimentary dysfunction. Excessive eating of sand or dirt can result in **sand impaction** of the caecum and colon.

Eating soil or sand (Geophagia)

Control:

1. Supplementary **ration** of bone meal should be provided to eliminate the possibility of **phosphorus** deficiency.
2. Affected animals can be examined for **anemia and worm infestation** and appropriate treatment provided when indicated.
3. Confined animals should be provided with enforced **exercise**.

Chewing and eating of wood (Lignophagia)

Causes:

1. The condition appears to be associated with a desire for roughage or cellulose.
2. Lack of roughage in the diet predisposes a horse to wood-chewing.
3. Horses fed on concentrate diets with low supply of roughage show the condition much more frequently than horses fed hay in abundance.

Lignophagia

Dangerous:

1. Wood-chewing can lead to **serious intestinal obstruction**, some splinters may be consumed and some can cause **damage within the mouth**.
2. Excessive **tooth wear** also occurs.
3. Affected animals may transmit the habit to associating horses. This can lead to the destruction of wooden fences, partitions and doors

Control:

1. Regular **creosote painting** of the wooden surfaces available to the horse.
2. use of balanced ration

Eating bedding:

Causes:

1. This habit may be due to **insufficiency of bulky food** (hay) or too small ration.
2. An **under worked horse** in a loose-box is more likely to eat his bedding as full work horse tied up at night in stall.

Remedy:

3. Give a full and **correct ration** for the work being done.
4. Use another kind of **bedding** which the horse will not eat
5. Apply a **muzzle** after feeding

Licking the walls

Causes:

1. Shortage of food.
2. Inadequate diet.
3. Digestive disturbances e.g. **internal parasites, mineral insufficiency or too over acidity**

Remedy:

4. Treatment of the digestive disturbance.
5. Parasites must be eliminated.
6. Put a lump of rock salt in the manger or **salt brick** and give the animal balanced mineral supplement containing calcium, phosphorus, magnesium

Throwing the food out of the manger

Some horses prefer to eat from the ground and will therefore throw their food out of the manger with their noses. Others do so to find the grain below the upper layers of chaff.

Remedy

1. By fixing a rope a cross the manager at either end.
2. Put the animal feed on a clean sheet, on the ground,
3. Put the food in a small quantity.
4. Wooden partition of manager.
5. Mixing of food.

Tearing clothes

The horse tears at his rug or quarter sheet.

Prevention:

Put on a muzzle, cradle, side-sticks may be applied

Sleeping while standing:

It is a bad habit rather than vice. The fear to lie down may be the result of:

1. Once have been cast in the stable
2. Overcrowding
3. The presence of rats in the stable.

Dangers:

1. The horse may **fall** causing injury to knees, fetlocks or lips and broken incisors.
2. If a neck rope or neck strap is used, the horse may be **strangled**.
3. The animal put an **extra strain on the muscles**, tendons and ligaments of the limbs, which render them more liable to sprain.

Sleeping while standing:

Remedy:

1. Put the horse in a loose-box with plenty of straw **bedding**.
2. Apply knee caps, fetlock boots, hock boots and **stable bandages**.
3. A strong **rope fixed** between the posts, the animal soon learn to put his weight on this rope.
4. **Slings** of horse to prevent weight on limbs until it accustom to put weight on slings then remove the slings

Vices led to economic losses

Kicking.

**Inter-
sucking
between
cows**

**Inter-
sucking
between
calves**

**Fence
breaking**

Kicking

During milking the cow kicks the milker so the collected milk be drooped on the ground

And economic losses.

Causes:

1. Nervous action as a result of lacking of food.
2. It may be due to any wound on the teats or udder.
3. Some animals are accustomed to be milked in a quit dark place, so if they are milked in a lighted sunny place it will kick or cattle are handled and milked from one side become aggressive when handled from other side which may affect milk production.
4. Rough handling.
5. Change of milker

Kicking

Remedy:

1. Plenty of **good quality food** palatable is put Infront of the animal at the time of milking.
2. Milking should be carried in a **dark quit place** away from any noise or crowdies.
3. **Treat any wound** or cracks on the teat or udder, by lubricating your hand before milking with zinc oxide ointment or Vaseline.
4. Apply hook strap
5. Anti-cow kicking and tail holding device
6. Loop from the tail around the right hind leg.
7. Tying a rope just above fetlock with one end, its other end is tied on a pointed piece of wood fixed in the ground or fixed to a ring.

Inter-sucking between cows

This behavior involves a cow suckling milk from its udder or the udder of other cow. Frequent sucking can lead to lose of milk, teat damage, pathological changes and deformation of the udder.

Inter-sucking between cows

Causes:

Cow suffers from **injuries in teat lick** its teat leading to sucking of milk so economic losses.

Means of cure:

1. The animal should be separated from others.
2. **Anti-sucking bit** is applied on nasal bridge and it must be removed at the time of feeding
3. Apply a muzzle, side stick or cradle.
4. Make partitions between the animals.
5. Apply **Anti-sucking shell** over the udder.

Fence breaking

Some animals have the habit of breaking the fences of their enclosures.

Causes:


1. Feeling that on the other side of the fence the **grasses** are more green or plenty.
2. Some bulls behave similar behavior when show **females in estrus** outside the fence.

Prevention:

1. Good arrangement.
2. Apply a **rubber** bag around the tip of the horn.
3. **Dehorning**.

Inter-sucking between calves

Calves separated from their mothers **suck and lick at their own bodies**, at objects in their pens and at parts of the bodies of other calves. They commonly suck on the navel, prepuce, scrotum, udder and ears of other animals.



Inter-sucking between calves

Causes:

1. **Wide** aperture of the nipple in case of artificial feeding.
2. Shorting the **time** of milking.
3. In case of **bucket feeding**, the calf sucks milk around the mouth of other calf which is called kissing.

Dangerous:

4. If the **coat** of the other calf is sucked significant quantities of hair may be ingested leading to formation of balls, or **bezoar** in the rumen which leads to digestive problems and even death.
5. If the penis is sucked, **urine** is often produced and then drunk by the suck calf. This can lead to liver disorders and to reduced nutrient intake.
6. Another adverse consequence of inter sucking is that the part of the calf sucked may become **inflamed**, damaged and infected.

Intersucking between calves

Control:

1. The feeding of calves with milk through **automatic nursers** with teats whose aperture are narrow, is such that the sucking times of each feed period is sufficiently prolonged will reduce the likelihood that other objects or calves will be sucked.
2. Young calves should be **housed individually** and tied shortly.
3. The supplement of **roughage**, such as 150 gm. of a supply of straw, can result in a reduction in inter sucking.
4. Urine drinking can be reduced by the plentiful **supply of water** from a normal drinker for the calves.
5. Apply **muzzle** and removed at the time of feeding.

Tongue-rolling

The tongue is extruded from the mouth and moved **by curling and uncurling outside or inside the mouth with no solid material present**. The tongue is extruded and rolled back into the open mouth, after which partial swallowing of the tongue and gulping of the air takes place

Tongue-rolling

Causes:

1. The condition occurs in all ages and breeds but younger cattle and certain breeds such as Brown Swiss are thought to exhibit it most frequently.
2. It is believed that there may be a **hereditary factor** in this condition but it is also believed that nutritional deficiencies can precipitate it.
3. It occurs more often in cows within herds which are continuously **confined indoors**.
4. When a feed is provided in a form **low in roughage** and not requiring significant oral activity in prehension and mastication, the condition may be encouraged.

Tongue-rolling

Control:

1. Insertion of a **metal ring** through the frenulum of the tongue.
2. The provision of diets improved by **salt mixtures**.
3. The provision of **freedom of movement**.
4. **Isolation** of the affected animals.
5. Apply **muzzle**.

Cannibalism (aggressive pecking)

Causes:

1. Nutritional disorders
2. **Crowding.**
3. Prolepses of oviduct.
4. **Excessive light.**
5. Poor ventilation.
6. **High temperature.**
7. Low humidity.
8. External parasites.
9. Genetic factors
10. Too little feeding and watering spaces.
11. increased number of males.



Prevention and control

-
1. **beak-trimming (debeaking)** . Beak-trimming involves the removal of the anterior part of the upper mandible.
 2. Limit the vision of birds. the use of red lamps or painting window panes red so it is difficult for birds to identify bloody sores that encourage more picking.
 3. Providing adequate feed and water space and protein supplement.
 4. Overcrowding should be avoided.
 5. Careful attention to ventilation as good ventilation lower heat and remove foul air lessening stress on birds and light intensity neither too bright nor too long.
 6. The scattering of pelted feed on the floor.

Egg-eating

1. The egg may be abnormal.
2. Soft shelled -egg.
3. Deficiency of Ca and P in the ration.
4. Inadequate nesting facilities.
5. Failure to collect eggs frequently.

Egg-eating

Prevention and control:

1. Beak-trimming.
2. Frequent collection of eggs.
3. Filling empty eggshells with unpleasant tasting substance.

Broodiness

hen would like to sit on some of her eggs to hatch allowing her body temperature to increase and often consuming less food and water than she would normally.

Causes:

1. Genetic factors.
2. The increase of prolactin hormone.
3. Accumulation of eggs in the nests encouraged the chicken to broodiness.

Broodiness

Control:

1. Placing the birds in small cages with a wire or slated floor (**Broody-coop**).
2. By bathing the bird in cold water.
3. Frequent collection of eggs.

Floor eggs

Causes:

1. Inadequate number of nests.
2. Insufficient nesting material in the nest.
3. Presence of broody hens.

Treatment:

1. Adequate number of nests should be provided.
2. Keeping the nesting material clear and ample.
3. Placing the nests in the pen before the birds started laying by 2weeks.
4. Placing the nests in a darker part of the house.

Destructiveness

Causes:

1. Absence of the owner from the home.
2. Lack of exercise.
3. Barrier frustration
4. Breed incidence: there may be a breed incidence factor as well Ex. **beagles; German shepherds** seem to be particularly prone to destructiveness.



Destructiveness

Treatment:

1. Negative reinforcement
2. Companionship
3. It is important that the dog get enough exercise.
4. Progesterone therapy for a short period (10-14 days) to calm the dog.
5. Behavioral modification

excessive Barking

Some dogs tied up outside, bark because of discomfort seek attention, or express boredom and separation anxiety.

Remedy

by shock collars that administer a shock whenever the animal barks have been used with some success but the severity of the shock is variable.

phobias


Fear of the loud noise of thunder, other dogs , child, moving truck or bicycle this called Trigger

Treatment: The best treatment for phobias is

systematic desensitization: These techniques work by **gradually exposing** the dog to a trigger at a low enough intensity that it remains calm (distance, volume or speed)

counterconditioning : This involves creating a **positive association** by pairing the trigger with a high-value reward

Systematic Desensitization with your dog



Counterconditioning with your dog




=



**Stress and
Fear**



+



=



**Repeat Many
Times**



=



**Relaxed or
Happy**

Spraying of urine

1. why do cats spray - to **mark their territory**

2. figure out if spraying vs urination

spraying is ejecting squirts of urine against vertical surfaces (like the couch) and

urinating involves squatting and urinating on horizontal surfaces

3. make sure no underlying **medical problem** - cystitis, stones, bladder tumor, arthritis, diabetes, kidney disease - these can either cause an increase in urine production or make it hard for your cat to use their litter box

4. **Castration** eliminated spraying behavior in 87% of the tom cats

Spraying of urine

reduce their stress

litter tray

A close-up photograph of numerous lavender flower spikes, showing individual small purple blossoms and green buds. The flowers are densely packed and fill the entire frame, creating a textured, purple and green background.

Thank you
