

# Gravitational Collapse

Tayyaba Noureen

SBASSE LUMS

Phy 442 - General Relativity, April 25, 2024

# Table of Contents

- 1 Idealizations
- 2 Exterior Geometry
- 3 Interior Geometry
- 4 Surface of the Star
- 5 Conclusions

- Star is static, spherically symmetric and non radiating

- Star is static, spherically symmetric and non radiating
- Vacuum outside star

- Star is static, spherically symmetric and non radiating
- Vacuum outside star
- Interior of star is a perfect pressureless fluid called dust

# Exterior Geometry

- Most general spherically symmetric vacuum solution of Einstein Equations is Schwarzschild metric - Birkhoff Theorem

# Exterior Geometry


- Most general spherically symmetric vacuum solution of Einstein Equations is Schwarzschild metric - Birkhoff Theorem
- Metric:

$$ds^2 = - \left( 1 - \frac{2M}{R} \right) dt^2 + \frac{1}{- \left( 1 - \frac{2M}{R} \right)} dr^2 + r^2 d\Omega^2$$

- Surface of star freely falls radially

$$r = \frac{R_i}{2}(1 + \cos \eta) \quad \tau = \frac{R_i}{2} \left( \frac{R}{2M} \right)^{1/2} (\eta + \sin \eta)$$


$$t = 2M \ln \left[ \frac{(R_i/2M - 1)^{1/2} + \tan \eta/2}{(R_i/2M - 1)^{1/2} - \tan \eta/2} \right] \\ + 2M(R_i/2M - 1)^{1/2}(\eta + (R_i/4M)(\eta + \sin \eta))$$





- Time elapsed for comoving observer:


$$\Delta\tau = \pi \sqrt{\frac{R_i^3}{8M}}.$$



- Time elapsed for comoving observer:

$$\Delta\tau = \pi\sqrt{\frac{R_i^3}{8M}}.$$

- Conclusion:
  - Event Horizon ( $R = 2M$ )
  - No communication beyond  $R = 2M$
  - External observer never sees the star crossing horizon



# Interior Geometry

- Interior is homogeneous and isotropic

# Interior Geometry

- Interior is homogeneous and isotropic
- Friedmann Solution

$$ds^2 = -d\tau^2 + a^2(\tau)[d\chi^2 + \sin^2 \chi(d\theta^2 + \sin^2 \theta d\phi^2)].$$

$$a = \frac{a_m}{2}(1 + \cos \eta)$$

$$\tau = \frac{a_m}{2}(\eta + \sin \eta)$$

# Interior Geometry

- Interior is homogeneous and isotropic
- Friedmann Solution

$$ds^2 = -d\tau^2 + a^2(\tau)[d\chi^2 + \sin^2 \chi(d\theta^2 + \sin^2 \theta d\phi^2)].$$

$$a = \frac{a_m}{2}(1 + \cos \eta)$$

$$\tau = \frac{a_m}{2}(\eta + \sin \eta)$$

- Energy Density

$$\rho = \frac{3a_m}{8\pi a^3} = \frac{3}{8\pi a_m^2} \left( \frac{1}{2}(1 + \cos \eta) \right)^{-3}$$

# Surface of Star

## First Junction Condition

- Induced metric on both side must be the same

$$h_{ab} = g_{\alpha\beta} e_a^\alpha e_b^\beta = g_{\alpha\beta} \frac{\partial x^\alpha}{\partial y^a} \frac{\partial x^\beta}{\partial y^b}$$

$$ds_\Sigma^2 = -d\tau^2 + a^2(\tau) \sin^2 \chi_0 d\Omega^2.$$

$$ds_\Sigma^2 = -(F \dot{T}^2 - F^{-1} \dot{R}^2) d\tau^2 + R^2(\tau) d\Omega^2$$

# Surface of Star

## First Junction Condition

- Induced metric on both side must be the same

$$h_{ab} = g_{\alpha\beta} e_a^\alpha e_b^\beta = g_{\alpha\beta} \frac{\partial x^\alpha}{\partial y^a} \frac{\partial x^\beta}{\partial y^b}$$

$$ds_\Sigma^2 = -d\tau^2 + a^2(\tau) \sin^2 \chi_0 d\Omega^2.$$

$$ds_\Sigma^2 = -(F\dot{T}^2 - F^{-1}\dot{R}^2)d\tau^2 + R^2(\tau)d\Omega^2$$

- Comparing the induced metrics we get:

$$R(\tau) = a(\tau) \sin \chi_0$$

$$F\dot{T}^2 - F^{-1}\dot{R}^2 = 1 \implies F\dot{T} = \sqrt{\dot{R}^2 + F} = \beta(R, \dot{R}).$$

## Second Junction Condition

- Extrinsic curvature on both sides must be the same

$$K_{ab} = \nabla_{\beta}(n_{\alpha})e_a^{\alpha}e_b^{\beta}$$



## Second Junction Condition

- Extrinsic curvature on both sides must be the same

$$K_{ab} = \nabla_\beta (n_\alpha) e_a^\alpha e_b^\beta$$

- Comparing the curvature tensor components we get:

$$\beta = E = \cos \chi_0$$

$$\dot{\beta} = 0$$

## Second Junction Condition

- Extrinsic curvature on both sides must be the same

$$K_{ab} = \nabla_\beta (n_\alpha) e_a^\alpha e_b^\beta$$

- Comparing the curvature tensor components we get:

$$\beta = E = \cos \chi_0$$

$$\dot{\beta} = 0$$

- Implies energy conservation

## Second Junction Condition

- Extrinsic curvature on both sides must be the same

$$K_{ab} = \nabla_{\beta}(n_{\alpha})e_a^{\alpha}e_b^{\beta}$$

- Comparing the curvature tensor components we get:

$$\beta = E = \cos \chi_0$$

$$\dot{\beta} = 0$$

- Implies energy conservation
- For smooth transition, hypersurface must be generated by geodesics of interior and exterior geometries.

# Conclusions

- General features of gravitational collapse are the same

# Conclusions

- General features of gravitational collapse are the same
- Horizon and Singularity theorems