Square-Wave Illusion

Physical simulation and inverse rendering

Design and Analysis of Experiments | Group 2
For teammate and professor review. Provisional target; no human data.

What has been completed?

All six computational steps in the supplied project outline are implemented: periodic surface geometry, light visibility and self-shadowing, spherical integration, Lambertian shading, von Mises-Fisher lighting, and joint optimization of surface heights and lighting. The repository includes source code, an executed notebook, saved results, numerical checks, and reproduction instructions.

Current finding


A bounded periodic height field can approximate the provisional triangle-wave target under the stated direct-light model. The free-surface candidate has RMSE **0.04155** at N=16. An additional N=32 search gives **0.02110**. Target mean luminance is one. These values describe image matching, not observer performance.

Model	Segments	Dense RMSE	R-squared
Flat control	16	0.34640	-0.0000
Sinusoid	16	0.05928	0.9707
Square-like	16	0.14801	0.8174
Free surface	16	0.04155	0.9856
Free: resolution check	32	0.02110	0.9963

Dense metrics use 16,384 light directions and 15 samples per segment. The N=32 result is a one-seed sensitivity case, not part of the three-seed N=16 comparison. R-squared is descriptive.

What is not established?

The fitted geometry is not constrained to be rectangular. A low residual does not show that it is the surface people perceive, identify a unique shape/light solution, prove global optimality, or establish a psychological explanation. Final interpretation requires agreement on the exact stimulus and admissible model.




Additional N=32 fit. The Z axis is exaggerated and the grayscale is illustrative.

1. Physical model

Geometry. One period contains N equally spaced height vertices, joined by planar segments with periodic closure. The cross-section repeats along X and extends along Y. An orthographic observer looks down Z. Surfaces are opaque and single-valued; vertical walls and overhangs are excluded.

Visibility. A light vector points toward its source. A point receives direct light only if the ray clears the maximum surface-elevation slope on that side. Checking vertices within one full period on each side includes neighboring cycles exactly for this geometry. Several samples are needed because a segment can be partly shadowed.



Example above: a low light source illuminates yellow points; dark points are blocked by the surface. The second plot multiplies binary visibility by the Lambertian cosine factor.


Lighting. Approximately equal-solid-angle Fibonacci points avoid uniform-latitude pole bias. A full-sphere von Mises-Fisher radiance profile has mean direction mu and concentration kappa. Zero concentration is uniform; larger values make the light more directional. Total angular radiance is fixed.

```
L = (rho/pi) sum [R(light) V max(0, normal.light) 4pi/M]
```

Constant albedo rho=1; output is relative linear luminance. No interreflection, specular reflection, texture, or calibrated display response is included. A flat white surface under uniform unit radiance has output one. See docs/METHODS.md for the full definitions and proof of the horizon construction.

2. Fitted candidate models

Each model is fitted to the same provisional triangle wave (period one, mean one, Michelson contrast 0.6). No free brightness gain or offset is used. Geometry appears on the left; target and rendered luminance appear on the right. These are synthetic modeling outputs.




The selected free candidate has maximum absolute height 0.03365 periods and maximum absolute slope 0.24840. Its light has azimuth 140.70 degrees, elevation 14.56 degrees, and kappa 21.628. These are one candidate set of parameters, not a unique reconstruction.

3. Optimization and numerical checks

The objective combines mean squared luminance error with a curvature penalty and penalties for height/slope bound violations. Mean height is fixed to zero. The N=16 free model has N-1 height coordinates plus three lighting coordinates; this removes the redundant vertical offset from the nominal N+3 outline.

Reference setting	Value
Height and slope limits	Absolute height <=0.25 periods; absolute slope <=6
Light bounds	Azimuth 0-180 degrees; elevation 5-85 degrees; kappa 0-30
Reference search	1,024 directions; 3 samples/segment; seeds 11, 29, 47
Solver	Differential evolution; bounded Powell refinement
Regularization	Mean squared periodic curvature; weight 0.00001
Refinement and evaluation	Refine at 2,048 directions and 5 samples; evaluate at 16,384 and 15

All free-surface global searches reached their iteration limit. Selected final local refinements terminated successfully; that stopping condition is not a proof of a global minimum. Free searches share the best fixed-shape initialization. Other initializations may yield different solutions.



Midpoint trap: the same N=16 candidate gives RMSE 0.00215 at segment midpoints, but 0.04155 across 15 samples per segment (0.04162 at 31). A midpoint-only claim would substantially overstate the fit. Increasing light directions from 16,384 to 32,768 changes the error from 0.04155 to 0.04154.

N=8 and N=32, changed smoothness, and a kappa upper bound of 60 were tested with one additional warm-started seed each. These are diagnostics, not exhaustive searches or inferential comparisons. Geometry resolution remains influential.

4. Validation and reproduction

15 of 15 physical/numerical checks passed. These include sphere-unit vectors, light normalization, the flat-surface analytic result, overhead illumination, the cosine law, lower-hemisphere blocking, nonnegativity, and invariance to height translation and light reflection along the ridges.

The shadow algorithm matches an independent explicit ray/segment intersection calculation at 576 sample/direction pairs containing both lit and shadowed points. Eight regression tests additionally check periodic normals, uniform radiance on tilted unoccluded facets, parameter encoding, invalid inputs, bounds, saved result consistency, and recovery of known synthetic kappa. All eight pass.

Open this	Purpose
README.md	Status, findings, links, and quick start
simulation_walkthrough.ipynb	Executed explanation with embedded plots and results
docs/METHODS.md	Geometry, light conventions, constraints, and objective
docs/RESULTS.md	Metrics, interpretation, and numerical sensitivity
docs/PROFESSOR_REVIEW.md	Decisions required before final interpretation
results/optimization/fits.json	All 12 reference searches and solver statuses
results/optimization/analysis.json	Dense evaluation, sensitivity, and convergence
results/optimization/*.csv	Numerical comparisons and sampled luminance profiles
configs/reference.json	Editable target and search configuration

Quick checks, from the repository root:

```
python validate.py
python -m unittest -v test_project.py
```

Full computational reproduction:

```
python run_all.py --full
```

The full pipeline regenerates demonstrations, fitting, analysis, summaries, the executed notebook, and tests. Generated outputs are replaced. The PDF is rebuilt separately with build_report.py and the optional requirements-report.txt dependencies. Saved reference versions are in requirements-reference.txt. Single-thread BLAS settings and fixed seeds improve reproducibility, but exact timings and final digits may vary across environments.

5. What remains before submission?

The computational prototype is ready for group review. It is not automatically the final course project: the question being tested must be agreed with the professor. No deadline, final author order, or behavioral-study requirement has been invented.

Confirm the target

Ideal triangle wave, specified harmonics, or measured luminance? Agree on phase, mean, contrast, and period.

Define the admissible shape

Can any bounded periodic graph count, or must the fit reproduce the particular reported square-like percept?

Review light and geometry assumptions

Approve the single vMF lobe, fixed angular radiance, source bounds, height/slope limits, and direct-only illumination.

Set the success criterion

Choose an acceptable whole-profile error and contrast tolerance. Then set final N, compute budget, and any further search/physics checks.

Finalize the course deliverables

Review group contributions, report length, presentation requirements, and how to discuss the collaborators through the instructor.

Literature context

Leguire, L. E., Blake, R., and Sloane, M. E. (1981). A Novel Illusion of Bars Made from Triangles. *Science, 212*, 1172-1175.

Wildzunas, R. M., and May, J. G. (1992). Phase shifts and the square-wave illusion. *Spatial Vision, 6*, 209-220.

May, J. G. (1994). Stimulus manipulations that reduce the square-wave illusion. *Spatial Vision, 8*, 329-340.

These papers motivate the question; their behavioral experiments are not reproduced here. Additional technical references and links are in docs/REFERENCES.md. Paper PDFs, private email text, and unpublished collaborator material are not redistributed.

Sharing the work

Upload the extracted project folder as the GitHub repository root. Start with the README, this PDF, and the executed notebook for teammate review; then revise before professor review. Follow docs/GITHUB_GUIDE.md. No remote repository has been created and no collaborator has been contacted by this package.