

# Optimal control in octopus: status report

Alberto Castro

ARAID Foundation and Institute for Biocomputation and Physics of Complex Systems (BIFI), Zaragoza (Spain)

octopus meeting, Jena, 26-30.01.2015



Instituto Universitario de Investigación  
**Biocomputación y Física  
de Sistemas Complejos**  
Universidad Zaragoza



# Outline

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary

# Outline

## What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

## What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

## Summary

## What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

## Control of quantum phenomena: past, present and future

**Constantin Brif, Raj Chakrabarti<sup>1</sup> and Herschel Rabitz<sup>2</sup>**

Department of Chemistry, Princeton University, Princeton, NJ 08544, USA

E-mail: [cbrif@princeton.edu](mailto:cbrif@princeton.edu), [rchakra@purdue.edu](mailto:rchakra@purdue.edu) and [hrabitz@princeton.edu](mailto:hrabitz@princeton.edu)

*New Journal of Physics* **12** (2010) 075008 (68pp)

# Basic idea, in a cartoon

## What is OCT?

What OCT calculations can we do with octopus now?


Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary



[R. J. Levis *et al*, Science **292**, 709 (2001)]

# Mono-chromatic lasers: IVR

## What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems


Different algorithms

Different targets

Different parametrizations

## Summary

- ▶ Lasers (coherent, monochromatic, intense light) promised to deliver precise control of quantum systems
- ▶ Initially, the first attempts to control molecules (i.e. “photo-selective chemistry”) were based on tuning the laser frequency to specific bonds
- ▶ Those attempts were seldom successful, due to “intramolecular vibrational redistribution”.



- ▶ Analogous problems will appear in other quantum control attempts, beyond molecular photo-chemistry.

# Pioneering schemes


## What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems  
Different algorithms  
Different targets  
Different parametrizations

## Summary


- **Interferences, and the “two pathway” scheme** Use of two monochromatic lasers with commensurate frequencies for creating quantum interference between two reaction pathways [P. Brumer and M. Shapiro, Chem. Phys. Lett. **126**, 541 (1986)].



- **Pump and dump** [D. J. Tannor and S. A. Rice J. Chem. Phys. **83**, 5013 (1985)]

Limitations: knowledge of the potential energy surfaces, competing processes.

- **STIRAP: STImulated Raman scattering involving Adiabatic Passage** [K. Bergmann *et al* Chem. Phys. Lett. **149**, 463 (1988)]



# Adaptive feedback control

## What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

## Summary

VOLUME 68, NUMBER 10

PHYSICAL REVIEW LETTERS

9 MARCH 1992

## Teaching Lasers to Control Molecules

Richard S. Judson<sup>(a)</sup>*Center for Computational Engineering, Sandia National Laboratories, Livermore, California 94551-0969*

Herschel Rabitz

*Department of Chemistry, Princeton University, Princeton, New Jersey 08544*

(Received 26 August 1991)

We simulate a method to teach a laser pulse sequences to excite specified molecular states. We use a learning procedure to direct the production of pulses based on "fitness" information provided by a laboratory measurement device. Over a series of pulses the algorithm learns an optimal sequence. The experimental apparatus, which consists of a laser, a sample of molecules, and a measurement device, acts as an analog computer that solves Schrödinger's equation *exactly*, in real time. We simulate an apparatus that learns to excite specified rotational states in a diatomic molecule.




# The learning loop

## What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems  
Different algorithms  
Different targets  
Different parametrizations

Summary



[H. Rabitz *et al*, Science **288**, 824 (2000)]

# Examples of AFC experiments: other

## What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

- ▶ Photo-dissociation reactions in molecules.
- ▶ High harmonic generation.
- ▶ Multi-photon ionization of atoms.
- ▶ Electronic excitation in molecules (fluorescence is used as the probe to build the merit function).
- ▶ Molecular alignment.
- ▶ Photo-induced electron transfer between molecules
- ▶ Photo-isomerization of molecules.
- ▶ etc.

# What optimal control theory is, in a nutshell.

## What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems  
Different  
algorithms  
Different targets  
Different  
parametrizations

Summary

Typical formulation of a (general) optimal control problem:

- Dynamical system:

$$\begin{aligned}\dot{x}(t) &= f(x(t), u(t), t) \\ x(0) &= x_0\end{aligned}$$

Typically,  $u = u(t)$ . But it can be a set of parameters whatsoever.

- Minimize the cost functional:

$$F[x, u] = F^{\text{terminal}}[x(T), u] + \int_0^T dt L(x(t), u(t))$$

- Since  $u \rightarrow x[u]$ , it amounts to minimizing

$$G[u] = F[x[u], u]$$

# What optimal control theory is, in a nutshell.

## What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

Problems to be answered by OCT:

- ▶ Design a good algorithm to find the optimal  $u$ .  
For example, find an efficient way to compute  $\nabla G[u]$ .
- ▶ How many solutions are there? Are there local minima?
- ▶ How stable are the minima with respect to small changes in  $u$ ? (find  $\nabla^2 G[u]$ )
- ▶ How stable are the minima with respect to small changes in the problem definition (changes in  $f$  or  $x_0$ )?
- ▶ Is the problem *controllable* at all?

# Essential theoretical results

## What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

- ▶ Pontryagin's minimum principle (1956)  
[V.G. Boltyanskii, R.V. Gamkrelidze, and L.S. Pontryagin, "Towards a theory of optimal processes", (Russian), Reports Acad. Sci. USSR **110**, 1 (1956)]  
It provides a *necessary* condition for the minimum – in practice, typically, an expression for  $\nabla G[u]$  so that the equation  $\nabla G[u] = 0$  can be posed.
- ▶ Hamilton-Jacobi-Bellman equation (1954)  
(Theory of "dynamic programming", Richard Bellman)  
[R.E Bellman, "Dynamic Programming and a new formalism in the calculus of variations" Proc. Nat. Acad. Sci. **40**, 231 (1954)]

# Hamiltonian systems, calculation of the gradient

## What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary

$$\begin{aligned}\dot{q}_i(t) &= \frac{\partial H}{\partial p_i}(q(t), p(t), u, t), \\ \dot{p}_i(t) &= -\frac{\partial H}{\partial q_i}(q(t), p(t), u, t).\end{aligned}$$

In condensed vector notation:

$$\begin{aligned}\dot{\mathbf{q}} &= \frac{\partial H}{\partial \mathbf{p}}, \\ \dot{\mathbf{p}} &= -\frac{\partial H}{\partial \mathbf{q}}.\end{aligned}$$

In this case,  $x$  is the vector  $[\mathbf{q}, \mathbf{p}]^\dagger$ .

The cost (or target) functional can be, for example, any expression in the form:

$$F[\mathbf{q}, \mathbf{p}, u] = J_1[\mathbf{q}(T), \mathbf{p}(T)] + J_2[u].$$

# Hamiltonian systems, calculation of the gradient

## What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary

$$\begin{aligned} \frac{\partial G}{\partial u_m} &= \frac{\partial J_2}{\partial u_m} \\ &\quad - \int_0^T dt \, \tilde{\mathbf{p}}(t) \cdot \frac{\partial}{\partial u_m} \frac{\partial H}{\partial \mathbf{p}}(\mathbf{q}(t), \mathbf{p}(t), u, t) \\ &\quad + \int_0^T dt \, \tilde{\mathbf{q}}(t) \cdot \frac{\partial}{\partial u_m} \frac{\partial H}{\partial \mathbf{q}}(\mathbf{q}(t), \mathbf{p}(t), u, t) \end{aligned}$$

The “costate”  $[\tilde{\mathbf{q}}, \tilde{\mathbf{p}}]^\dagger$  is itself a Hamiltonian system, determined by the quadratic Hamiltonian:

$$\begin{aligned} \tilde{H}(\tilde{q}, \tilde{p}, q, p, u, t) = \\ \frac{1}{2} \tilde{\mathbf{q}}^t \mathbf{H}^{qq}(q, p, u, t) \tilde{\mathbf{q}} + \frac{1}{2} \tilde{\mathbf{q}}^t \mathbf{H}^{qp}(q, p, u, t) \tilde{\mathbf{p}} + \\ + \frac{1}{2} \tilde{\mathbf{q}}^t \mathbf{H}^{pq}(q, p, u, t) \tilde{\mathbf{p}} + \frac{1}{2} \tilde{\mathbf{q}}^t \mathbf{H}^{pp}(q, p, u, t) \tilde{\mathbf{p}}. \end{aligned}$$

# Hamiltonian systems, calculation of the gradient

## What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary

The matrices  $\mathbf{H}^{xy}$  are defined as:

$$H_{ij}^{qq} = \frac{\partial^2 H}{\partial q_i \partial q_j}(q, p, u, t)$$

$$H_{ij}^{qp} = \frac{\partial^2 H}{\partial q_i \partial p_j}(q, p, u, t)$$

$$H_{ij}^{pq} = \frac{\partial^2 H}{\partial p_i \partial q_j}(q, p, u, t)$$

$$H_{ij}^{pp} = \frac{\partial^2 H}{\partial p_i \partial p_j}(q, p, u, t)$$



# Quantum problems, calculation of the gradient

## What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

$$\hat{H} = \hat{H}[u_1, \dots, u_M; t]$$

$$\begin{aligned} i \frac{d}{dt} |\Psi(t)\rangle &= \hat{H}[u; t] |\Psi(t)\rangle \\ |\Psi(t_0)\rangle &= |\Psi_0\rangle \end{aligned}$$

$$\Psi(t_0) \longrightarrow \Psi[u](t) \longrightarrow \Psi[u](T)$$

Maximize a quantity

$$F = F[\Psi[u](t)],$$

that depends on the system evolution, or final state, or both.

# Quantum problems, calculation of the gradient

## What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

## Summary

$$F[\Psi, u] = J_1[\Psi(T)] + J_2[u]$$

$$G[u] = F[\Psi[u], u]$$

$$\frac{\partial G}{\partial u_m} = \frac{\partial J_2}{\partial u_m} + 2\text{Im} \int_0^T dt \langle \chi(t) | \frac{\partial \hat{H}}{\partial u_m} | \Psi(t) \rangle ,$$

where the “costate”  $\chi$  verifies:

$$i \frac{d}{dt} |\chi(t)\rangle = \hat{H}(t) |\chi(t)\rangle ,$$

$$|\chi(T)\rangle = \frac{\delta}{\delta \Psi^*(T)} F[\Psi(T)]$$

# Outline

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary

# Outline

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary

## What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems  
Different  
algorithms  
Different targets  
Different  
parametrizations

## Summary

# Different dynamical systems: (1) "Standard" Schrödinger equation.

Example: [Zhu *et al.*, J. Chem. Phys. **108**, 1953 (1999)]  
(06-zbr98.test)

Transition between the ground and one excited vibrational state in a Morse potential:

$$V(r) = D_0[e^{-\beta(r-r_0)} - 1]^2 - D_0$$

$$H(t) = T_0 + V(r) - \mu_0 r e^{-r/r^*} \epsilon(t)$$




Fig. 1. Potential energy curves for nitrogen. Energy in wave-number and nuclear separation in Angstrom units. Numbers on curves refer to Table I.

[Morse, Phys. Rev. B **34**, 57 (1929)]

## What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems  
Different  
algorithms  
Different targets  
Different  
parametrizations

## Summary

# Different dynamical systems: (1) "Standard" Schrödinger equation.




FIG. 1. Optimized objective functional  $J_{\beta}^{(N)}$  and transition probability  $|P_{\beta}^{(N)}|^2$  for the transition of  $v=0 \rightarrow v=1$ . Rapid monotonic convergence is found.




FIG. 2. Optimized electric field time dependence for the transition of  $v=0 \rightarrow v=1$ . It is almost a pure cosine function.

# Different dynamical systems: (2) TDDFT.

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

- ▶ We have a system of  $N$  electrons, driven by an external potential  $v_{\text{ext}}(\vec{r}, t, \mathbf{u})$ .
- ▶ The time-dependent density is therefore determined by  $\mathbf{u}$ :

$$\mathbf{u} \longrightarrow n[\mathbf{u}](\vec{r}, t) = \langle \Psi[\mathbf{u}](t) | \hat{n}(\vec{r}) | \Psi[\mathbf{u}](t) \rangle$$

- ▶ The objective is to maximize some function  $G$  of the *control parameters*  $\mathbf{u}$ , defined in terms of a functional of the density:

$$G[\mathbf{u}] = \tilde{F}[n[\mathbf{u}], \mathbf{u}].$$

- ▶ Since the definition is given in terms of the density, everything can be reformulated for the Kohn-Sham system, and the optimization will be equivalent. Since we use the Kohn-Sham substitution, we may use the Kohn-Sham orbitals instead:

$$F[\varphi[\mathbf{u}], \mathbf{u}] \equiv \tilde{F}[n[\mathbf{u}], \mathbf{u}], \quad n[\mathbf{u}](\vec{r}, t) = \sum |\varphi_i[\mathbf{u}](\vec{r}, t)|^2.$$

# Different dynamical systems: (2) TDDFT.

## What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary

- Definition of a target in terms of the Kohn-Sham orbitals:

$$F = F[\underline{\varphi}, \underline{u}] \quad \Rightarrow \quad G[\underline{u}] = F[\underline{\varphi}[\underline{u}], \underline{u}] = \tilde{F}[n[\underline{u}], \underline{u}].$$

- Optimal control theory equations for TDDFT (terminal target only):

$$\begin{aligned} \nabla_u G[\underline{u}] &= \nabla_u F[\underline{\varphi}[\underline{u}], \underline{u}] + \\ &2\text{Im} \left[ \sum_{i=1}^N \int_0^T dt \langle \lambda_i[\underline{u}](t) | \nabla_u \hat{H}[n[\underline{u}](t), u, t] | \underline{\varphi}_i[\underline{u}](t) \rangle \right] \end{aligned}$$

$$\dot{\underline{\varphi}}[\underline{u}](t) = -i \underline{\hat{H}}[n(t), u, t] \underline{\varphi}[\underline{u}](t),$$

$$\underline{\varphi}_u(0) = \underline{\varphi}_0,$$

$$\dot{\underline{\lambda}}[\underline{u}](t) = -i \left[ \underline{\hat{H}}[n(t), u, t] + \underline{\hat{K}}[\underline{\varphi}[\underline{u}](t)] \right] \underline{\lambda}[\underline{u}](t),$$

$$\underline{\lambda}[\underline{u}](T) = \frac{\delta F}{\delta \underline{\varphi}^*} [\underline{\varphi}[\underline{u}](T), \underline{u}].$$



# Different dynamical systems: (2) TDDFT.

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

$$\dot{\underline{\lambda}}[u](t) = -i \left[ \underline{\hat{H}}^\dagger[n[u](t), u, t] + \underline{\hat{K}}[\underline{\varphi}[u](t)] \right] \underline{\lambda}[u](t),$$

$$\dot{\lambda}_i[u](t) = -i \hat{H}^\dagger[n[u](t), u, t] \lambda_i[u](t) - i \sum_{j=1}^N \hat{K}_{ij}[\underline{\varphi}[u](t)] \lambda_j[u](t)$$

$$\begin{aligned} & \langle \vec{r} | \hat{K}_{ij}[\underline{\varphi}[u](t)] | \lambda_j[u](t) \rangle = \\ & -2i\varphi_i[u](\vec{r}, t) \text{Im} \left[ \int d^3r' \lambda_j[u]^*(\vec{r}', t) f_{\text{Hxc}}[n[u](t)](\vec{r}, \vec{r}') \varphi_j[u](\vec{r}', t) \right] \end{aligned}$$

$$f_{\text{Hxc}}[n[u](t)](\vec{r}, \vec{r}') = \frac{1}{|\vec{r} - \vec{r}'|} + f_{\text{xc}}[n[u](t)](\vec{r}, \vec{r}')$$

# Different dynamical systems: (3) Ehrenfest dynamics, with TDDFT (or not).

Coupled electron-ion model: Ehrenfest dynamics:

$$\hat{H}[q, p, u, t] = H_{\text{clas}}[q, p, u, t] \hat{I} + \hat{H}_{\text{quantum}}[q, p, u, t].$$

$$\begin{aligned} \dot{q}_a(t) = & \frac{\partial H_{\text{clas}}}{\partial p_a}[q(t), p(t), u, t] \\ & + \langle \Psi(t) | \frac{\partial \hat{H}_{\text{quantum}}}{\partial p_a}[q(t), p(t), u, t] | \Psi(t) \rangle \end{aligned}$$

$$\begin{aligned} \dot{p}_a(t) = & -\frac{\partial \hat{H}_{\text{clas}}}{\partial q_a}[q(t), p(t), u, t] \\ & - \langle \Psi(t) | \frac{\partial \hat{H}_{\text{quantum}}}{\partial q_a}[q(t), p(t), u, t] | \Psi(t) \rangle \end{aligned}$$

$$\dot{\Psi}(x, t) = -i \hat{H}_{\text{quantum}}[q(t), p(t), u, t] \Psi(x, t),$$

[Castro and Gross, J. Phys. A **47**, 025204 (2014)]

# Outline

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary

# Different algorithms: (1) Gradient-less algorithms

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

- ▶ Simpler approaches: *direct* or *gradient-less* algorithms. They only require a means to compute  $G[u]$  (i.e. a method to propagate the dynamical equation and compute the resulting cost or target functional).
  - ▶ The most fashionable, the families of *evolutionary* or *genetic* algorithms.
  - ▶ Our choices:
    - ▶ The *simplex* algorithm [J.A. Nelder and R. Mead, Computer Journal **7**, 308 (1965)], and
    - ▶ the NEWUOA algorithm [M. J. D. Powell, IMA J. Numer. Anal. **28**, 649 (2008)].

## Different algorithms: (2) “QOCT-specific” algorithms designed for real-space parametrizations.

- ▶ ZR98: [Zhu and Rabitz, J. Chem. Phys. **109**, 385 (1998)]
- ▶ ZBR98: [Zhu, Botina, and Rabitz, J. Chem. Phys. **108**, 1953 (1998)]
- ▶ Krotov: [V. F. Krotov, Automat. Remote Control **34**, 1863 (1973)]
- ▶ MT03: [Maday and Turinici, J. Chem. Phys. **118**, 8191 (2003)]
- ▶ WG05: [Werschnik and Gross, J. Opt. B **7**, S300 (2005)]

All of these assume that the parameters  $u_1, \dots, u_M$  are the real-time discretization values of the control function  $\varepsilon(t)$ :  $\varepsilon(t_1), \varepsilon(t_2), \dots, \varepsilon(t_M)$ . It is difficult to establish constraints.

## What is OCT?


What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems  
Different  
algorithms  
Different targets  
Different  
parametrizations

## Summary

## Different algorithms: (3) The usual BFGS

In fact, one can use all the gradient-based methods implemented in the GSL library (conjugate gradients, etc.). The parametrization of the control field can then be arbitrary, although only some options have been implemented.



[Castro and Gross, PRL **109**, 153603 (2012)]

# Outline

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

**Different targets**

Different  
parametrizations

Summary

What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

**Different targets**

Different parametrizations

Summary

# Different targets

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

**Different targets**

Different  
parametrizations

Summary

- ▶ State-to-state transitions.

$$J_1[\Psi] = \langle \Psi(T) | \Psi_O \rangle \langle \Psi_O | \Psi(T) \rangle.$$

- ▶ “Ground-state transformation”

$$\Psi = \det[\psi_1, \dots, \psi_N]$$

$$|\Psi_O\rangle = \hat{a}_{a_1}^\dagger \hat{a}_{i_1} \dots \hat{a}_{a_k}^\dagger \hat{a}_{i_k} |\Psi\rangle$$

- ▶ “Excited state” population

$$|\Psi_O\rangle = \sum_{ai} c_{ai} \hat{a}_a^\dagger \hat{a}_i |\Psi\rangle$$



# Different targets

What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

**Different targets**


Different parametrizations

Summary

## ► Ionization

$$J_1[\Psi] = 1 - \sum_i |\langle \Psi(T) | \Psi_i \rangle|^2.$$

$$J_1[\Psi] = \int_{r \in U} d^3r n(\vec{r}, T).$$



[Castro *et al.*, EPL **87**, 53001 (2009)]

# Different targets

What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary

## ► Density, current

$$J_1[\Psi] = \alpha \int d^3r a(n(\vec{r}, T)) + \beta \int d^3r b(\vec{j}(\vec{r}, T)).$$




FIG. 1. (Color online) 1D QW. The overlap between the controlled density  $n$  and the target density  $n_{tg}$  is shown in the top panel. For minimized current ( $w_c = 10$ ) the overlap with the target (found in the inset) at terminal time  $T = 300$  a.u. is better. Additionally, density  $n$  is significantly more stable than without current control ( $w_c = 0$ ). The respective laser pulses are reported in the lower panel.

[Kammerlander *et al.*, PRA **83**, 043413 (2011)]

# Different targets

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

**Different targets**

Different  
parametrizations

Summary

- ▶ Time-dependent targets.

$$J_1[\Psi] = \int_0^T dt \, G(\Psi(t)) .$$


In these cases, the backwards propagation includes an inhomogeneous term.

# Different targets

- High-harmonic generation.

$$J_1[\Psi] = \int_0^\infty d\omega \alpha(\omega) H(\omega),$$

$$H(\omega) = \int_0^T dt |e^{-i\omega t} \frac{d^2}{dt^2} x(t)|^2.$$



# Different targets

What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations


Summary

- Quantum-classical targets.

$$F = F[q, p, \Psi]$$

For example, for the dissociation of two atoms:

$$F = F[\vec{R}_1, \vec{R}_2, \vec{P}_1, \vec{P}_2, \Psi] = (\vec{R}_1 - \vec{R}_2)^2.$$



**Ultrafast and optimal:** optimal ultra-short pulses are theoretically constructed by the combination of time-dependent density functional theory and optimal control, optimizing the classical opposing force on the nuclei for the hydrogen molecule (see picture; top panel: internuclear distance and the bound electrons count; bottom panel: opposing force; gray region: pulse duration).

[Castro, ChemPhysChem **14**, 1488 (2014)]

# Outline

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary

# Different parametrizations

- ▶ Real-time parametrization

$$u_1, u_2, \dots = \varepsilon(t_1), \varepsilon(t_2), \dots$$

- ▶ Fourier coefficients

$$u_1, u_2, \dots = \{a_n, b_n\}.$$

$$\varepsilon(t) = a_0 + \sum_n a_1 \cos(w_n t) + b_n \sin(w_n t)$$

One may want to force:

$$\int_0^T dt \, \varepsilon(t) = 0 \implies a_0 = 0,$$

$$\varepsilon(0) = \varepsilon(T) = 0 \implies \sum_n a_n = 0.$$

- ▶ Constant-fluence constraint.

$$\int_0^T dt \, \varepsilon^2(t) = C,$$

In this case,  $u_1, u_2, \dots$  are certain hyper-spherical angles that are obtained after some transformations.

What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary

# Outline

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

What is OCT?

What OCT calculations can we do with octopus now?

Different dynamical systems

Different algorithms

Different targets

Different parametrizations

Summary



# Summary

What is OCT?

What OCT  
calculations can we  
do with octopus  
now?

Different  
dynamical systems

Different  
algorithms

Different targets

Different  
parametrizations

Summary

Systems	Algorithms	Targets	Parametrizations
Schrödinger	ZR98	State projection	Real-time
TDKS	ZBR98	Density	Fourier
Ehrenfest + wave function	WG05	Current	Fourier (con- strained)
Ehrenfest + TDDFT	MT03	HHG	
	Krotov	Ionization	
	BFGS, CG (GSL)	Classical vars.	
	Nelder-Mead	Spin	
	NEWUOA		