

**Remark.**  $\mathbf{N} = \{1, 2, 3, \dots\}$ .

Let  $F$  be an **ordered field** (for example,  $\mathbf{Q}$  and  $\mathbf{R}$  are ordered fields). Suppose we have a sequence  $(I_n)_{n \in \mathbf{N}}$  of closed bounded intervals

$$I_n = [a_n, b_n] = \{x \in F : a_n \leq x \leq b_n\}.$$

Consider the following two properties.

- (i) For all  $n \in \mathbf{N}$ ,  $I_{n+1} \subseteq I_n$ .
- (ii) For all  $\varepsilon > 0$  there exists an  $N \in \mathbf{N}$  such that  $|I_n| := b_n - a_n < \varepsilon$ .

If  $(I_n)_{n \in \mathbf{N}}$  satisfies property (i), then  $(I_n)_{n \in \mathbf{N}}$  is known as a sequence of **nested intervals**. If  $(I_n)_{n \in \mathbf{N}}$  satisfies both properties (i) and (ii), then  $(I_n)_{n \in \mathbf{N}}$  is known as a sequence of **shrinking nested intervals**.

If  $F$  is such that any sequence  $(I_n)_{n \in \mathbf{N}}$  of shrinking nested intervals has a singleton intersection, i.e.  $\bigcap_{n=1}^{\infty} I_n = \{x\}$  for some  $x \in F$ , then  $F$  is said to have the **nested interval property**.

Let us first show that if  $F$  has the least-upper-bound property, then  $F$  has the nested interval property. Since  $\mathbf{R}$  is the unique ordered field with the least-upper-bound property, the proposition we want to prove is the following.

**Proposition 1.**  $\mathbf{R}$  has the nested interval property.

*Proof.* Let  $(I_n)_{n \in \mathbf{N}}$  be a sequence of shrinking nested intervals, where  $I_n = [a_n, b_n]$ , and let  $A$  be the set of left endpoints, i.e.  $A = \{a_n : n \in \mathbf{N}\}$ . Note that for any  $m \in \mathbf{N}$ ,  $b_m$  is an upper bound of  $A$ :

- if  $n \leq m$ , then  $a_n \leq a_m \leq b_m$ ;
- if  $m < n$ , then  $a_n \leq b_n \leq b_m$ .

(The sequence of left endpoints is non-decreasing and the sequence of right endpoints is non-increasing since the intervals are nested.) Hence  $x := \sup A$  exists in  $\mathbf{R}$  and satisfies  $a_n \leq x \leq b_n$  for each  $n \in \mathbf{N}$ , i.e.  $x \in \bigcap_{n=1}^{\infty} I_n$ . To see that this  $x$  is unique, suppose that  $x_1, x_2 \in \bigcap_{n=1}^{\infty} I_n$  and without loss of generality suppose that  $x_1 \leq x_2$ . Let  $\varepsilon > 0$  be given. Then there exists an  $N \in \mathbf{N}$  such that  $b_N - a_N < \varepsilon$ . Since  $x_1, x_2 \in I_N$  for each  $n \in \mathbf{N}$ , we have

$$a_N \leq x_1, x_2 \leq b_N \implies x_2 - x_1 \leq b_N - a_N < \varepsilon.$$

Hence  $0 \leq x_2 - x_1 < \varepsilon$  for any  $\varepsilon > 0$ ; it follows that  $x_1 = x_2$ . □

Note that the proof just given shows that if  $(I_n)_{n \in \mathbf{N}}$  is only a sequence of nested intervals in  $\mathbf{R}$ , not necessarily shrinking, then  $\bigcap_{n=1}^{\infty} I_n$  is non-empty.

Next, let us show that if  $F$  has the nested interval property, then  $F$  has the least-upper-bound property, i.e.  $F = \mathbf{R}$ .

**Proposition 2.** If  $F$  has the nested interval property, then  $F$  has the least-upper-bound property.

*Proof.* Let  $E \subseteq F$  be non-empty and bounded above by some  $b_1 \in F$ . If  $E$  has a maximum  $x$ , then  $\sup E = x$  and we are done. Otherwise, we shall use an induction argument to construct a sequence  $(I_n)_{n \in \mathbf{N}}$  of shrinking nested intervals. Pick some  $a_1 \in E$ ; it must be the case that  $a_1$  is not an upper bound of  $E$  since  $E$  has no maximum. Let  $I_1 = [a_1, b_1]$ . Then:

- $a_1$  is not an upper bound of  $E$ ;
- $b_1$  is an upper bound of  $E$ ;
- $|I_1| = 2^0(b_1 - a_1)$ .

Suppose that after  $N$  steps we have chosen intervals  $I_n = [a_n, b_n]$ ,  $1 \leq n \leq N$ , such that

- $a_1 \leq \dots \leq a_N$  are not upper bounds of  $E$ ;
- $b_N \leq \dots \leq b_1$  are upper bounds of  $E$ ;
- $|I_n| = 2^{-(n-1)}(b_1 - a_1)$  for  $1 \leq n \leq N$ .

Let  $m = \frac{a_N + b_N}{2}$ , the midpoint of the interval  $I_N$ . If  $m$  is not an upper bound of  $E$ , set

$$a_{N+1} = m, b_{N+1} = b_N, \text{ and } I_{N+1} = [a_{N+1}, b_{N+1}].$$

If  $m$  is an upper bound of  $E$ , set

$$a_{N+1} = a_N, b_{N+1} = m, \text{ and } I_{N+1} = [a_{N+1}, b_{N+1}].$$

In either case, we have chosen intervals  $I_n = [a_n, b_n]$ ,  $1 \leq n \leq N + 1$ , such that

- $a_1 \leq \dots \leq a_{N+1}$  are not upper bounds of  $E$ ;
- $b_{N+1} \leq \dots \leq b_1$  are upper bounds of  $E$ ;
- $|I_n| = 2^{-(n-1)}(b_1 - a_1)$  for  $1 \leq n \leq N + 1$ .

In this way we obtain a sequence  $(I_n)_{n \in \mathbf{N}}$  of intervals  $I_n = [a_n, b_n]$  such that

- $a_1 \leq \dots \leq a_n \leq \dots$  are not upper bounds of  $E$ ;

- $\cdots \leq b_n \leq \cdots \leq b_1$  are upper bounds of  $E$ ;
- $|I_n| = 2^{-(n-1)}(b_1 - a_1)$  for all  $n \in \mathbf{N}$ .

Hence  $(I_n)_{n \in \mathbf{N}}$  is a sequence of shrinking nested intervals. By assumption,  $F$  has the nested interval property, so there exists an  $x \in F$  such that  $\bigcap_{n=1}^{\infty} I_n = \{x\}$ . We claim that  $x = \sup E$ . For  $y \in E$ , suppose  $x < y$ . Then there is an  $N \in \mathbf{N}$  such that

$$b_N - a_N < y - x \implies x + (b_N - a_N) < y.$$

Since  $x \in \bigcap_{n=1}^{\infty} I_n$ , we have

$$a_N \leq x \implies 0 \leq x - a_N \implies b_N \leq x + (b_N - a_N) < y.$$

This is a contradiction since  $b_N$  is an upper bound of  $E$ . It follows that  $y \leq x$ , so that  $x$  is an upper bound of  $E$ . Suppose that  $z \in F$  is such that  $z < x$ . There is an  $N \in \mathbf{N}$  such that

$$b_N - a_N < x - z \implies z < x - (b_N - a_N).$$

Since  $x \in \bigcap_{n=1}^{\infty} I_n$ , we have

$$x \leq b_N \implies x - b_N \leq 0 \implies x - (b_N - a_N) \leq a_N \implies z < a_N.$$

It follows that  $z$  is not an upper bound of  $E$  since  $a_N$  is not an upper bound of  $E$ . We may conclude that  $x$  is the least upper bound of  $E$ , i.e.  $x = \sup E$ .  $\square$

So for ordered fields, the nested interval property and the least-upper-bound property are equivalent. In light of the fact that  $\mathbf{R}$  is the unique ordered field with the least-upper-bound property, we see that  $\mathbf{R}$  is the unique ordered field with the nested interval property.