Blood culture time to positivity and mortality in bacteremia: a systematic review and Bayesian meta-analysis

Monica Basso Russell E. Lewis

2026-10-30

Abstract

Background. Blood culture time to positivity (TTP) has been proposed as a prognostic marker in bacteremia, but studies use different thresholds, pathogens, and outcomes. We synthesized the association between short versus long TTP and clinical outcomes.

Methods. We searched PubMed, Embase, Scopus, the Cochrane Library, and Web of Science (updated May 2026) and extracted effect sizes from 100 studies meeting strict eligibility criteria. Log odds ratios for all-cause mortality (short versus long TTP) were pooled with a Bayesian random-effects model. We examined pathogen class, TTP cutpoint, publication bias, and a sensitivity analysis adding studies meeting more inclusive criteria. Persistent bacteremia and microbiological clearance were analyzed as secondary outcomes.

Results. Fifty-eight studies (72 effect sizes) reported mortality. Short TTP was associated with higher mortality (pooled OR 2.06; 95% credible interval [CrI], 1.67 to 2.55), with substantial heterogeneity (median I^2 62%; 95% prediction interval, 0.5 to 8.6). The association was larger in Gram-negative (OR 2.66; 95% CrI, 1.72 to 4.06) than Gram-positive bacteremia (OR 1.48; 95% CrI, 0.85 to 2.58), although the credible intervals overlapped. Higher TTP cutpoints were associated with weaker associations overall, but this was not consistent across subgroups. The funnel plot was asymmetric, and trim-and-fill gave an adjusted OR of 1.51 (95% CI, 1.23 to 1.86). Short TTP was also associated with persistent bacteremia (OR 1.74; 95% CrI, 1.09 to 2.80); data on clearance were sparse.

Conclusions. Short TTP is associated with higher mortality, particularly in Gram-negative bacteremia, but heterogeneity, cutpoint dependence, probable confounding, and small-study effects limit its use as a stand-alone prognostic tool. Standardized thresholds and adjusted, individual-patient analyses are needed.

Introduction

Bloodstream infection is a major cause of morbidity and mortality, and early risk stratification remains a practical clinical problem. Automated continuous-monitoring blood culture systems record the time to positivity (TTP), the interval from the start of incubation to the instrument's detection of microbial growth. Because growth kinetics depend on the starting inoculum and on organism-specific factors, shorter TTP has been proposed as a readily available, laboratory-generated

marker of high bacterial burden and severe infection.

Individual cohorts have examined whether TTP carries prognostic information in specific settings, including *Staphylococcus aureus* bacteremia, where TTP has been explored as a marker of endocarditis and poor outcome (1–3); Gram-negative bacteremia, including *Pseudomonas aeruginosa* (4, 5) and *Escherichia coli* (6); candidemia (7); and bacteremia in neutropenic patients, where TTP has been suggested as a stewardship tool (8). These studies differ in organism mix, patient population, outcome definition, and, importantly, the TTP threshold used to define "short" versus "long" TTP. Whether the association is consistent across pathogens, whether it varies with the chosen cutpoint, and how large it is once adjusted estimates are considered remain unclear.

We therefore updated a systematic review of the association between blood culture TTP and clinical outcomes in bacteremia. Our primary objective was to estimate the association between short versus long TTP and all-cause mortality using a Bayesian random-effects meta-analysis. Secondary objectives were to examine heterogeneity by pathogen class and by TTP cutpoint, to assess publication bias and the robustness of the findings to study inclusion criteria, and to summarize associations with persistent bacteremia and other markers of clinical failure.

Materials and Methods

Protocol and registration

The review is reported according to the PRISMA 2020 statement and the PRISMA-S extension for searches. A protocol was prepared for registration in PROSPERO (registration number: [PROSPERO registration number to be inserted]). The review builds on a master's thesis at the University of Padova (F. Pangallo, 2025/2026; supervisor R. E. Lewis) that produced the original 57-study extraction database. Because that work was substantially complete before the

protocol was registered, registration is retrospective for the baseline studies, and the 2026 update (additional search, screening, extraction, and the inclusive-criterion re-screen) was added afterwards. The analysis plan was developed during the thesis; deviations from the protocol are listed in the supplementary notebook (Methods and PRISMA).

Search strategy and study selection

Five databases were searched (PubMed, Embase, Scopus, Cochrane Library, and Web of Science), with an original search and a May 2026 update [insert run dates and per-database record counts]. The searches yielded 4,605 records, with 3,196 unique records after de-duplication. CINAHL, trial registers, and preprint servers were not searched, and no reference-list searching or citation tracking was performed; these are deviations from the registered protocol. Candidate records were triaged for relevance [confirm the role of the Elicit semantic search tool and PubMed enrichment in this step, and reconcile Alvarez 2012 and Peri 2026, which project notes record as handsearch or snowball identifications], and 355 reports were sought for full-text review. Of these, 53 were not retrieved after pre-screening as ineligible (mainly conference abstracts, non-English papers, and methods-only papers), and 302 full-text reports were assessed (5 additional PDFs duplicated baseline studies). Under the strict criterion 43 reports were included, which added to the 57 baseline studies gave 100 studies; 259 were excluded (104 ineligible design, 65 no TTP measured, 54 no quantitative TTP-outcome association, 29 no clinical outcome, 5 duplicates, 2 not bloodstream infection). A second, inclusive re-screen of the 54 reports excluded only for lacking a quantitative association identified 40 further studies (sensitivity pool, 140 studies in total). Eligibility criteria and the full search strings are in the supplementary notebook (Methods and PRISMA) and `search_strategies.md`. Figure 1 shows study flow.




Figure 1: PRISMA 2020 flow diagram for the original search and the May 2026 update.

Data extraction and risk of bias

Data were extracted into a structured relational workbook with linked tables for study characteristics, population, TTP reporting (continuous, dichotomized, or categorical; units; cutpoints), and outcomes (events and totals by TTP group, reported effect estimates, adjustment status, and outcome definitions). TTP was recorded in hours. Where a study reported both 2×2 data and an effect estimate, the 2×2 data were preferred. The baseline 57 studies were extracted by the thesis author [confirm whether extraction was independently verified]. For the 2026 update, full-text screening and data extraction were performed by a large language model agent working from the full-text PDFs and the written eligibility criteria and extraction codebook. Each of the 43 strict-pool extractions was then independently verified field by field against the source PDF by a separate agent that had not performed the original extraction; this led to 7 corrections before analysis. Studies using TTP only as part of a catheter-related bloodstream infection case definition were excluded from effect-size pooling. The human contribution to screening and extraction decisions (spot-checking, adjudication, or second human review) is described in the AI-use statement below [confirm], and we did not perform dual independent human screening or report inter-rater agreement.

Risk of bias was assessed for every included study with the Quality in Prognosis Studies (QUIPS) tool across six domains (participation, attrition, prognostic factor measurement, outcome measurement, confounding, and statistical analysis and reporting), each rated as low, moderate, or high, with an overall rating per study (notebook). For the update these ratings were generated by the extraction agent and were not independently re-rated by a second human.

Statistical analysis

Log odds ratios for mortality (short versus long TTP) were computed with `metafor::escalc()` from 2×2 tables where available, otherwise back-calculated from reported OR, HR, or RR with 95%

confidence intervals. Effect sizes were pooled with a Bayesian random-effects model (`brms`; study-level random intercepts) with weakly informative priors ($\mu \sim \mathcal{N}(0, 1)$, $\tau \sim$ Half-Cauchy$(0, 0.5)$). Four chains of 4,000 iterations (1,000 warm-up) were run with `adapt_delta = 0.99`. Convergence was assessed with $\hat{R}$ and effective sample sizes. Secondary analyses examined pathogen class, TTP cutpoint (meta-regression), publication bias, and a sensitivity pool that added studies meeting a more inclusive criterion. All code and data are available in the project repository.

Results

Study selection and characteristics

The primary (strict) pool comprised 100 studies, of which 58 reported mortality and contributed 72 mortality effect sizes. Overall risk of bias (QUIPS) was low in 24, moderate in 62, high in 13, and critical in 1 studies (notebook).

Primary analysis

Short TTP was associated with higher mortality (pooled OR 2.06 (95% CrI, 1.67 to 2.55); Figure 2, Table 1). The posterior probability that short TTP increases mortality was 100.0%, and heterogeneity was substantial (median $I^2 = 62\%$).

Table 1: Pooled effect of short vs long TTP on mortality (OR > 1 = short TTP harmful). Primary pool only (strict criterion).

Parameter	Value
Median OR (short vs long TTP)	2.06
95% credible interval	[1.67, 2.55]
95% prediction interval (new study)	[0.50, 8.41]

Table 1: Pooled effect of short vs long TTP on mortality (OR > 1 = short TTP harmful). Primary pool only (strict criterion).

Parameter	Value
P(short TTP harmful)	100.0%
Between-study SD ( )	0.69
Heterogeneity (I^2)	62%

116 Source: Time-to-Positivity and Mortality in Bacteremia




Figure 2: Forest plot of OR (short vs long TTP) on mortality — primary pool. Each row is a single effect size; multi-arm studies appear multiple times. Pooled estimate from the Bayesian model in dark navy.

117 Source: Time-to-Positivity and Mortality in Bacteremia

Pathogen class




Figure 3: **Figure 1.** Posterior distributions of the pooled odds ratio (OR) for all-cause mortality comparing short versus long time-to-positivity (TTP), stratified by pathogen class. Each density represents the full posterior from a Bayesian random-effects meta-analysis (72 mortality effect sizes; 100 studies). The filled point marks the posterior median; the horizontal bar spans the 95% credible interval (CrI). The vertical dashed line indicates OR = 1 (no effect). 'Short' and 'long' TTP were defined by individual studies using study-specific thresholds (median cutpoint 12 h; range 2–72 h). Outcome was predominantly 30-day mortality. Overall: OR = 2.06 (95% CrI 1.67–2.55); Gram-negative: OR = 2.66 (95% CrI 1.72–4.06); Gram-positive: OR = 1.48 (95% CrI 0.85–2.58). OR $>$ 1 indicates higher mortality with shorter TTP.

Source: Time-to-Positivity and Mortality in Bacteremia

Pathogen class was available for 52 of 72 mortality effect sizes (Gram-negative, 26 effect sizes from 17 studies; Gram-positive, 13 from 11; fungal, 6 from 6; mixed or polymicrobial, 7 from 5). The remaining 20 effect sizes had no pathogen classification and were excluded from the stratified models. Short TTP was associated with higher mortality in Gram-negative bacteremia (OR 2.66 (95% CrI, 1.72 to 4.06)) and, with a smaller point estimate and a credible interval that included 1, in Gram-positive bacteremia (OR 1.48 (95% CrI, 0.85 to 2.58); Figure 3, Table 2). In a meta-regression with both classes, the odds ratio in Gram-positive studies was 0.61 times that in Gram-negative

studies (95% CrI 0.36 to 1.02), with a posterior probability of 97% that the association is weaker in Gram-positive studies. This interval is wide and just includes 1, so the evidence for a difference between classes is suggestive rather than conclusive. For fungal bacteremia the pooled estimate was imprecise (OR 1.31 (95% CrI, 0.92 to 2.12), 6 studies). Heterogeneity remained considerable within both Gram-negative (median I^2 61%) and Gram-positive (74%) strata.

Table 2: Pooled OR by pathogen subgroup.

Subgroup	N effect sizes	Median OR	95% CrI	P(short TTP harmful)
Gram-positive	13	1.48	[0.85, 2.58]	92.7%
Gram-negative	26	2.66	[1.72, 4.06]	100.0%

Source: Time-to-Positivity and Mortality in Bacteremia

TTP cutpoint

A TTP cutpoint was reported for 42 of 72 mortality effect sizes (29 studies); the remainder had no cutpoint recorded in the extraction database. Cutpoints ranged from 2 to 72 h (median 12 h). Across all effect sizes with a cutpoint, the log odds ratio changed by -0.035 (95% CrI, -0.056 to -0.015) per additional hour, so higher thresholds were associated with weaker associations (Figure 4). When the analysis was restricted to cutpoints of 24 h or less, the slope was attenuated and its interval included zero (-0.024 (95% CrI, -0.075 to 0.028)), so the gradient depends on the few studies using long thresholds.

Within pathogen class the pattern differed. Among Gram-negative effect sizes (median cutpoint 10 h) the slope was positive (0.062 (95% CrI, -0.005 to 0.130); Table 3), whereas among Gram-positive effect sizes it was negative with an interval that included zero (-0.029 (95% CrI, -0.058 to 0.004);

Table 4). The overall negative gradient therefore does not hold within Gram-negative bacteremia, and these class-specific analyses rest on small numbers of effect sizes.




Figure 4: TTP cutpoint meta-regression. The line is the Bayesian posterior median predicted OR by cutpoint; ribbon is the 95% CrI. Dark points are observed 2×2-derived effect sizes (size precision).

Source: Time-to-Positivity and Mortality in Bacteremia

Table 3: Gram-negative cutpoint meta-regression slope and predicted OR at key thresholds.

Parameter	Estimate	95% CrI	P(direction)
Slope (log-OR per hour above median)	0.062	[-0.005, 0.130]	96.8%
Predicted OR at cutpoint 8 h	3.03	[2.02, 4.52]	100.0%
Predicted OR at cutpoint 10 h	3.43	[2.40, 4.87]	100.0%
Predicted OR at cutpoint 13 h	4.14	[2.84, 5.93]	100.0%
Predicted OR at cutpoint 18 h	5.65	[3.17, 9.95]	100.0%

147 Source: Time-to-Positivity and Mortality in Bacteremia

Table 4: Gram-positive cutpoint meta-regression: slope comparison between full model and sensitivity analysis excluding Minejima2019 (72 h).

Model	N studies	Slope (log-OR/h)	P(slope < 0)
Full (8–72 h)	7	-0.029 [-0.058, 0.004]	96.5%
Sensitivity (excl. Minejima2019)	6	-0.081 [-0.323, 0.214]	75.4%

148 Source: Time-to-Positivity and Mortality in Bacteremia

149 Publication bias

150 The funnel plot of mortality effect sizes was asymmetric (Figure 5). Egger's regression test indicated
151 small-study effects ($t = 6.19$, $df = 70$, $p < 0.0001$; Table 5), and the regression-based estimate for a
152 study with zero standard error was close to no association (OR 0.99; 95% CI 0.97 to 1.01). Begg's
153 rank correlation test did not detect asymmetry ($p = 0.47$). The trim-and-fill method imputed 19
154 missing studies and gave an adjusted pooled OR of 1.51 (95% CI, 1.23 to 1.86), lower than the
155 unadjusted frequentist estimate (OR 2.05). These tests treat all effect sizes as independent, assume
156 that asymmetry is due to selective reporting, and have limited reliability when heterogeneity is high,
157 so they cannot distinguish publication bias from genuine heterogeneity, such as larger associations in
158 smaller studies that used more extreme or optimized cutpoints. The evidence nevertheless suggests
159 that the pooled association may be overestimated.




Figure 5: Contour-enhanced funnel plot (primary pool, mortality). Studies in white zones are not statistically significant at the labelled threshold; shaded outer zones indicate increasing levels of significance. Asymmetry toward the right (short TTP harmful) would suggest positive-result publication bias.

160 Source: Time-to-Positivity and Mortality in Bacteremia

Table 5: Egger's regression test for funnel-plot asymmetry (primary pool, mortality; metafor::regtest, weighted regression on the standard error). The t-statistic tests asymmetry; the limit estimate is the predicted effect for a hypothetical study with zero standard error.

Statistic	Value
Asymmetry test t-statistic	6.19
df	70
p-value	<1e-04
Limit estimate as SE -> 0 (log OR)	-0.010
95% CI (log OR)	[-0.034, 0.014]

Table 5: Egger's regression test for funnel-plot asymmetry (primary pool, mortality; metafor::regtest, weighted regression on the standard error). The t-statistic tests asymmetry; the limit estimate is the predicted effect for a hypothetical study with zero standard error.

Statistic	Value
Limit estimate as SE -> 0 (OR)	0.99

Source: Time-to-Positivity and Mortality in Bacteremia

Sensitivity analyses

Adding the studies that met only the inclusive criterion increased the number of studies reporting mortality to 67 (84 effect sizes) and gave a pooled OR of 1.92 (95% CrI, 1.58 to 2.35), similar to the primary estimate (Figure 6). Pooled estimates by data source were consistent in direction. Relative to 2×2-derived effect sizes (49 effect sizes), the log odds ratio for effect sizes back-calculated from adjusted reported estimates (15 effect sizes) was -0.26 (95% CrI, -0.68 to 0.16) and for unadjusted reported estimates (8 effect sizes) was -0.12 (95% CrI, -0.62 to 0.38). Both intervals included zero. Directly comparing unadjusted with adjusted reported estimates, the difference in log odds ratio was 0.12 (95% CrI, -0.39 to 0.62). Association strength tended to decrease with publication year (change in log odds ratio per year, -0.04 (95% CrI, -0.08 to 0.00)), with an interval that only just included zero. None of these moderator analyses showed a clear difference, but all were underpowered relative to the observed heterogeneity.




Figure 6: Posterior densities of pooled OR — primary pool vs sensitivity pool (with inclusive-only studies added).

Source: Time-to-Positivity and Mortality in Bacteremia

Clinical failure outcomes

Outcomes other than mortality were reported far less often (Table 6). Persistent bacteremia was the most frequently reported, with 12 effect sizes from 12 studies, followed by microbiological clearance (4 effect sizes, 3 studies). No usable effect sizes were available for relapse (0).

Short TTP was associated with higher odds of persistent bacteremia (pooled OR 1.74 (95% CrI, 1.09 to 2.80); Figure 7), with substantial heterogeneity (median I^2 69%). Outcome definitions were not uniform: on review of the outcome definitions, 6 effect sizes defined persistent bacteremia by duration of positive cultures, 3 defined it by endocarditis or valve involvement, and 3 used other definitions. In a meta-regression by definition group, endocarditis-based definitions had a log odds ratio 0.59 higher

than duration-based definitions (95% CrI -0.19 to 1.24), which is compatible with no difference given the small number of effect sizes per group. The pooled persistent-bacteremia estimate should therefore be interpreted as an association with a heterogeneous composite of outcomes.

For microbiological clearance, the pooled estimate was too imprecise to interpret (OR 1.63 (95% CrI, 0.24 to 9.47)), reflecting only 4 effect sizes with very large between-study heterogeneity (median I^2 89%).

Table 6: Available effect sizes per clinical-failure outcome (strict pool).

Outcome	Effect sizes	Unique studies
Persistent bacteremia	12	12
Microbiological clearance	4	3
Relapse	0	0

Source: TTP and Clinical Failure Outcomes




Figure 7: Forest plot of OR (short vs long TTP) for persistent bacteremia. NB: many studies coded under 'persistent_bacteremia' actually report endocarditis (the Codebook's outcome_type enum lacks 'endocarditis'). Review outcome_definition in each row before interpretation.

Source: TTP and Clinical Failure Outcomes

Discussion

In this Bayesian random-effects meta-analysis, short TTP was associated with higher mortality in bacteremia (pooled OR 2.06 (95% CrI, 1.67 to 2.55); 58 studies, 72 effect sizes). The posterior probability that the association is in the direction of harm was 100.0%. Between-study heterogeneity was moderate to substantial (median I^2 62%), and the 95% prediction interval for a new study (0.51 to 8.50) is wide and includes no association. The pooled estimate is therefore best read as an

average association, not as the effect expected in any particular setting. Adding studies that met only a more inclusive criterion gave a similar estimate (OR 1.92 (95% CrI, 1.58 to 2.35)).

Pathogen class. The association was larger in Gram-negative bacteremia (OR 2.66 (95% CrI, 1.72 to 4.06)) than in Gram-positive bacteremia (OR 1.48 (95% CrI, 0.85 to 2.58)), for which the credible interval included 1. The two intervals overlap, so this contrast should be considered hypothesis-generating rather than confirmed. A plausible explanation is that TTP in Gram-positive infection, notably *S. aureus*, is influenced by factors other than burden of disease, such as the source of infection and the presence of endovascular foci, whereas in Gram-negative infection it may track inoculum and host response more directly. This explanation is speculative and was not tested directly here.

Study design and adjustment. Effect sizes derived from 2×2 tables (OR 2.28 (95% CrI, 1.74 to 3.05)) were larger than those back-calculated from reported effect estimates (OR 1.71 (95% CrI, 1.28 to 2.36)). Meta-regression by data source did not show a clear difference, and the intervals included zero, so the data are compatible with either similar or lower adjusted associations. Crude 2×2 comparisons may carry more confounding by severity of illness, source, or pathogen, in which case the pooled estimate could overstate the independent prognostic contribution of TTP, but the groups also differ in other respects.

TTP cutpoint. Studies defined "short" TTP using different thresholds (median cutpoint 12 h), and 30 of 72 mortality effect sizes did not have a reported cutpoint. In meta-regression among the remaining effect sizes, the log OR decreased by 0.036 per additional hour of cutpoint (95% CrI 0.015 to 0.056), meaning that stricter (shorter) thresholds were associated with larger odds ratios overall (Figure 4). This gradient was not robust: it was attenuated when cutpoints above 24 h were excluded, and it was reversed within Gram-negative studies. Because thresholds are often chosen

post hoc to maximize discrimination in the study sample, the gradient may partly reflect optimistic cutpoint selection or differences in pathogen mix, and it should not be interpreted as validating any single clinical threshold.

Other outcomes. For persistent bacteremia, short TTP was associated with higher odds (OR 1.74 (95% CrI, 1.09 to 2.80)), but outcome definitions varied widely between studies. Evidence for microbiological clearance was too sparse and imprecise to interpret (OR 1.63 (95% CrI, 0.24 to 9.47)) (notebook).

Strengths and limitations. Strengths include a multi-database search with a May 2026 update, a distinction between strict and inclusive study pools, a Bayesian framework that yields prediction intervals and direct probability statements, and a fully reproducible code base. Limitations are substantial. Most included studies were retrospective cohorts and risk of bias was rated moderate or worse in most (76 of 100 studies). Several studies contributed multiple effect sizes, which we handled with study-level random intercepts but which do not remove all dependence. The TTP threshold, outcome timing, and adjustment sets were not standardized, and adjustment status varied across estimates. Blood culture system, bottle type, transport delay, and prior antibiotic exposure affect TTP and were incompletely reported, so residual confounding by measurement conditions is likely. The funnel plot was asymmetric and Egger's test indicated small-study effects (Figure 5, Table 5). Trim-and-fill reduced the pooled estimate to OR 1.51, so the primary estimate may overstate the association, although asymmetry can also arise from heterogeneity and cutpoint optimization, and these tests do not account for multiple effect sizes per study. Finally, the meta-analysis is based on aggregate data, which prevents examination of TTP as a continuous predictor within patients.

Implications. Short TTP is associated with higher mortality on average, but heterogeneity, the dependence on cutpoint, and probable confounding limit its use as a stand-alone prognostic

tool. These results support using TTP as one contextual signal, particularly in Gram-negative bacteremia, alongside clinical severity and pathogen identification. Individual-patient-data analyses with standardized TTP measurement, and prospective validation of pathogen-specific thresholds, are needed before TTP can be used to guide treatment decisions.

Funding

No specific funding was received for this study. The authors are supported by their institutional salaries (University of Padova). [Confirm.]

Conflicts of interest

R.E.L. serves as an Associate Editor at the Journal of Antimicrobial Chemotherapy. The other author declares no conflicts of interest. [Confirm with all co-authors.]

Data and code availability

The extraction workbook, analysis code, and rendered notebooks are included in the manuscript project. [Insert repository and DOI, for example an OSF project, once deposited.]

Use of artificial intelligence

A large language model agent was used to assist with full-text screening and data extraction for the 2026 update, and a second, independent agent verified the extractions against the source PDFs, as described in the Methods. [State the model name and version, how authors reviewed outputs, and who is responsible for the final data. Confirm the wording against ASM's current policy on AI use.] Analysis code was written and run in R and Stan. The authors take responsibility for the content

of the manuscript.

Acknowledgments

[None to declare / add acknowledgments.]

References

1. Sowden D, Anstey C, Faddy M. 2008. Blood culture time to positivity as a predictor of mortality in community acquired Methicillin-susceptible Staphylococcus aureus bacteremia. Journal of Infection 56:295–296.

2. Kahn F, Resman F, Bergmark S, Filiptsev P, Nilson B, Gilje P, Rasmussen M. 2021. Time to blood culture positivity in Staphylococcus aureus bacteraemia to determine risk of infective endocarditis. Clinical Microbiology and Infection 27:1345.e7–1345.e12.

3. Strömdahl M, Hagman K, Hedman K, Westman A, Hedenstierna M, Ursing J. 2025. Time to Staphylococcus aureus Blood Culture Positivity as a Risk Marker of Infective Endocarditis: A Retrospective Cohort Study. Clinical Infectious Diseases 80:727–734.

4. Willmann M, Kuebart I, Vogel W, Flesch I, Markert U, Marschal M, Schröppel K, Autenrieth IB, Hölzl F, Peter S. 2013. Time to positivity as prognostic tool in patients with Pseudomonas aeruginosa bloodstream infection. Journal of Infection 67:416–423.

5. Tang P-C, Lee C-C, Li C-W, Li M-C, Ko W-C, Lee N-Y. 2017. Time-to-positivity of blood culture: An independent prognostic factor of monomicrobial Pseudomonas aeruginosa bacteremia. Journal of Microbiology, Immunology and Infection 50:486–493.

6. Martínez JA, Soto S, Fabrega A, Almela M, Mensa J, Soriano A, Marco F, Jimenez De Anta MT, Vila J. 2006. Relationship of phylogenetic background, biofilm production, and time to detection of growth in blood culture vials with clinical variables and prognosis associated with Escherichia coli bacteremia. Journal of Clinical Microbiology 44:1468–1474.

7. Keighley C, Pope AL, Marriott D, Chen SC-A, Slavin MA. 2023. Time-to-positivity in bloodstream infection for Candida species as a prognostic marker for mortality. Medical Mycology 61.

8. Puerta-Alcalde P, Cardozo C, Suárez-Lledó M, Rodríguez-Núñez O, Morata L, Fehér C, Marco F, Del Río A, Martínez JA, Mensa J, Rovira M, Esteve J, Soriano A, Garcia-Vidal C. 2019. Current time-to-positivity of blood cultures in febrile neutropenia: a tool to be used in stewardship de-escalation strategies. Clinical Microbiology and Infection 25:447–453.

Supplementary appendix: model specification and diagnostics

Model specification

All pooled and moderator models were Bayesian random-effects models fitted with `brms`: $y_i \mid \mathrm{SE}_i \sim \mathcal{N}(\mu + u_s,\ \mathrm{SE}_i^2)$ with study-level random intercepts $u_s \sim \mathcal{N}(0, \tau^2)$, where y_i is the log odds ratio for effect size i and SE_i its standard error. Priors were $\mu \sim \mathcal{N}(0, 1)$, $\tau \sim \text{Half-Cauchy}(0, 0.5)$, and $\mathcal{N}(0, 0.5)$ for moderator coefficients. Each model used 4 chains of 4,000 iterations (1,000 warm-up;

12,000 post-warm-up draws), `adapt_delta = 0.99`, and `max_treedepth = 12`. The persistent bacteremia and microbiological clearance models were fitted with the CmdStan backend and all others with rstan. Convergence was judged by $\hat{R} < 1.01$, bulk and tail effective sample sizes, divergent transitions, and visual inspection of trace and rank plots. Code is in `R/02_fit_models.R` and the notebooks.

Convergence across all fitted models

Table 7: Convergence diagnostics for every fitted model. Parameters are population-level effects and the between-study SD; study-level random effects are not summarized.

Model	Max R-hat	Min bulk ESS	Min tail ESS	Divergent transitions	Treedepth limit hits
fit_2x2	1.001	3334	4906	0	0
fit_adjustment	1.004	1702	2781	0	0
fit_base	1.003	1429	2831	0	0
fit_clr	1.000	2748	3191	2	0
fit_cutpoint_le24	1.001	3806	5910	0	0
fit_cutpoint	1.001	3854	5342	0	0
fit_es_source	1.003	1978	3585	0	0
fit_fungal	1.001	3388	4296	8	0
fit_gneg_cutpoint	1.000	3219	3938	0	0
fit_gpos_cut-point_sens	1.001	2110	3010	0	0
fit_gpos_cutpoint	1.001	2304	3438	1	0
fit_gram_neg	1.001	2557	3765	0	0

Table 7: Convergence diagnostics for every fitted model. Parameters are population-level effects and the between-study SD; study-level random effects are not summarized.

Model	Max R-hat	Min bulk ESS	Min tail ESS	Divergent transitions	Treedepth limit hits
fit_gram_pos	1.002	2650	4142	0	0
fit_metareg	1.003	3164	4357	0	0
fit_mixed_source	1.002	2246	3595	0	0
fit_pb_defgroup	1.001	2866	4597	0	0
fit_pb	1.002	2135	3478	0	0
fit_reported	1.003	1958	3610	0	0
fit_src_binary	1.001	2795	5091	0	0
fit_year	1.003	1572	3301	0	0
fit_base (sensitivity pool)	1.002	1789	2906	0	0

Source: Article Notebook

The maximum $\hat{R}$ across models was 1.004 and the minimum bulk ESS was 1,429. Divergent transitions occurred in 11 post-warm-up draws in total, confined to these models: fit_clr, fit_fungal, fit_gpos_cutpoint. Estimates from models with divergent transitions, which have few effect sizes, should be interpreted cautiously.

Primary pool: pooled model

Table 8: Convergence diagnostics for the primary pooled model.

Parameter	R	ESS_bulk
b_Intercept	1.003	1429
sd_study_id__Intercept	1.002	2043
sigma	NA	NA
Intercept	1.003	1429
r_study_id[Alvarez2012,Intercept]	1.001	10557
r_study_id[Bae2021,Intercept]	1.000	12229
r_study_id[Balaji2024,Intercept]	1.002	2836
r_study_id[Barr2022,Intercept]	1.000	11548
r_study_id[Blackberg2022,Intercept]	1.001	11388
r_study_id[Blackberg2023,Intercept]	1.001	9980
r_study_id[Bo2011,Intercept]	1.000	7848
r_study_id[Chavez2022,Intercept]	1.003	1426
r_study_id[Chen2018,Intercept]	1.000	6414
r_study_id[Chen2020,Intercept]	1.000	14148
r_study_id[Chen2023,Intercept]	1.000	13576
r_study_id[Cheng2022,Intercept]	1.000	13904
r_study_id[Cilloniz2017,Intercept]	1.000	7851
r_study_id[Deguchi2024,Intercept]	1.000	12353
r_study_id[Fenoll2022,Intercept]	1.000	10865
r_study_id[Hamilton2021,Intercept]	1.003	1432
r_study_id[Hou2023,Intercept]	1.001	7382

Table 8: Convergence diagnostics for the primary pooled model.

Parameter	R	ESS_bulk
r_study_id[Hsieh2022,Intercept]	1.001	3480
r_study_id[IrigoyenSierakowski2024,Intercept]	1.000	16462
r_study_id[Jayabalakrishnan2022,Intercept]	1.000	16215
r_study_id[Ji2020,Intercept]	1.000	6152
r_study_id[Kahn2021,Intercept]	1.000	6419
r_study_id[Keighley2023,Intercept]	1.000	16962
r_study_id[Kim2010,Intercept]	1.001	5819
r_study_id[Kim2013,Intercept]	1.000	9157
r_study_id[Laupland2022,Intercept]	1.002	1610
r_study_id[Laupland2024,Intercept]	1.003	1478
r_study_id[Liao2009,Intercept]	1.000	8344
r_study_id[Lin2016,Intercept]	1.000	13218
r_study_id[Marco2025,Intercept]	1.001	4825
r_study_id[Marra2006,Intercept]	1.000	13785
r_study_id[Minejima2019,Intercept]	1.001	8153
r_study_id[Nunes2013,Intercept]	1.000	13928
r_study_id[Ok2013,Intercept]	1.000	12816
r_study_id[Palmer2013,Intercept]	1.001	15976
r_study_id[PapadimitriouOlivgeris2023,Intercept]	1.000	5065
r_study_id[Paquette2021,Intercept]	1.003	1576
r_study_id[Peralta2007,Intercept]	1.000	10735

Table 8: Convergence diagnostics for the primary pooled model.

Parameter	R	ESS_bulk
r_study_id[PerezRodriguez2024,Intercept]	1.000	10843
r_study_id[Prinsloo2023,Intercept]	1.000	9664
r_study_id[PuertaAlcalde2019,Intercept]	1.001	6856
r_study_id[Recio2020,Intercept]	1.003	1490
r_study_id[Rojas2013,Intercept]	1.001	14005
r_study_id[Rolo2022,Intercept]	1.000	8569
r_study_id[Sachu2024,Intercept]	1.000	11112
r_study_id[Savithri2011,Intercept]	1.000	16329
r_study_id[Simeon2019,Intercept]	1.001	5714
r_study_id[Sowden2008,Intercept]	1.000	12698
r_study_id[Takahashi2022,Intercept]	1.000	13238
r_study_id[Tang2017,Intercept]	1.001	7028
r_study_id[Taur2010,Intercept]	1.003	1440
r_study_id[Tsai2025,Intercept]	1.003	1462
r_study_id[Turkeltaub2024,Intercept]	1.000	14546
r_study_id[Willmann2013,Intercept]	1.000	12819
r_study_id[Zhang2016,Intercept]	1.000	7888
r_study_id[Zhang2020,Intercept]	1.000	9920
r_study_id[Zhang2021,Intercept]	1.001	7441
r_study_id[Zouggari2022,Intercept]	1.000	10868

Source: Time-to-Positivity and Mortality in Bacteremia




Figure 8: Trace plots for key parameters of the primary pooled model. Well-mixed chains (caterpillar appearance) indicate good convergence.

Source: Time-to-Positivity and Mortality in Bacteremia




Figure 9: Rank histograms for key parameters. Uniform bars across all chains indicate no chain-mixing issues.

295 Source: Time-to-Positivity and Mortality in Bacteremia




Figure 10: R dot plot for all parameters of the primary pooled model. All values should fall below 1.01 (dashed line).

296 Source: Time-to-Positivity and Mortality in Bacteremia

Sensitivity pool: pooled model




Figure 11: Trace plots for key parameters of the sensitivity (n=140) model.

Source: Time-to-Positivity and Mortality in Bacteremia




Figure 12: Rank histograms for key parameters of the sensitivity (n=140) model.

Source: Time-to-Positivity and Mortality in Bacteremia




Figure 13: R dot plot for all parameters of the sensitivity (n=140) model. All values should fall below 1.01 (dashed line).

Source: Time-to-Positivity and Mortality in Bacteremia

Moderator and sensitivity analyses

Table 9: All moderator and sensitivity analyses (primary pool). Δ log-OR is the change in pooled log-OR per unit of the moderator; negative values shrink the short-TTP harm signal.

Moderator	Δ log-OR	95% CrI	P(direction)
TTP cutpoint (per 10 h)	-0.355	[-0.561, -0.153]	99.9%
Pathogen: Gram+ vs Gram−	-0.500	[-1.016, 0.024]	97.0%
Adjustment: adjusted vs unadjusted	0.123	[-0.385, 0.624]	68.7%
Publication year (per 5 yr)	-0.203	[-0.414, 0.001]	97.5%

Source: Time-to-Positivity and Mortality in Bacteremia