

# Definition of Algorithm

---

Lin Chen

Email: [Lin.Chen@ttu.edu](mailto:Lin.Chen@ttu.edu)

Grader: [zulfi.khan@ttu.edu](mailto:zulfi.khan@ttu.edu)




TEXAS TECH  
UNIVERSITY.

# Algorithm

Algorithm is like a recipe

- It consists of finite number of instructions
- Each instruction is expressed by a finite number of symbols
- If it is carried out without error, it produces the same result in finite number of steps
- It demands no insight, intuition, or ingenuity, on the part of the one carrying out it



In principle, we are able to execute the code by hand, the machine just does that for us

# Algorithm

---

## Hilbert's 10<sup>th</sup> problem

Determine whether a given polynomial has an integral root by "*a process according to which it can be determined by a finite number of operations*"



# Algorithm

## Hilbert's 10<sup>th</sup> problem

Determine whether a given polynomial has an integral root by "*a process according to which it can be determined by a finite number of operations*"

Polynomials:

$$6x^3yz^2 + 3xy^2 - x^3 - 10$$

Root of a polynomials:

$$x = 5, y = 3, \text{ and } z = 0.$$

# Algorithm

---

## Hilbert's 10<sup>th</sup> problem

Determine whether a given polynomial has an integral root by "*a process according to which it can be determined by a finite number of operations*"

With an intuitive definition of an algorithm such a problem cannot be addressed

# Algorithm

## Hilbert's 10<sup>th</sup> problem

Determine whether a given polynomial has an integral root by "*a process according to which it can be determined by a finite number of operations*"

With an intuitive definition of an algorithm such a problem cannot be addressed

<i>Intuitive notion of algorithms</i>	equals	<i>Turing machine algorithms</i>
---	--------	--------------------------------------

No Turing machine can decide whether an arbitrary given polynomial has an integral root

# Algorithm

## Hilbert's 10<sup>th</sup> problem

Determine whether a given polynomial has an integral root by "*a process according to which it can be determined by a finite number of operations*"

$$D_1 = \{p \mid p \text{ is a polynomial over } x \text{ with an integral root}\}.$$

$M_1 =$  "The input is a polynomial  $p$  over the variable  $x$ .

1. Evaluate  $p$  with  $x$  set successively to the values 0, 1,  $-1$ , 2,  $-2$ , 3,  $-3$ ,  $\dots$ . If at any point the polynomial evaluates to 0, *accept*."

# Algorithm

## Hilbert's 10<sup>th</sup> problem

Determine whether a given polynomial has an integral root by "*a process according to which it can be determined by a finite number of operations*"

$$D_1 = \{p \mid p \text{ is a polynomial over } x \text{ with an integral root}\}.$$

$M_1$  = "The input is a polynomial  $p$  over the variable  $x$ .

1. Evaluate  $p$  with  $x$  set successively to the values 0, 1, -1, 2, -2, 3, -3, ... If at any point the polynomial evaluates to 0, *accept*."

TM  $M_1$  recognizes  $D_1$ . Does it decide  $D_1$



# Algorithm

## Hilbert's 10<sup>th</sup> problem

Determine whether a given polynomial has an integral root by "*a process according to which it can be determined by a finite number of operations*"

$$D_1 = \{p \mid p \text{ is a polynomial over } x \text{ with an integral root}\}.$$

$M_1$  = "The input is a polynomial  $p$  over the variable  $x$ .

1. Evaluate  $p$  with  $x$  set successively to the values 0, 1,  $-1$ , 2,  $-2$ , 3,  $-3$ ,  $\dots$ . If at any point the polynomial evaluates to 0, *accept*."

$M_1$  also decides  $D_1$  because we know when it must terminate: for single variable  $k$ -term polynomial with integral coefficients, the integral root has an absolute value at most  $kc_{max}$ , if it exists

# Algorithm

## Hilbert's 10<sup>th</sup> problem

Determine whether a given polynomial has an integral root by "*a process according to which it can be determined by a finite number of operations*"

For multi-variable polynomials, there is no such an upper bound, and a similar TM only recognizes.

$$x^3 + y^3 + z^3 = k$$

**Roger Heath-Brown conjecture (1992): every  $k$  unequal to 4 or 5 modulo 9 has infinitely many representations as the sum of three cubes**

Is it even true for  $k$  within 100? Unknown for 33 and 42 till 2016

# Algorithm

## Hilbert's 10<sup>th</sup> problem

Determine whether a given polynomial has an integral root by "*a process according to which it can be determined by a finite number of operations*"

For multi-variable polynomials, there is no such an upper bound, and a similar TM only recognizes.

$$x^3 + y^3 + z^3 = k$$

**Roger Heath-Brown conjecture (1992): every  $k$  unequal to 4 or 5 modulo 9 has infinitely many representations as the sum of three cubes**

Is it even true for  $k$  within 100? Unknown for 42 till 2019

# Algorithm

## Hilbert's 10<sup>th</sup> problem

Determine whether a given polynomial has an integral root by "*a process according to which it can be determined by a finite number of operations*"

For multi-variable polynomials, there is no such an upper bound, and a similar TM only recognizes.

$$x^3 + y^3 + z^3 = k$$

**Roger Heath-Brown conjecture (1992): every  $k$  unequal to 4 or 5 modulo 9 has infinitely many representations as the sum of three cubes**

Is it even true for  $k$  within 100? Unknown for 42 till 2019

A solution through supercomputer running for years:

$$(-80538738812075974)^3 + 80435758145817515^3 + 12602123297335631^3 = 42$$

# Algorithm

“I feel relieved,” Booker **said**. “In this game, it’s impossible to be sure that you’ll find something. It’s a bit like trying to predict earthquakes, in that we have only rough probabilities to go by. So, we might find what we’re looking for with a few months of searching, or it might be that the solution isn’t found for another century.”

For multi-variable polynomials, there is no such an upper bound, and a similar TM only recognizes.

$$x^3 + y^3 + z^3 = k$$

**Roger Heath-Brown conjecture (1992): every  $k$  unequal to 4 or 5 modulo 9 has infinitely many representations as the sum of three cubes**

Is it even true for  $k$  within 100? Unknown for 42 till 2019

A solution through supercomputer running for years:

$$(-80538738812075974)^3 + 80435758145817515^3 + 12602123297335631^3 = 42$$