




1 stgam: An R package for GAM-based varying 2 coefficient models

3 Lex Comber ^{1*}, Paul Harris ^{2*}, and Chris Brunsdon ^{3*}

4 1 School of Geography, University of Leeds, UK 2 Rothamsted Research, North Wyke, UK 3 National
5 Centre for Geocomputation, Maynooth University, Ireland * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright
and release the work under a
Creative Commons Attribution 4.0
International License ([CC BY 4.0](#))

6 Summary

7 We are often interested in understanding how and where statistical relationships vary over
8 space, and how they change over time. Quantifying such *process heterogeneity* (spatial and or
9 temporal) can be done using *varying coefficient* models. The opportunity to undertake such
10 space-time analyses are greater due to the increased generation and availability of data that
11 include both spatial and temporal attributes (e.g. GPS coordinates and time-stamps). The
12 **stgam** package provides a framework for creating regression models using Generalized Additive
13 Models (GAMs) ([Hastie & Tibshirani, 1990](#)) in which the relationships between the response
14 (dependent) variable and individual predictor (independent) variables are allowed to vary over
15 space, time or both.

Reference

Hastie, T., & Tibshirani, R. (1990). Generalized additive models. Chapman hall & CRC.
Monographs on Statistics & Applied Probability. Chapman and Hall/CRC, 1.