

Geometry¹

First Year, Computer science

Assoc. Prof. Cornel-Sebastian PINTEA

“Babeş-Bolyai” University
Faculty of Mathematics and Computer Sciences
Cluj-Napoca, Romania

Lecture 12, 28.05.2014

¹These notes are not in a final form. They are continuously being improved

Transformations

Transformations of the
plane

Translations

Scaling about the origine

Reflections

Rotations

Transformations

Transformations of the plane

Translations

Scaling about the origine

Reflections

Rotations

Definition 1.1

An affine transformation of the plane is a mapping

$$L : \mathbb{R}^2 \longrightarrow \mathbb{R}^2, L(x, y) = (ax + by + c, dx + ey + f), \quad (1.1)$$

for some constant real numbers a, b, c, d, e, f .

By using the matrix language, the action of the map L can be written in the form

$$L(x, y) = [x \ y] \begin{bmatrix} a & d \\ b & e \end{bmatrix} + [c \ f].$$

The affine transformation L can be also identified with the map $L^c : \mathbb{R}^{2 \times 1} \longrightarrow \mathbb{R}^{2 \times 1}$ given by

$$\begin{aligned} L^c \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) &= \begin{bmatrix} ax + by + c \\ dx + ey + f \end{bmatrix} = \begin{bmatrix} a & b \\ d & e \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} c \\ f \end{bmatrix} \\ &= [L] \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} c \\ f \end{bmatrix}, \text{ where } [L] = \begin{bmatrix} a & b \\ d & e \end{bmatrix}. \end{aligned}$$

Lemma 1.1

If $(aB - bA)^2 + (dB - eA)^2 > 0$, then the affine transformation (1.1) maps the line (d) $Ax + By + C = 0$ to the line

$$(eA - dB)x + (aB - bA)y + (bf - ce)A - (af - cd)B + (ae - bd)C = 0.$$

If $aB - bA = dB - eA = 0$, then $ae - bd = 0$ and L is the constant map $\left(\frac{cB - bC}{B}, \frac{fB - eC}{B}\right)$.

Definition 1.3

An affine transformation (1.1) is said to be singular if

$$\begin{vmatrix} a & b \\ d & e \end{vmatrix} = 0 \text{ i.e. } ae - bd = 0.$$

and non-singular otherwise.

Transformations

Transformations of the
plane

Translations

Scaling about the origine

Reflections

Rotations

Note that the affine transformation L is nonsingular if and only if it is invertible. In such a case the inverse L^{-1} is a non-singular affine transformation and $[L^{-1}] = [L]^{-1}$.

Definition 1.4

The translation of vector $(h, k) \in \mathbb{R}^2$ is the affine transformation

$$T(h, k) : \mathbb{R}^2 \longrightarrow \mathbb{R}^2, [T(h, k)](x, y) = (x + h, y + k).$$

Thus

$$[T(h, k)]^c \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} x + h \\ y + k \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} h \\ k \end{bmatrix},$$

i.e.

$$[T(h, k)] = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Note that the translation $T(h, k)$ is non-singular (invertible) and $(T(h, k))^{-1} = T(-h, -k)$.

Transformations

Transformations of the plane

Translations

Scaling about the origine

Reflections

Rotations

Transformations

Transformations of the
plane

Translations

Scaling about the origine

Reflections

Rotations

Definition 1.5

The scaling about the origine by non-zero scaling factors $(s_x, s_y) \in \mathbb{R}^2$ is the affine transformation

$$S(s_x, s_y) : \mathbb{R}^2 \longrightarrow \mathbb{R}^2, [S(s_x, s_y)](x, y) = (s_x \cdot x, s_y \cdot y).$$

Thus

$$[S(s_x, s_y)]^c \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} s_x \cdot x \\ s_y \cdot y \end{bmatrix} = \begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix},$$

i.e.

$$[S(s_x, s_y)] = \begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix}.$$

Note that the scaling about the origine by non-zero scaling factors $(s_x, s_y) \in \mathbb{R}^2$ is non-singular (invertible) and $(S(s_x, s_y))^{-1} = S(s_x^{-1}, s_y^{-1})$.

Definition 1.6

The reflections about the x -axis and the y -axis respectively are the affine transformation

$$S_x, S_y : \mathbb{R}^2 \longrightarrow \mathbb{R}^2, \quad S_x(x, y) = (x, -y), \quad S_y = (-x, y).$$

Thus

$$[S_x^c] \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} x \\ -y \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix},$$

i.e.

$$[S_x] = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}. \text{ Similarly } [S_y] = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Note that $S_x = S(-1, 1)$ and $S_y = S(1, -1)$. Thus the two reflections are non-singular (invertible) and $S_x^{-1} = S_x$, $S_y^{-1} = S_y$.

Transformations

Transformations of the
plane

Translations

Scaling about the origine

Reflections

Rotations

Definition 1.7

The reflection $S_l : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ about the line l maps a given point M to the point M' defined by the property that l is the perpendicular bisector of the segment MM' . One can show that the action of the reflection about the line $l : ax + by + c = 0$ is

$$S_l(x, y) = \left(\frac{b^2 - a^2}{a^2 + b^2}x - \frac{2ab}{a^2 + b^2}y - \frac{2ac}{a^2 + b^2}, -\frac{2ab}{a^2 + b^2}x + \frac{a^2 - b^2}{a^2 + b^2}y - \frac{2bc}{a^2 + b^2} \right)$$

$$\begin{aligned} \text{Thus } [S_l^c] \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) &= \begin{bmatrix} \frac{b^2 - a^2}{a^2 + b^2}x - \frac{2ab}{a^2 + b^2}y - \frac{2ac}{a^2 + b^2} \\ -\frac{2ab}{a^2 + b^2}x + \frac{a^2 - b^2}{a^2 + b^2}y - \frac{2bc}{a^2 + b^2} \end{bmatrix} \\ &= \begin{bmatrix} \frac{b^2 - a^2}{a^2 + b^2} & -\frac{2ab}{a^2 + b^2} \\ -\frac{2ab}{a^2 + b^2} & \frac{a^2 - b^2}{a^2 + b^2} \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} - \begin{bmatrix} \frac{2ac}{a^2 + b^2} \\ \frac{2bc}{a^2 + b^2} \end{bmatrix}, \end{aligned}$$

i.e. $[S_l] = \frac{1}{a^2 + b^2} \begin{bmatrix} b^2 - a^2 & -2ab \\ -2ab & a^2 - b^2 \end{bmatrix}$. Note that the reflection S_l is non-singular (invertible) and $S_l^{-1} = S_l$.

Definition 1.8

The rotation $R_\theta : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ about the origine through an angle θ maps a point $M(x, y)$ into a point $M'(x', y')$ with the properties that the segments $[OM]$ and $[OM']$ are congruent and the $m(\widehat{MOM'}) = \theta$. If $\theta > 0$ the rotation is supposed to be anticlockwise and for $\theta < 0$ the rotation is clockwise. If $(x, y) = (r \cos \varphi, r \sin \varphi)$, then the coordinates of the rotated point are $(r \cos(\theta + \varphi), r \sin(\theta + \varphi)) = (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta)$, i.e. $R_\theta(x, y) = (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta)$.

$$\begin{aligned} \text{Thus } [R_\theta^c] \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) &= \begin{bmatrix} x \cos \theta - y \sin \theta \\ x \sin \theta + y \cos \theta \end{bmatrix} \\ &= \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}, \end{aligned}$$

i.e. $[R_\theta] = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$. Note that the rotation R_θ is non-singular (invertible) and $R_\theta^{-1} = R_{-\theta}$.