

# Geometry<sup>1</sup>

## First Year, Computer science

Assoc. Prof. Cornel-Sebastian PINTEA

“Babeş-Bolyai” University  
Faculty of Mathematics and Computer Sciences  
Cluj-Napoca, Romania

Lecture 1, 26.02.2014

---

<sup>1</sup>These notes are not in a final form. They are continuously being improved

## Vectors and their vector structure

Vectors

Operations with vectors

The addition of vectors

The multiplication of  
vectors with scalars

The vector structure on the  
set of vectors

## Vectors and their vector structure

### Vectors

### Operations with vectors

The addition of vectors

The multiplication of vectors with scalars

The vector structure on the set of vectors

# Vectors and their vector structure. Vectors


Cursul 1

Conf. Dr. PINTEA  
Cornel-Sebastian

Let  $\mathcal{P}$  be the three dimensional physical space in which we can talk about points, lines, planes and various relations among them. If  $(A, B) \in \mathcal{P} \times \mathcal{P}$  is an ordered pair, then  $A$  is called the *original point* or the *origin* and  $B$  is called the *terminal point* or the *extremity* of  $(A, B)$ .

## Definition 1.1

The ordered pairs  $(A, B)$ ,  $(C, D)$  are said to be *equipollent*, written  $(A, B) \sim (C, D)$ , if the segments  $[AD]$  and  $[BC]$  have the same midpoint.



Vectors and their  
vector structure

Vectors

Operations with vectors

The addition of vectors

The multiplication of

vectors with scalars

The vector structure on the  
set of vectors

## Remark 1.2

If  $A, B, C, D$  are not collinear points, then  $(A, B) \sim (C, D)$  if and only if  $ABDC$  is a parallelogram. In fact the length of the segments  $[AB]$  and  $[CD]$  is the same whenever  $(A, B) \sim (C, D)$ .

## Proposition 1.3

*If  $(A, B)$  is an ordered pair and  $O \in \mathcal{P}$  is a given point, then there exists a unique point  $X$  such that  $(A, B) \sim (O, X)$ .*

## Proposition 1.4

*The equipollence relation is an equivalence relation on  $\mathcal{P} \times \mathcal{P}$ .*

Vectors and their  
vector structure

Vectors

Operations with vectors

The addition of vectors

The multiplication of  
vectors with scalarsThe vector structure on the  
set of vectors

## Definition 1.5

*The equivalence classes with respect to the equipollence relation are called (free) vectors.*

Denote by  $\overrightarrow{AB}$  the equivalence class of the ordered pair  $(A, B)$ , that is  $\overrightarrow{AB} = \{(X, Y) \in \mathcal{P} \times \mathcal{P} \mid (X, Y) \sim (A, B)\}$  and let  $\mathcal{V} = \mathcal{P} \times \mathcal{P} / \sim = \{\overrightarrow{AB} \mid (A, B) \in \mathcal{P} \times \mathcal{P}\}$  be the set of (free) vectors. The *length* or the *magnitude* of the vector  $\overrightarrow{AB}$ , denoted by  $\|\overrightarrow{AB}\|$  or by  $|\overrightarrow{AB}|$ , is the length of the segment  $[AB]$ .

## Remark 1.6

*If two ordered pairs  $(A, B)$  and  $(C, D)$  are equipollent, i.e. the vectors  $\overrightarrow{AB}$  and  $\overrightarrow{CD}$  are equal, then they have the same length, the same direction and the same sense. In fact a vector is determined by these three items.*

Vectors and their  
vector structure

Vectors

Operations with vectors

The addition of vectors

The multiplication of  
vectors with scalars

The vector structure on the  
set of vectors



## Proposition 1.7

1.  $\overrightarrow{AB} = \overrightarrow{CD} \Leftrightarrow \overrightarrow{AC} = \overrightarrow{BD}$ .
2.  $\forall A, B, O \in \mathcal{P}, \exists ! X \in \mathcal{P}$  such that  $\overrightarrow{AB} = \overrightarrow{OX}$ .
3.  $\overrightarrow{AB} = \overrightarrow{A'B'}, \overrightarrow{BC} = \overrightarrow{B'C'} \Rightarrow \overrightarrow{AC} = \overrightarrow{A'C'}$ .

## Definition 1.8

If  $O, M \in \mathcal{P}$ , the the vector  $\overrightarrow{OM}$  is denoted by  $\vec{r}_M$  and is called the *position vector of  $M$  with respect to  $O$* .

## Corollary 1.9

The map  $\varphi_O : \mathcal{P} \rightarrow \mathcal{V}$ ,  $\varphi_O(M) = \vec{r}_M$  is one-to-one and onto, i.e bijective.

Vectors and their  
vector structure

Vectors

Operations with vectors

The addition of vectors

The multiplication of  
vectors with scalars

The vector structure on the  
set of vectors

## Vectors and their vector structure

### Vectors

#### Operations with vectors

##### The addition of vectors

##### The multiplication of vectors with scalars

##### The vector structure on the set of vectors

In this section we shall define and study the the addition operation of vectors alongside the multiplication of vectors with scalars.

# The addition of vectors

Cursul 1

Conf. Dr. PINTEA  
Cornel-Sebastian

Let  $\vec{a}, \vec{b} \in \mathcal{V}$  and  $O \in \mathcal{P}$  be such that  $\vec{a} = \overrightarrow{OA}$ ,  $\vec{b} = \overrightarrow{OB}$ . The vector  $\overrightarrow{OB}$  is called the *sum* of the vectors  $\vec{a}$  and  $\vec{b}$  and is written  $\overrightarrow{OB} = \overrightarrow{OA} + \overrightarrow{AB} = \vec{a} + \vec{b}$ .




Figure: 1

Vectors and their  
vector structure

Vectors

Operations with vectors

The addition of vectors

The multiplication of  
vectors with scalars

The vector structure on the  
set of vectors



# The addition of vectors

Cursul 1

Conf. Dr. PINTEA  
Cornel-Sebastian

Let  $O'$  be another point and  $A', B' \in \mathcal{P}$  be such that  $\overrightarrow{O'A'} = \vec{a}$ ,  $\overrightarrow{A'B'} = \vec{b}$ . Since  $\overrightarrow{OA} = \overrightarrow{O'A'}$  and  $\overrightarrow{AB} = \overrightarrow{A'B'}$  it follows, according to Proposition 1.4 (3), that  $\overrightarrow{OB} = \overrightarrow{O'B'}$ . Therefore the vector  $\vec{a} + \vec{b}$  is independent on the choice of the point  $O$ .

## Proposition 2.1

*The set  $\mathcal{V}$  endowed to the binary operation  $\mathcal{V} \times \mathcal{V} \rightarrow \mathcal{V}$ ,  $(\vec{a}, \vec{b}) \mapsto \vec{a} + \vec{b}$ , is an abelian group whose zero element is the vector  $\overrightarrow{AA} = \overrightarrow{BB} = \vec{0}$  and the opposite of  $\overrightarrow{AB}$ , denoted by  $-\overrightarrow{AB}$ , is the vector  $\overrightarrow{BA}$ .*

In particular, then addition operation is associative, the vector  $(\vec{a} + \vec{b}) + \vec{c} = \vec{a} + (\vec{b} + \vec{c})$  is usually denoted by  $\vec{a} + \vec{b} + \vec{c}$ . Moreover the expression

$$((\cdots(\vec{a}_1 + \vec{a}_2) + \vec{a}_3 + \cdots + \vec{a}_n) \cdots), \quad (1.1)$$

Vectors and their  
vector structure

Vectors

Operations with vectors

The addition of vectors

The multiplication of  
vectors with scalars

The vector structure on the  
set of vectors

# The addition of vectors

Cursul 1

Conf. Dr. PINTEA  
Cornel-Sebastian

is independent on the distribution of paranthesis and it is usually denoted by  $\vec{a}_1 + \vec{a}_2 + \cdots + \vec{a}_n$ .

## Example 2.2

*If  $A_1, A_2, A_3, \dots, A_n \in \mathcal{P}$  are some given points, then  $\vec{A_1A_2} + \vec{A_2A_3} + \cdots + \vec{A_{n-1}A_n} = \vec{A_1A_n}$ . This shows that  $\vec{A_1A_2} + \vec{A_2A_3} + \cdots + \vec{A_{n-1}A_n} + \vec{A_nA_1} = \vec{0}$ , namely the sum of vectors constructed on the edges of a closed broken line is zero.*

## Example 2.3

*Consider the parallelograms in  $\mathcal{P}$ ,  $A_1A_2A_3A_4$ ,  $B_1B_2B_3B_4$ , and  $M_1, M_2, M_3, M_4$  the midpoints of the segments  $[A_1B_1]$ ,  $[A_2B_2]$ ,  $[A_3B_3]$ ,  $[A_4B_4]$  respectively. Show that:*

- ▶  $2 \vec{M_1M_2} = \vec{A_1A_2} + \vec{B_1B_2}$  and  $2 \vec{M_3M_4} = \vec{A_3A_4} + \vec{B_3B_4}$ .
- ▶  $M_1, M_2, M_3, M_4$  are the vertices of a parallelogram.

Vectors and their  
vector structure

Vectors

Operations with vectors

The addition of vectors

The multiplication of  
vectors with scalars

The vector structure on the  
set of vectors



# The addition of vectors

Cursul 1

Conf. Dr. PINTEA  
Cornel-Sebastian

## Corolarul 2.4

If  $\vec{a} = \overrightarrow{OA}$ ,  $\vec{b} = \overrightarrow{OB}$  are given vectors, there exists a unique vector  $\vec{x} \in \mathcal{V}$  such that  $\vec{a} + \vec{x} = \vec{b}$ . In fact  $\vec{x} = \vec{b} + (-\vec{a}) = \overrightarrow{AB}$  and is denoted by  $\vec{b} - \vec{a}$ .




Figure: 1

Vectors and their  
vector structure

Vectors

Operations with vectors

The addition of vectors

The multiplication of  
vectors with scalars

The vector structure on the  
set of vectors

# The multiplication of vectors with scalars

Cursul 1

Conf. Dr. PINTEA  
Cornel-Sebastian

Let  $\alpha \in \mathbb{R}$  be a scalar and  $\vec{a} = \overrightarrow{OA} \in \mathcal{V}$  be a vector. We define the vector  $\alpha \cdot \vec{a}$  as follows:  $\alpha \cdot \vec{a} = \vec{0}$  if  $\alpha = 0$  or  $\vec{a} = \vec{0}$ ; if  $\vec{a} \neq \vec{0}$  and  $\alpha > 0$ , there exists a unique point on the half line  $]OA$  such that  $\|OB\| = \alpha \cdot \|OA\|$  and define  $\alpha \cdot \vec{a} = \overrightarrow{OB}$ ; if  $\alpha < 0$  we define  $\alpha \cdot \vec{a} = -(|\alpha| \cdot \vec{a})$ . The external binary operation

$$\mathbb{R} \times \mathcal{V} \rightarrow \mathcal{V}, (\alpha, \vec{a}) \mapsto \alpha \cdot \vec{a}$$

is called the *multiplication of vectors with scalars*.

## Proposition 2.5

*The following properties hold:*

$$(v1) \quad (\alpha + \beta) \cdot \vec{a} = \alpha \cdot \vec{a} + \beta \cdot \vec{a}, \quad \forall \alpha, \beta \in \mathbb{R}, \vec{a} \in \mathcal{V}.$$

$$(v2) \quad \alpha \cdot (\vec{a} + \vec{b}) = \alpha \cdot \vec{a} + \alpha \cdot \vec{b}, \quad \forall \alpha \in \mathbb{R}, \vec{a}, \vec{b} \in \mathcal{V}.$$

$$(v3) \quad \alpha \cdot (\beta \cdot \vec{a}) = (\alpha\beta) \cdot \vec{a}, \quad \forall \alpha, \beta \in \mathbb{R}.$$

$$(v4) \quad 1 \cdot \vec{a} = \vec{a}, \quad \forall \vec{a} \in \mathcal{V}.$$

Vectors and their  
vector structure

Vectors

Operations with vectors

The addition of vectors

The multiplication of  
vectors with scalars

The vector structure on the  
set of vectors



# The vector structure on the set of vectors

Cursul 1

Conf. Dr. PINTEA  
Cornel-Sebastian

## Theorem 2.6

*The set of (free) vectors endowed with the addition binary operation of vectors and the external binary operation of multiplication of vectors with scalars is a real vector space.*

Vectors and their  
vector structure

Vectors

Operations with vectors

The addition of vectors

The multiplication of  
vectors with scalars

The vector structure on the  
set of vectors

## Example 2.7

*If  $A'$  is the midpoint of the edge  $[BC]$  of the triangle  $ABC$ , then  $\overrightarrow{AA'} = \frac{1}{2}(\overrightarrow{AB} + \overrightarrow{AC})$ .*

## Example 2.8

*If  $O$  is a fixed point in  $\mathcal{P}$ , then the position vector  $\vec{r}_G = \overrightarrow{OG}$  of the centroid  $G$  with respect to the point  $O$  is  $\vec{r}_G = \frac{1}{3}(\vec{r}_A + \vec{r}_B + \vec{r}_C)$ .*