

Geometry¹

First Year, Computer science

Assoc. Prof. Cornel-Sebastian PINTEA

“Babeş-Bolyai” University
Faculty of Mathematics and Computer Sciences
Cluj-Napoca, Romania

Lecture 3, 12.03.2014

¹These notes are not in a final form. They are continuously being improved

Cartesian and affine reference systems

The cartesian equations of the straight lines and planes

Analytic conditions of parallelism

Pencils of planes

Cartesian and
affine reference
systems

The cartesian
equations of the
straight lines and
planes

Analytic conditions
of parallelism

Pencils of planes

Cartesian and affine reference systems

Lecture 3

Assoc. Prof.
Cornel-Sebastian
PINTEA

Cartesian and
affine reference
systems

The cartesian
equations of the
straight lines and
planes

Analytic conditions
of parallelism

Pencils of planes

A basis of the direction $\vec{\pi}$ of the plane π , or for the vector space \mathcal{V} is an ordered basis $[\vec{e}, \vec{f}]$ of π , or an ordered basis $[\vec{u}, \vec{v}, \vec{w}]$ of \mathcal{V} .

If $b = [\vec{u}, \vec{v}, \vec{w}]$ is a basis of \mathcal{V} and $\vec{x} \in \mathcal{V}$, recall that the column vector of \vec{x} with respect to b is being denoted by $[\vec{x}]_b$. In other words

$$[\vec{x}]_b = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}.$$

whenever $\vec{x} = x_1\vec{u} + x_2\vec{v} + x_3\vec{w}$.

Definition 1.1

A cartesian reference system of the space \mathcal{P} , is a system $R = (O, \vec{u}, \vec{v}, \vec{w})$ where O is a point from \mathcal{P} called the origin of the reference system and $b = [\vec{u}, \vec{v}, \vec{w}]$ is a basis of the vector space \mathcal{V} .

Denote by E_1, E_2, E_3 the points for which $\vec{u} = \overrightarrow{OE_1}$, $\vec{v} = \overrightarrow{OE_2}$, $\vec{w} = \overrightarrow{OE_3}$.

Definition 1.2

The system of points (O, E_1, E_2, E_3) is called the affine reference system associated to the cartesian reference system $R = (O, \vec{u}, \vec{v}, \vec{w})$.

The straight lines OE_i , $i \in \{1, 2, 3\}$, oriented from O to E_i are called *the coordinate axes*. The coordinates x, y, z of the position vector $\vec{r}_M = \overrightarrow{OM}$ with respect to the basis $[\vec{u}, \vec{v}, \vec{w}]$ are called the coordinates of the point M with respect to the cartesian system R written $M(x, y, z)$.

Also, for the column matrix of coordinates of the vector \vec{r}_M we are going to use the notation $[M]_R$. In other words, if $\vec{r}_M = x\vec{u} + y\vec{v} + z\vec{w}$, then

$$[M]_R = [\overrightarrow{OM}]_b = \begin{pmatrix} x \\ y \\ z \end{pmatrix}.$$

Remark 1.3

If $A(x_A, y_A, z_A)$, $B(x_B, y_B, z_B)$ are two points, then

$$\begin{aligned} \overrightarrow{AB} &= \overrightarrow{OB} - \overrightarrow{OA} \\ &= x_B\vec{u} + y_B\vec{v} + z_B\vec{w} - (x_A\vec{u} + y_A\vec{v} + z_A\vec{w}) \\ &= (x_B - x_A)\vec{u} + (y_B - y_A)\vec{v} + (z_B - z_A)\vec{w}, \end{aligned}$$

i.e. the coordinates of the vector \overrightarrow{AB} are being obtained by performing the differences of the coordinates of the points A and B .

The cartesian equations of the straight lines and planes

Let Δ be a straight line passing through the point $A_0(x_0, y_0, z_0)$ which is parallel to the vector $\vec{d}(p, q, r)$. Its vector equation is

$$\vec{r}_M = \vec{r}_{A_0} + \lambda \vec{d}. \quad (2.1)$$

Denoting by x, y, z the coordinates of the generic point M of the straight line Δ , its vector equation (2.1) is equivalent to the following system of relations

$$\begin{cases} x = x_0 + \lambda p \\ y = y_0 + \lambda q \\ z = z_0 + \lambda r \end{cases}, \lambda \in \mathbb{R} \quad (2.2)$$

The relations (2.2) are being called the *parametric equations* of the straight line Δ and they are equivalent to the following relations

$$\frac{x - x_0}{p} = \frac{y - y_0}{q} = \frac{z - z_0}{r} \quad (2.3)$$

If $r = 0$, for instance, the canonical equations of the straight line Δ are

$$\frac{x - x_0}{p} = \frac{y - y_0}{q} \wedge z = z_0.$$

If $A(x_A, y_A, z_A)$, $B(x_B, y_B, z_B)$ are different points of the straight line Δ , then $\overrightarrow{AB} (x_B - x_A, y_B - y_A, z_B - z_A)$ is a director vector of Δ , its canonical equations having, in this case, the form

$$\frac{x - x_A}{x_B - x_A} = \frac{y - y_A}{y_B - y_A} = \frac{z - z_A}{z_B - z_A}. \quad (2.4)$$

Cartesian and
affine reference
systems

The cartesian
equations of the
straight lines and
planes

Analytic conditions
of parallelism

Pencils of planes

Te cartesian equations of the planes

Lecture 3

Assoc. Prof.
Cornel-Sebastian
PINTEA

Let $A_0(x_0, y_0, z_0) \in \mathcal{P}$ and $\vec{d}_1(p_1, q_1, r_1), \vec{d}_2(p_2, q_2, r_2) \in \mathcal{V}$ be linearly independent vectors, that is

$$\text{rang} \begin{pmatrix} p_1 & q_1 & r_1 \\ p_2 & q_2 & r_2 \end{pmatrix} = 2.$$

The vector equation of the plane π passing through A_0 which is parallel to the vectors $\vec{d}_1(p_1, q_1, r_1), \vec{d}_2(p_2, q_2, r_2)$ is

$$\vec{r}_M = \vec{r}_{A_0} + \lambda_1 \vec{d}_1 + \lambda_2 \vec{d}_2, \quad \lambda_1, \lambda_2 \in \mathbb{R}. \quad (2.5)$$

If we denote by x, y, z the coordinates of the generic point M of the plane π , then the vector equation (2.5) is the equivalent to the following system of relations

$$\begin{cases} x = x_0 + \lambda_1 p_1 + \lambda_2 p_2 \\ y = y_0 + \lambda_1 q_1 + \lambda_2 q_2 \\ z = z_0 + \lambda_1 r_1 + \lambda_2 r_2 \end{cases}, \quad \lambda_1, \lambda_2 \in \mathbb{R}. \quad (2.6)$$

Cartesian and
affine reference
systems

The cartesian
equations of the
straight lines and
planes

Analytic conditions
of parallelism

Pencils of planes

The relations (2.6) represent a characterization of the points of the plane π called the *parametric equations* of the plane π . More precisely, the compatibility of the linear system (2.6) with the unknowns λ_1, λ_2 is a necessary and sufficient condition for the point $M(x, y, z)$ to be contained within the plane π . On the other hand the compatibility of the linear system (2.6) is equivalent to the relations

$$\begin{vmatrix} x - x_0 & y - y_0 & z - z_0 \\ p_1 & q_1 & z_1 \\ p_2 & q_2 & z_2 \end{vmatrix} = 0. \quad (2.7)$$

and express the fact that the rank of the matrix of the system is equal to the rank of the extended matrix of the system. The condition (2.7) is a characterization of the points of the plane π expressed in terms of the cartesian coordinates of the generic point M and is called the *cartesian equation* of the plane π .

If $A(x_A, y_A, z_A)$, $B(x_B, y_B, z_B)$, $C(x_C, y_C, z_C)$ are noncollinear points, then the plane (ABC) determined by the three points can be viewed as the plane passing through the point A which is parallel to the vectors $\vec{d}_1 = \overrightarrow{AB}$, $\vec{d}_2 = \overrightarrow{AC}$. The coordinates of the vectors \vec{d}_1 și \vec{d}_2 are

$(x_B - x_A, y_B - y_A, z_B - z_A)$ and $(x_C - x_A, y_C - y_A, z_C - z_A)$ respectively.

Thus, the equation of the plane (ABC) is

$$\begin{vmatrix} x - x_A & y - y_A & z - z_A \\ x_B - x_A & y_B - y_A & z_B - z_A \\ x_C - x_A & y_C - y_A & z_C - z_A \end{vmatrix} = 0, \quad (2.8)$$

or, echivalently

$$\begin{vmatrix} x & y & z & 1 \\ x_A & y_A & z_A & 1 \\ x_B & y_B & z_B & 1 \\ x_C & y_C & z_C & 1 \end{vmatrix} = 0. \quad (2.9)$$

One can put the equation (2.7) in the form

$$A(x - x_0) + B(y - y_0) + C(z - z_0) = 0 \text{ or} \quad (2.10)$$

$$Ax + By + Cz + D = 0, \quad (2.11)$$

where the coefficients A, B, C satisfy the relation $A^2 + B^2 + C^2 > 0$. It is also easy to show that every equation of the form (2.11) represents the equation of a plane. Indeed, if $A \neq 0$, then the equation (2.11) is equivalent to

$$\begin{vmatrix} x + \frac{D}{A} & y & z \\ B & -A & 0 \\ C & 0 & -A \end{vmatrix} = 0.$$

We observe that one can put the equation (2.10) in the form

$$AX + BY + CZ = 0 \quad (2.12)$$

where $X = x - x_0$, $Y = y - y_0$, $Z = z - z_0$ are the coordinates of the vector $\overrightarrow{A_0M}$.

Cartesian and
affine reference
systems

The cartesian
equations of the
straight lines and
planes

Analytic conditions
of parallelism

Pencils of planes

Analytic conditions of parallelism

Lecture 3

Assoc. Prof.
Cornel-Sebastian
PINTEA

Cartesian and
affine reference
systems

The cartesian
equations of the
straight lines and
planes

Analytic conditions
of parallelism

Pencils of planes

The equation $AX + BY + CZ = 0$ is a necessary and sufficient condition for the vector $\overrightarrow{A_0M}(X, Y, Z)$ to be contained within the direction of the plane

$$\pi : A(x - x_0) + B(y - y_0) + C(z - z_0) = 0.$$

Thus the *equation of the director subspace*

$$\vec{\pi} = \{\overrightarrow{A_0M} \mid M \in \pi\} \text{ is } AX + BY + CZ = 0.$$

Proposition 3.1

The straight line

$$\Delta : \frac{x - x_0}{p} = \frac{y - y_0}{q} = \frac{z - z_0}{r}$$

is parallel to the plane $\pi : Ax + By + Cz + D = 0$ iff

$$Ap + Bq + Cr = 0 \tag{3.1}$$



Proposition 3.2

Consider the planes $(\pi_1) A_1x + B_1y + C_1z + D_1 = 0$, $(\pi_2) A_2x + B_2y + C_2z + D_2 = 0$. Then $\dim(\vec{\pi}_1 \cap \vec{\pi}_2) \in \{1, 2\}$ and the following statements are equivalent

1. $\pi_1 \parallel \pi_2$.
2. $\dim(\vec{\pi}_1 \cap \vec{\pi}_2) = 2$, i.e. $\vec{\pi}_1 = \vec{\pi}_2$.
3. $\text{rang} \begin{pmatrix} A_1 & B_1 & C_1 \\ A_2 & B_2 & C_2 \end{pmatrix} = 1$.
4. The vectors $(A_1, B_1, C_1), (A_2, B_2, C_2) \in \mathbb{R}^3$ are linearly dependent.

Cartesian and
affine reference
systems

The cartesian
equations of the
straight lines and
planes

Analytic conditions
of parallelism

Pencils of planes

Corollary 3.3

Consider the planes $(\pi_1) A_1x + B_1y + C_1z + D_1 = 0$, $(\pi_2) A_2x + B_2y + C_2z + D_2 = 0$. The following statements are equivalent

1. $\pi_1 \nparallel \pi_2$.
2. $\dim(\vec{\pi}_1 \cap \vec{\pi}_2) = 1$.
3. $\text{rang} \begin{pmatrix} A_1 & B_1 & C_1 \\ A_2 & B_2 & C_2 \end{pmatrix} = 2$.
4. *The vectors (A_1, B_1, C_1) , $(A_2, B_2, C_2) \in \mathbb{R}^3$ are linearly independent.*

By using Proposition (3.1) we shall find a necessary and sufficient condition for a vector to be contained within the direction of a straight line which is given as the intersection of two planes.

Consider the planes $(\pi_1) A_1x + B_1y + C_1z + D_1 = 0$,
 $(\pi_2) A_2x + B_2y + C_2z + D_2 = 0$ such that

$$\text{rang} \begin{pmatrix} A_1 & B_1 & C_1 \\ A_2 & B_2 & C_2 \end{pmatrix} = 2,$$

alongside their intersection straight line $\Delta = \pi_1 \cap \pi_2$ of equations

$$(\Delta) \begin{cases} A_1x + B_1y + C_1z + D_1 = 0 \\ A_2x + B_2y + C_2z + D_1 = 0. \end{cases}$$

Thus, $\vec{\Delta} = \vec{\pi}_1 \cap \vec{\pi}_2$ and therefore, by means of Proposition (3.1), it follows that the equations of $\vec{\Delta}$ are

$$(\vec{\Delta}) \begin{cases} A_1X + B_1Y + C_1Z = 0 \\ A_2X + B_2Y + C_2Z = 0. \end{cases} \quad (3.2)$$

Cartesian and
affine reference
systems

The cartesian
equations of the
straight lines and
planes

Analytic conditions
of parallelism

Pencils of planes

By solving the system (3.2) one can therefore deduce that $\vec{d}(p, q, r) \in \vec{\Delta} \Leftrightarrow \exists \lambda \in \mathbb{R}$ such that

$$(p, q, r) = \lambda \left(\left| \begin{array}{cc} B_1 & C_1 \\ B_2 & C_2 \end{array} \right|, \left| \begin{array}{cc} C_1 & A_1 \\ C_2 & A_2 \end{array} \right|, \left| \begin{array}{cc} A_1 & B_1 \\ A_2 & B_2 \end{array} \right| \right). \quad (3.3)$$

The relation is usually (3.3) written in the form

$$\overline{\left| \begin{array}{cc} B_1 & C_1 \\ B_2 & C_2 \end{array} \right|} = \overline{\left| \begin{array}{cc} C_1 & A_1 \\ C_2 & A_2 \end{array} \right|} = \overline{\left| \begin{array}{cc} A_1 & B_1 \\ A_2 & B_2 \end{array} \right|}. \quad (3.4)$$

Let us mention that the chosen values for (p, q, r) are usually precisely

$$\left| \begin{array}{cc} B_1 & C_1 \\ B_2 & C_2 \end{array} \right|, \left| \begin{array}{cc} C_1 & A_1 \\ C_2 & A_2 \end{array} \right| \text{ și } \left| \begin{array}{cc} A_1 & B_1 \\ A_2 & B_2 \end{array} \right|.$$

Cartesian and
affine reference
systems

The cartesian
equations of the
straight lines and
planes

Analytic conditions
of parallelism

Pencils of planes

Definition 4.1

The collection of all planes containing a given straight line

$$(\Delta) \begin{cases} A_1x + B_1y + C_1z + D_1 = 0 \\ A_2x + B_2y + C_2z + D_2 = 0 \end{cases}$$

is called the pencil of planes through Δ .

Proposition 4.2

The plane π belongs to the pencil of planes through the straight line Δ if and only if there exists $\lambda, \mu \in \mathbb{R}$ such that the equation of the plane π is

$$\lambda(A_1x + B_1y + C_1z + D_1) + \mu(A_2x + B_2y + C_2z + D_2) = 0. \quad (4.1)$$

Cartesian and
affine reference
systems

The cartesian
equations of the
straight lines and
planes

Analytic conditions
of parallelism

Pencils of planes

Remark 4.3

The family of planes

$$A_1x + B_1y + C_1z + D_1 + \lambda(A_2x + B_2y + C_2z + D_2) = 0,$$

where λ covers the whole real line \mathbb{R} , is the so called reduced pencil of planes through Δ and it consists in all planes through Δ except the plane of equation $A_2x + B_2y + C_2z + D_2 = 0$.