

Geometry¹

First Year, Computer science

Assoc. Prof. Cornel-Sebastian PINTEA

“Babeş-Bolyai” University
Faculty of Mathematics and Computer Sciences
Cluj-Napoca, Romania

Lecture 9, 07.05.2014

¹These notes are not in a final form. They are continuously being improved

Conics

The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets

Conics

The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets

Definiția 1.1

The parabola is a plane curve defined to be the geometric locus of the points in the plane, whose distance to a fixed line d is equal to its distance to a fixed point F .

The line d is the *director line* and the point F is the *focus*. The distance between the focus and the director line is denoted by p and represents the *parameter* of the parabola.

Consider a Cartesian system of coordinates xOy , in which $F\left(\frac{p}{2}, 0\right)$ and $d : x = -\frac{p}{2}$. If $M(x, y)$ is an arbitrary point of the parabola, then it verifies

$$|MN| = |MF|,$$

where N is the orthogonal projection of M on Oy .

Conics

The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets

Thus, the coordinates of a point of the parabola verify

$$\sqrt{\left(x + \frac{p}{2}\right)^2 + 0} = \sqrt{\left(x - \frac{p}{2}\right)^2 + y^2} \Leftrightarrow$$

$$\Leftrightarrow \left(x + \frac{p}{2}\right)^2 = \left(x - \frac{p}{2}\right)^2 + y^2 \Leftrightarrow$$

$$\Leftrightarrow x^2 + px + \frac{p^2}{4} = x^2 - px + \frac{p^2}{4} + y^2,$$

and the equation of the parabola is

$$y^2 = 2px. \quad (1.1)$$

Remark 1.2

The equation (1.1) is equivalent to $y = \pm\sqrt{2px}$, so that the parabola is symmetric with respect to the x -axis.

Representing the graph of the function $f : [0, \infty) \rightarrow [0, \infty)$ and using the symmetry of the curve with respect to the x -axis, one obtains the graph of the parabola. One has

$$f'(x) = \frac{p}{\sqrt{2px_0}}; \quad f''(x) = -\frac{p}{2x\sqrt{2x}}.$$

x	0	∞			
$f'(x)$		+	+	+	+
$f(x)$	0	\nearrow			
$f''(x)$	-	-	-	-	-

Finally, one can easily show that the equation of tangent to the parabola $\mathcal{P} : y^2 = 2px$ at one of its point $M_0(x_0, y_0)$ is $T_{M_0}(\mathcal{P}) : yy_0 = p(x + x_0)$.

Conics

The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets

The *ellipsoid* is the quadric surface given by the equation

$$\mathcal{E} : \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} - 1 = 0, \quad a, b, c \in \mathbb{R}_+^*. \quad (2.1)$$

- ▶ The coordinate planes are all planes of symmetry of \mathcal{E} since, for an arbitrary point $M(x, y, z) \in \mathcal{E}$, its symmetric points with respect to these planes, $M_1(-x, y, z)$, $M_2(x, -y, z)$ and $M_3(x, y, -z)$ belong to \mathcal{E} ; therefore, the coordinate axes are axes of symmetry for \mathcal{E} and the origin O is the center of symmetry of the ellipsoid (2.1);
- ▶ The traces in the coordinates planes are ellipses of equations

$$\left\{ \begin{array}{l} \frac{y^2}{b^2} + \frac{z^2}{c^2} - 1 = 0 \\ x = 0 \end{array} \right., \quad \left\{ \begin{array}{l} \frac{x^2}{a^2} + \frac{z^2}{c^2} - 1 = 0 \\ y = 0 \end{array} \right., \quad \left\{ \begin{array}{l} \frac{x^2}{a^2} + \frac{y^2}{b^2} - 1 = 0 \\ z = 0. \end{array} \right.$$

Conics

The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets

Conics

The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets

- ▶ The sections with planes parallel to xOy are given by setting $z = \lambda$ in (2.1). Then, a section is of equations

$$\begin{cases} \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 - \frac{\lambda^2}{c^2} \\ z = \lambda \end{cases}.$$

- ▶ If $|\lambda| < c$, the section is an ellipse

$$\begin{cases} \frac{x^2}{\left(a\sqrt{1 - \frac{\lambda^2}{c^2}}\right)^2} + \frac{y^2}{\left(b\sqrt{1 - \frac{\lambda^2}{c^2}}\right)^2} = 1 \\ z = \lambda \end{cases};$$

- ▶ If $|\lambda| = c$, the intersection is reduced to one (tangency) point $(0, 0, \lambda)$;
- ▶ If $|\lambda| > c$, the plane $z = \lambda$ does not intersect the ellipsoid \mathcal{E} .

The sections with planes parallel to xOz or yOz are obtained in a similar way.

Conics


The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets



Hyperboloids of One Sheet

Lecture 9

Assoc. Prof.
Cornel-Sebastian
PINTEA

The surface of equation

$$\mathcal{H}_1 : \frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} - 1 = 0, \quad a, b, c \in \mathbb{R}_+^*, \quad (2.2)$$

is called *hyperboloid of one sheet*.

- ▶ The coordinate planes are planes of symmetry for \mathcal{H}_1 ; hence, the coordinate axes are axes of symmetry and the origin O is the center of symmetry of \mathcal{H}_1 ;
- ▶ The intersections with the coordinates planes are, respectively, of equations

$$\left\{ \begin{array}{l} \frac{y^2}{b^2} - \frac{z^2}{c^2} - 1 = 0 \\ x = 0 \end{array} \right. ; \left\{ \begin{array}{l} \frac{x^2}{a^2} - \frac{z^2}{c^2} - 1 = 0 \\ y = 0 \end{array} \right. ; \left\{ \begin{array}{l} \frac{x^2}{a^2} + \frac{y^2}{b^2} - 1 = 0 \\ z = 0 \end{array} \right. ;$$

a hyperbola a hyperbola an ellipse

Conics

The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets

- The intersections with planes parallel to the coordinate planes are

$$\left\{ \begin{array}{l} \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1 - \frac{\lambda^2}{a^2} \\ x = \lambda \end{array} \right. ; \left\{ \begin{array}{l} \frac{x^2}{a^2} - \frac{z^2}{c^2} = 1 - \frac{\lambda^2}{b^2} \\ y = \lambda \end{array} \right. ;$$

hyperbolas hyperbolas

$$\left\{ \begin{array}{l} \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 + \frac{\lambda^2}{c^2} \\ z = \lambda \end{array} \right. ;$$

ellipses

Remark: The surface \mathcal{H}_1 contains two families of lines, as

$$\frac{x^2}{a^2} - \frac{z^2}{c^2} = 1 - \frac{y^2}{b^2} \Leftrightarrow \left(\frac{x}{a} + \frac{z}{c} \right) \left(\frac{x}{a} - \frac{z}{c} \right) = \left(1 + \frac{y}{b} \right) \left(1 - \frac{y}{b} \right).$$

The equations of the two families of lines are:

Conics

The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets

Conics

The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets

$$d_\lambda : \begin{cases} \lambda \left(\frac{x}{a} + \frac{z}{c} \right) = 1 + \frac{y}{b} \\ \frac{x}{a} - \frac{z}{c} = \lambda \left(1 - \frac{y}{b} \right) \end{cases}, \lambda \in \mathbb{R},$$
$$d'_\mu : \begin{cases} \mu \left(\frac{x}{a} + \frac{z}{c} \right) = 1 - \frac{y}{b} \\ \frac{x}{a} - \frac{z}{c} = \mu \left(1 + \frac{y}{b} \right) \end{cases}, \mu \in \mathbb{R}.$$

Through any point on \mathcal{H}_1 pass two lines, one line from each family.

Conics


The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets



Hiperboloidul cu o pânză H_1

Hyperboloids of Two Sheets

Lecture 9

Assoc. Prof.
Cornel-Sebastian
PINTEA

The *hyperboloid of two sheets* is the surface of equation

$$\mathcal{H}_2 : \frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} + 1 = 0, \quad a, b, c \in \mathbb{R}_+^*. \quad (2.3)$$

- ▶ The coordinate planes are planes of symmetry for \mathcal{H}_1 , the coordinate axes are axes of symmetry and the origin O is the center of symmetry of \mathcal{H}_1 ;
- ▶ The intersections with the coordinates planes are, respectively,

$$\left\{ \begin{array}{l} \frac{y^2}{b^2} - \frac{z^2}{c^2} + 1 = 0 \\ x = 0 \end{array} \right., \quad \left\{ \begin{array}{l} \frac{x^2}{a^2} - \frac{z^2}{c^2} + 1 = 0 \\ y = 0 \end{array} \right., \quad \left\{ \begin{array}{l} \frac{x^2}{a^2} + \frac{y^2}{b^2} + 1 = 0 \\ z = 0 \end{array} \right. ;$$

a hyperbola; a hyperbola the empty set

Conics

The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets

- The intersections with planes parallel to the coordinate planes are

$$\left\{ \begin{array}{l} \frac{y^2}{b^2} - \frac{z^2}{c^2} = -1 - \frac{\lambda^2}{a^2} \\ x = \lambda \\ \text{hyperbolas} \end{array} \right., \quad \left\{ \begin{array}{l} \frac{x^2}{a^2} - \frac{z^2}{c^2} = -1 - \frac{\lambda^2}{b^2} \\ y = \lambda \\ \text{hyperbolas} \end{array} \right.$$

$$\text{and } \left\{ \begin{array}{l} \frac{x^2}{a^2} + \frac{y^2}{b^2} = -1 + \frac{\lambda^2}{c^2} \\ z = \lambda \end{array} \right. .$$

- If $|\lambda| > c$, the section is an ellipse;
- If $|\lambda| = c$, the intersection reduces to a point $(0, 0, \lambda)$;
- If $|\lambda| < c$, one obtains the empty set.

Conics

The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets

Conics


The Parabola

Quadrics


The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets



hiperbolidul
cu două pânze H_2



hiperboloizii H_1 , H_2
și conul asimptot comun

Conics





The Parabola

Quadrics

The ellipsoid

Hyperboloids of One Sheet

Hyperboloids of Two Sheets

-  [AnTo] Andrica, D., Țopan, L., Analytic geometry, Cluj University Press, 2004.
-  [GaRa] Galbură Gh., Radó, F., Geometrie, Editura didactică și pedagogică-București, 1979.
-  [Pi] Pinte, C. Geometrie. Elemente de geometrie analitică. Elemente de geometrie diferențială a curbelor și suprafețelor, Presa Universitară Clujeană, 2001.
-  [ROGV] Radó, F., Orban, B., Groze, V., Vasiu, A., Culegere de Probleme de Geometrie, Lit. Univ. "Babeș-Bolyai", Cluj-Napoca, 1979.