

Geometry¹

First Year, Computer science

Assoc. Prof. Cornel-Sebastian PINTEA

“Babeş-Bolyai” University
Faculty of Mathematics and Computer Sciences
Cluj-Napoca, Romania

Lecture 2, 05.03.2014

¹These notes are not in a final form. They are continuously being improved

Straight lines and planes

Dependence and independence of vectors.
The directions of the straight lines and planes
The vector equation of the straight lines and planes

Straight lines and planes

Dependence and independence of vectors. The directions of the straight lines and planes

The vector equation of the straight lines and planes

Dependence and independence of vectors

Lecture 2

Assoc. Prof.
Cornel-Sebastian
PINTEA

Definition 1.1

1. The vectors \overrightarrow{OA} , \overrightarrow{OB} are said to be *collinear* if the points O, A, B are collinear. Otherwise the vectors \overrightarrow{OA} , \overrightarrow{OB} are said to be *noncollinear*.
2. The vectors \overrightarrow{OA} , \overrightarrow{OB} , \overrightarrow{OC} are said to be *coplanar* if the points O, A, B, C are coplanar. Otherwise the vectors \overrightarrow{OA} , \overrightarrow{OB} , \overrightarrow{OC} are *noncoplanar*.

Straight lines and
planes

Dependence and
independence of vectors.
The directions of the
straight lines and planes

The vector equation of the
straight lines and planes

Remark 1.2

1. The vectors \overrightarrow{OA} , \overrightarrow{OB} are linearly (in)dependent if and only if they are (non)collinear.
2. The vectors \overrightarrow{OA} , \overrightarrow{OB} , \overrightarrow{OC} are linearly (in)dependent if and only if they are (non)coplanar.



Straight lines and
planes

Dependence and
independence of vectors.
The directions of the
straight lines and planes

The vector equation of the
straight lines and planes

Proposition 1.3

The vectors \overrightarrow{OA} , \overrightarrow{OB} , \overrightarrow{OC} form a basis of \mathcal{V} if and only if they are noncoplanar.

Corolar 1.4

The dimension of the vector space of free vectors \mathcal{V} is three.

The directions of the straight lines and planes

Lecture 2

Assoc. Prof.
Cornel-Sebastian
PINTEA

Proposition 1.5

Let Δ be a straight line and let $A \in \Delta$ be a given point.

The set $\vec{\Delta} = \{\vec{AM} \mid M \in \Delta\}$ is an one dimensional subspace of \mathcal{V} . It is independent on the choice of A on the line Δ and is called the director subspace of Δ or the direction of Δ .

Remark 1.6

The straight lines Δ , Δ' are parallel if and only if $\vec{\Delta} = \vec{\Delta}'$

Definition 1.7

We call *director vector* of the straight line Δ every nonzero vector $\{\vec{d}\} \in \vec{\Delta}$.

If $\vec{d} \in \mathcal{V}$ is a nonzero vector and $A \in \mathcal{P}$ is a given point, then there exists a unique straight line which passes

Straight lines and
planes

Dependence on
independence of vectors.
The directions of the
straight lines and planes

The vector equation of the
straight lines and planes

through A and has the direction $\langle \vec{d} \rangle$. This straight line is $\Delta = \{M \in \mathcal{P} \mid \overrightarrow{AM} \in \langle \vec{d} \rangle\}$. Δ is called the straight line which passes through A and is parallel to the vector \vec{d} .

Proposition 1.8

Let π be a plane and let $A \in \pi$ be a given point. The set $\vec{\pi} = \{\overrightarrow{AM} \in \mathcal{V} \mid M \in \pi\}$ is a two dimensional subspace of \mathcal{V} . It is independent on the position of A inside π and is called the director subspace, the director plane or the direction of the plane π .

Remark 1.9

- The planes π, π' are parallel if and only if $\vec{\pi} = \vec{\pi}'$.
- If \vec{d}_1, \vec{d}_2 are two linearly independent vectors and $A \in \mathcal{P}$ is a fixed point, then there exists a unique plane $\pi = \{M \in \mathcal{P} \mid \overrightarrow{AM} \in \langle \vec{d}_1, \vec{d}_2 \rangle\}$ passing through A and having the direction $\langle \vec{d}_1, \vec{d}_2 \rangle$. It is equally said to be parallel to the vectors \vec{d}_1 and \vec{d}_2 .

Straight lines and planes

Dependence an independence of vectors.
The directions of the straight lines and planes

The vector equation of the straight lines and planes

The vector equation of the straight lines and planes

Let Δ be a straight line and let $A \in \Delta$ be a given point.

$$\vec{r}_M = \overrightarrow{OM} = \overrightarrow{OA} + \overrightarrow{AM} = \vec{r}_A + \overrightarrow{AM}.$$

Thus

$$\begin{aligned}\{\vec{r}_M \mid M \in \Delta\} &= \{\vec{r}_A + \overrightarrow{AM} \mid M \in \Delta\} \\ &= \vec{r}_A + \{\overrightarrow{AM} \mid M \in \Delta\} \\ &= \vec{r}_A + \vec{\Delta}.\end{aligned}$$

Similarly, for a plane π and $B \in \pi$ a given point, then

$$\{\vec{r}_M \mid M \in \pi\} = \vec{r}_B + \vec{\pi}.$$

Generally speaking, a subset X of a vector space is called *affine variety* if either $X = \emptyset$ or there exists $a \in V$ and a vector subspace U of V , such that $X = a + U$.

$$\dim(X) = \begin{cases} -1 & \text{dacă } X = \emptyset \\ \dim(U) & \text{dacă } X = a + U, \end{cases}$$

Straight lines and planes

Dependence and independence of vectors.
The directions of the straight lines and planes

The vector equation of the straight lines and planes

Proposition 1.10

The bijection φ_o transforms the straight lines and the planes of the space \mathcal{P} into the one and two dimensional affine varieties of the vector space \mathcal{V} .

Let Δ be a straight line, let π be a plane, $\{\vec{d}\}$ be a basis of $\vec{\Delta}$ and let $[\vec{d}_1, \vec{d}_2]$ be a basis of $\vec{\pi}$. Then for $A \in \Delta$, we obtain the equivalence $M \in \Delta$ if and only if there exists $\lambda \in \mathbb{R}$ such that

$$\vec{r}_M = \vec{r}_A + \lambda \vec{d}. \quad (1.1)$$

The relation (1.1) is called *the vector equation* of the straight line Δ . Similarly, for $B \in \pi$, we obtain the equivalence $M \in \pi$ if and only if there exists $\lambda_1, \lambda_2 \in \mathbb{R}$ such that

$$\vec{r}_M = \vec{r}_B + \lambda_1 \vec{d}_1 + \lambda_2 \vec{d}_2. \quad (1.2)$$

The relation (1.2) is called the *vector equation* of the plane π .

Straight lines and
planes

Dependence and
independence of vectors.
The directions of the
straight lines and planes

The vector equation of the
straight lines and planes

Straight lines and
planes

Dependence on
independence of vectors.
The directions of the
straight lines and planes

The vector equation of the
straight lines and planes

Proposition 1.11

If A, B are different points of a straight line Δ , then its vector equation can be put in the form

$$\vec{r}_M = (1 - \lambda)\vec{r}_A + \lambda\vec{r}_B, \lambda \in \mathbb{R}. \quad (1.3)$$

Proposition 1.12

If A, B, C are three noncolinear points within the plane π , then the vector equation of the plane π can be put in the form

$$\vec{r}_M = (1 - \lambda_1 - \lambda_2)\vec{r}_A + \lambda_1\vec{r}_B + \lambda_2\vec{r}_C, \lambda_1, \lambda_2 \in \mathbb{R}. \quad (1.4)$$

Example 1.13

Consider the angle BOB' and the points $A \in [OB]$, $A' \in [OB']$. We shall show that

$$\vec{r}_M = m \frac{1-n}{1-mn} \vec{u} + n \frac{1-m}{1-mn} \vec{v},$$

where $\{M\} = AB' \cap A'B$, $\vec{u} = \overrightarrow{OA}$, $\vec{v} = \overrightarrow{OA'}$, $\overrightarrow{OB} = m \overrightarrow{OA}$ and $\overrightarrow{OB'} = n \overrightarrow{OA'}$, i.e.

$$\overrightarrow{OM} = m \frac{1-n}{1-mn} \overrightarrow{OA} + n \frac{1-m}{1-mn} \overrightarrow{OA'}.$$

The vector equations of the lines AB' and $A'B$ are:

$$AB' : \vec{r}_x = (1-\lambda)\vec{r}_A + \lambda\vec{r}_{B'}, \quad A'B : \vec{r}_y = (1-\mu)\vec{r}_{A'} + \mu\vec{r}_B,$$

or, equivalently

$$AB' : \vec{r}_x = (1-\lambda)\vec{u} + \lambda n \vec{v}, \quad A'B : \vec{r}_y = (1-\mu)\vec{v} + \mu m \vec{u}.$$

Straight lines and
planes

Dependence and
independence of vectors.
The directions of the
straight lines and planes

The vector equation of the
straight lines and planes




Figure: 1

Since $\{M\} = AB' \cap A'B$, it follows that \vec{r}_M both a representation in the form $(1 - \lambda)\vec{u} + \lambda n\vec{v}$ and a representation in the form $(1 - \mu)\vec{v} + \mu m\vec{u}$, i.e.

$$\vec{r}_M = (1 - \lambda)\vec{u} + \lambda n\vec{v} = (1 - \mu)\vec{v} + \mu m\vec{u}, \quad \lambda, \mu \in \mathbb{R}.$$

The linear independence of the vectors $\vec{u} = \overrightarrow{OA}$, $\vec{v} = \overrightarrow{OA'}$ leads us to the compatible linear system

$$\begin{aligned} 1 - \lambda &= \mu m \\ \lambda n &= 1 - \mu, \end{aligned}$$

whose solution is

$$\lambda = \frac{1 - m}{1 - mn}, \mu = \frac{1 - n}{1 - mn},$$

i.e.

$$1 - \mu = n \frac{1 - m}{1 - mn}.$$

Thus,

$$\vec{r}_M = m \frac{1 - n}{1 - mn} \vec{u} + n \frac{1 - m}{1 - mn} \vec{v}.$$

Straight lines and planes

Dependence an
independence of vectors.
The directions of the
straight lines and planes

The vector ecuation of the
straight lines and planes

Exemple 1.14

(Newton's theorem) *The midpoints of a the diagonals of a complet quadrilater are collinear.*

Consider the convex quadrilater $OABC$ with pairwise unparallel opposite sides. Let us also consider $\{D\} = OC \cap AB$ and $\{E\} = OA \cap BC$. The figure $OABCDE$ is called *complete quadrilateral*, and its diagonals are OB , AC and DE .

