

Geometry¹

First Year, Computer science

Assoc. Prof. Cornel-Sebastian PINTEA

“Babeş-Bolyai” University
Faculty of Mathematics and Computer Sciences
Cluj-Napoca, Romania

Lecture 10, 14.05.2014

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

¹These notes are not in a final form. They are continuously being improved

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder, Hyperbolic Cylinder, Parabolic Cylinder

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

The equation of the tangent plane and the normal line

Lecture 10

Assoc. Prof.
Cornel-Sebastian
PINTEA

Quadrics

The equation of the tangent
plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

Proposition 1.1 ([Pi, p. 256])

If $D \subseteq \mathbb{R}^3$ is an open set and $F : D \longrightarrow \mathbb{R}$ is a C^1 -smooth function, then the gradient vector field $\text{grad}F := (F_x, F_y, F_z)$ is normal to the level sets of F at every point

$M_0(x_0, y_0, z_0) \in D$ where $(\text{grad}F)(x_0, y_0, z_0)$ is nonzero. More precisely, $(\text{grad}F)(x_0, y_0, z_0)$ is a normal vector of the plane $T_{M_0}(F^{-1}(F(x_0, y_0, z_0)))$ tangent to the level set $F^{-1}(F(x_0, y_0, z_0))$ at its point M_0 .

Proposition 1.2

Let $D \subseteq \mathbb{R}^3$ be an open set and $F : D \rightarrow \mathbb{R}$ be a C^1 -smooth function. If $(\text{grad} F)(x_0, y_0, z_0) \neq 0$ at some point $M_0(x_0, y_0, z_0) \in D$, then the equation of the plane $T_{M_0}(S)$ -tangent to the implicit surface

$$S : F(x, y, z) = F(x_0, y_0, z_0)$$

at its point $M_0(x_0, y_0, z_0) \in S$ is

$$F_x(x_0, y_0, z_0)(x - x_0) + F_y(x_0, y_0, z_0)(y - y_0) + F_z(x_0, y_0, z_0)(z - z_0) = 0.$$

The equations of the normal line to S at M_0 are

$$\frac{x - x_0}{F_x(x_0, y_0, z_0)} = \frac{y - y_0}{F_y(x_0, y_0, z_0)} = \frac{z - z_0}{F_z(x_0, y_0, z_0)}.$$

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

Exemple 1.3

The equations of the tangent plane and the normal line to the ellipsoid

$$\mathcal{E} : \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1,$$

at its point $M_0(x_0, y_0, z_0)$ are

$$T_{M_0}(\mathcal{E}) : \frac{x_0 x}{a^2} + \frac{y_0 y}{b^2} + \frac{z_0 z}{c^2} = 1$$

$$N_{M_0}(\mathcal{E}) : \frac{a^2}{x_0}(x - x_0) = \frac{b^2}{y_0}(y - y_0) = \frac{c^2}{z_0}(z - z_0).$$

Indeed, the gradient of the function $F : \mathbb{R}^3 \rightarrow \mathbb{R}$,

$$F(x, y, z) = \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} - 1 \text{ is } (\text{grad} F) = \left(\frac{2x}{a^2}, \frac{2y}{b^2}, \frac{2z}{c^2} \right)$$

and

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

$\frac{1}{2}(\text{grad} F)(x_0, y_0, z_0) = \left(\frac{x_0}{a^2}, \frac{y_0}{b^2}, \frac{z_0}{c^2} \right)$ is a normal vector to the tangent plane $T_{M_0}(\mathcal{E})$ to the ellipsoid $\mathcal{E} = F^{-1}(0)$ at $M_0(x_0, y_0, z_0)$.

Exemple 1.4

The equations of the tangent plane and the normal line to the hyperboloid of one sheet

$$\mathcal{H}_1 : \frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1,$$

at its point $M_0(x_0, y_0, z_0)$ are

$$T_{M_0}(\mathcal{H}_1) : \frac{x_0 x}{a^2} + \frac{y_0 y}{b^2} - \frac{z_0 z}{c^2} = 1$$

$$N_{M_0}(\mathcal{H}_1) : \frac{a^2}{x_0}(x - x_0) = \frac{b^2}{y_0}(y - y_0) = \frac{c^2}{z_0}(z_0 - z).$$

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

Exemple 1.5

The equations of the tangent plane and the normal line to the hyperboloid of two sheets

$$\mathcal{H}_2 : \frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} + 1 = 0,$$

at its point $M_0(x_0, y_0, z_0)$ are

$$T_{M_0}(\mathcal{H}_2) : \frac{x_0 x}{a^2} + \frac{y_0 y}{b^2} - \frac{z_0 z}{c^2} = -1$$

$$N_{M_0}(\mathcal{H}_2) : \frac{a^2}{x_0}(x - x_0) = \frac{b^2}{y_0}(y - y_0) = \frac{c^2}{z_0}(z_0 - z).$$

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

The surface of equation

$$\mathcal{C} : \frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0, \quad a, b, c \in \mathbb{R}_+^*, \quad (1.1)$$

is called *elliptic cone*.

- ▶ The coordinate planes are planes of symmetry for \mathcal{C} , the coordinate axes are axes of symmetry and the origin O is the center of symmetry of \mathcal{C} ;
- ▶ The intersections with the coordinates planes are

$$\left\{ \begin{array}{l} \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0 \\ x = 0 \\ \text{two lines} \end{array} \right., \quad \left\{ \begin{array}{l} \frac{x^2}{a^2} - \frac{z^2}{c^2} - 1 = 0 \\ y = 0 \\ \text{two lines} \end{array} \right.,$$

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

$$\left\{ \begin{array}{l} \frac{x^2}{a^2} + \frac{y^2}{b^2} = 0 \\ z = 0 \\ \text{the origin } O(0, 0, 0). \end{array} \right.$$

- The intersections with planes parallel to the coordinate planes are

$$\left\{ \begin{array}{l} \frac{y^2}{b^2} - \frac{z^2}{c^2} = -\frac{\lambda^2}{a^2} \\ x = \lambda \\ \text{hyperbolas} \end{array} \right. ; \quad \left\{ \begin{array}{l} \frac{x^2}{a^2} - \frac{z^2}{c^2} = -\frac{\lambda^2}{b^2} \\ y = \lambda \\ \text{hyperbolas.} \end{array} \right.$$

$$\left\{ \begin{array}{l} \frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{\lambda^2}{c^2} \\ z = \lambda \\ \text{ellipses} \end{array} \right.$$

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

The surface of equation

$$\mathcal{P}_e : \frac{x^2}{p} + \frac{y^2}{q} = 2z, \quad p, q \in \mathbb{R}_+^*, \quad (1.2)$$

is called *elliptic paraboloid*.

- ▶ The planes xOz and yOz are planes of symmetry;
- ▶ The traces in the coordinate planes are

$$\left\{ \begin{array}{l} \frac{y^2}{q} = 2z \\ x = 0 \\ \text{a parabola} \end{array} \right., \left\{ \begin{array}{l} \frac{x^2}{p} = 2z \\ y = 0 \\ \text{a parabola} \end{array} \right., \left\{ \begin{array}{l} \frac{x^2}{p} + \frac{y^2}{q} = 0 \\ z = 0 \\ \text{the origin } O(0, 0, 0). \end{array} \right.$$

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

- The intersection with the planes parallel to the

coordinate planes are
$$\begin{cases} \frac{x^2}{p} + \frac{y^2}{q} = 2\lambda \\ z = \lambda \end{cases},$$

- If $\lambda > 0$, the section is an ellipse;
- If $\lambda = 0$, the intersection reduces to the origin;
- If $\lambda < 0$, one has the empty set;

and

$$\begin{cases} \frac{y^2}{q} = 2z - \frac{\lambda^2}{p} \\ x = \lambda \\ \text{parabolas} \end{cases}; \quad \begin{cases} \frac{x^2}{p} = 2z - \frac{\lambda^2}{q} \\ y = \lambda \\ \text{parabolas} \end{cases};$$

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

Quadrics

The equation of the tangent plane and the normal line


The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder



The *hyperbolic paraboloid* is the surface given by the equation

$$\mathcal{P}_h : \frac{x^2}{p} - \frac{y^2}{q} = 2z, \quad p, q \in \mathbb{R}_+^*. \quad (1.3)$$

- ▶ The planes xOz and yOz are planes of symmetry;
- ▶ The traces in the coordinate planes are, respectively,

$$- \begin{cases} \frac{y^2}{q} = 2z \\ x = 0 \\ \text{a parabola} \end{cases} ; \quad \begin{cases} \frac{x^2}{p} = 2z \\ y = 0 \\ \text{a parabola} \end{cases} ; \quad \begin{cases} \frac{x^2}{p} - \frac{y^2}{q} = 0 \\ z = 0 \\ \text{two lines.} \end{cases} ;$$

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

- The intersection with the planes parallel to the coordinate planes are

$$\left\{ \begin{array}{l} -\frac{y^2}{q} = 2z + \frac{\lambda^2}{p} \\ x = \lambda \\ \text{parabolas} \end{array} \right. ; \quad \left\{ \begin{array}{l} \frac{x^2}{p} = 2z - \frac{\lambda^2}{q} \\ y = \lambda \\ \text{parabolas.} \end{array} \right.$$

$$\left\{ \begin{array}{l} \frac{x^2}{p} - \frac{y^2}{q} = 2\lambda \\ z = \lambda \\ \text{hyperbolas} \end{array} \right.$$

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

Remark: The hyperbolic paraboloid admits two families of rectilinear generatrices. Since

$$\left(\frac{x}{\sqrt{p}} + \frac{y}{\sqrt{q}} \right) \left(\frac{x}{\sqrt{p}} - \frac{y}{\sqrt{q}} \right) = 2z,$$

then the two families are, respectively, of equations

$$d_\lambda : \begin{cases} \frac{x}{\sqrt{p}} - \frac{y}{\sqrt{q}} = \lambda \\ \lambda \left(\frac{x}{\sqrt{p}} + \frac{y}{\sqrt{q}} \right) = 2z \end{cases}, \lambda \in \mathbb{R}^* \text{ and}$$

$$d'_\mu : \begin{cases} \frac{x}{\sqrt{p}} + \frac{y}{\sqrt{q}} = \mu \\ \mu \left(\frac{x}{\sqrt{p}} - \frac{y}{\sqrt{q}} \right) = 2z \end{cases}, \mu \in \mathbb{R}^*.$$

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

Quadrics

The equation of the tangent plane and the normal line



The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder



Singular Quadrics: Elliptic Cylinder, Hyperbolic Cylinder, Parabolic Cylinder

Lecture 10

Assoc. Prof.
Cornel-Sebastian
PINTEA

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder

- ▶ The *elliptic cylinder* is the surface of equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - 1 = 0, \quad a, b > 0. \quad (1.4)$$

or

$$\frac{x^2}{a^2} + \frac{z^2}{c^2} - 1 = 0, \quad \frac{y^2}{b^2} + \frac{z^2}{c^2} - 1 = 0$$

- ▶ The *hyperbolic cylinder* is the surface of equation

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} - 1 = 0, \quad a, b > 0, \quad (1.5)$$

or

$$\frac{x^2}{a^2} - \frac{z^2}{c^2} - 1 = 0, \quad \frac{y^2}{b^2} - \frac{z^2}{c^2} - 1 = 0.$$

- ▶ The *parabolic cylinder* is the surface of equation

$$y^2 = 2px, \quad p > 0, \quad (\text{or an alternative equation}). \quad (1.6)$$

Quadrics

The equation of the tangent plane and the normal line


The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid


Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder




Cilindrul eliptic

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$




Cilindrul hiperbolic

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$







Cilindrul parabolic

$$y^2 = 2px$$



Conul

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0$$

-  [AnTo] Andrica, D., Țopan, L., Analytic geometry, Cluj University Press, 2004.
-  [GaRa] Galbură Gh., Radó, F., Geometrie, Editura didactică și pedagogică-București, 1979.
-  [Pi] Pinte, C. Geometrie. Elemente de geometrie analitică. Elemente de geometrie diferențială a curbelor și suprafețelor, Presa Universitară Clujeană, 2001.
-  [ROGV] Radó, F., Orban, B., Groze, V., Vasii, A., Culegere de Probleme de Geometrie, Lit. Univ. "Babeș-Bolyai", Cluj-Napoca, 1979.

Quadrics

The equation of the tangent plane and the normal line

The elliptic cone

The elliptic paraboloid

The hyperbolic paraboloid

Singular Quadrics

Elliptic Cylinder,
Hyperbolic Cylinder,
Parabolic Cylinder