

# Inner metric structure of complex surface singularities

Lorenzo Fantini

Université Aix-Marseille

Lille, May 16th, 2019

Joint work with André Belotto da Silva and Anne Pichon (arXiv:1905.01677)

# Local structure of a singularity

A long history: Wirtinger 1895, Milnor 1968...

$X$  complex variety,  
 $0 \in X$  isolated singularity


$$(X, 0) \hookrightarrow (\mathbb{C}^N, 0)$$

Topology: the Conical Structure Theorem

$$0 < \varepsilon \ll 1 \implies X \cap B(0, \varepsilon) \stackrel{\text{homeo}}{\sim} \text{Cone}(X \cap S(0, \varepsilon))$$

Inner metric on  $(X, 0)$

$$d_{\text{inner}}(x, y) = \inf_{\substack{\gamma: [0,1] \rightarrow X, \\ \gamma(0)=x, \gamma(1)=y}} \{\text{length}(\gamma)\}$$



I'm interested in the metric germ, not only on its bi-Lipschitz class !

(Mostovski 1985/ $\mathbb{C}$ , Parusiński 1987/ $\mathbb{R}$ , Birbrair–Neumann–Pichon 2014)

# Local structure of a singularity

A long history: Wirtinger 1895, Milnor 1968...

$X$  complex variety,  
 $0 \in X$  isolated singularity


$$(X, 0) \hookrightarrow (\mathbb{C}^N, 0)$$

Topology: the Conical Structure Theorem

$$0 < \varepsilon \ll 1 \implies X \cap B(0, \varepsilon) \stackrel{\text{homeo}}{\sim} \text{Cone}(X \cap S(0, \varepsilon))$$

Inner metric on  $(X, 0)$

$$d_{\text{inner}}(x, y) = \inf_{\substack{\gamma: [0,1] \rightarrow X, \\ \gamma(0)=x, \gamma(1)=y}} \{\text{length}(\gamma)\}$$



I'm interested in the metric germ, not only on its bi-Lipschitz class !

(Mostovski 1985/ $\mathbb{C}$ , Parusiński 1987/ $\mathbb{R}$ , Birbrair–Neumann–Pichon 2014)

# Local structure of a singularity

A long history: Wirtinger 1895, Milnor 1968...

$X$  complex variety,  
 $0 \in X$  isolated singularity


$$(X, 0) \hookrightarrow (\mathbb{C}^N, 0)$$

## Topology: the Conical Structure Theorem

$$0 < \varepsilon \ll 1 \implies X \cap B(0, \varepsilon) \stackrel{\text{homeo}}{\sim} \text{Cone}(X \cap S(0, \varepsilon))$$

## Inner metric on $(X, 0)$

$$d_{\text{inner}}(x, y) = \inf_{\substack{\gamma: [0,1] \rightarrow X, \\ \gamma(0)=x, \gamma(1)=y}} \{\text{length}(\gamma)\}$$



I'm interested in the metric germ, not only on its bi-Lipschitz class !

(Mostovski 1985/ $\mathbb{C}$ , Parusiński 1987/ $\mathbb{R}$ , Birbrair–Neumann–Pichon 2014)

# Local structure of a singularity

A long history: Wirtinger 1895, Milnor 1968...

$X$  complex variety,  
 $0 \in X$  isolated singularity


$$(X, 0) \hookrightarrow (\mathbb{C}^N, 0)$$

## Topology: the Conical Structure Theorem

$$0 < \varepsilon \ll 1 \implies X \cap B(0, \varepsilon) \stackrel{\text{homeo}}{\sim} \text{Cone}(X \cap S(0, \varepsilon))$$

## Inner metric on $(X, 0)$

$$d_{\text{inner}}(x, y) = \inf_{\substack{\gamma: [0,1] \rightarrow X, \\ \gamma(0)=x, \gamma(1)=y}} \{\text{length}(\gamma)\}$$



I'm interested in the metric germ, not only on its bi-Lipschitz class !

(Mostovski 1985/ $\mathbb{C}$ , Parusiński 1987/ $\mathbb{R}$ , Birbrair–Neumann–Pichon 2014)

# Local structure of a singularity

A long history: Wirtinger 1895, Milnor 1968...

$X$  complex variety,  
 $0 \in X$  isolated singularity


$$(X, 0) \hookrightarrow (\mathbb{C}^N, 0)$$

## Topology: the Conical Structure Theorem

$$0 < \varepsilon \ll 1 \implies X \cap B(0, \varepsilon) \stackrel{\text{homeo}}{\sim} \text{Cone}(X \cap S(0, \varepsilon))$$

## Inner metric on $(X, 0)$

$$d_{\text{inner}}(x, y) = \inf_{\substack{\gamma: [0,1] \rightarrow X, \\ \gamma(0)=x, \gamma(1)=y}} \{\text{length}(\gamma)\}$$




I'm interested in the **metric germ**, not only on its bi-Lipschitz class !

(Mostovski 1985/ $\mathbb{C}$ , Parusiński 1987/ $\mathbb{R}$ , Birbrair–Neumann–Pichon 2014)

# Inner rates

I will focus on the case of surfaces.



The inner rate  $\mathcal{I}(E)$  of  $E$  is the contact order between the two curve  $\pi_*\gamma$  and  $\pi_*\gamma'$  on  $(X, 0)$ , with respect to the inner metric:

$$d_{\text{inner}}(\pi_*\gamma \cap S_{\mathbb{C}^n}(0, \varepsilon), \pi_*\gamma' \cap S_{\mathbb{C}^n}(0, \varepsilon)) \approx \varepsilon^{\mathcal{I}(E)}$$

Interpretation:


The inner rate  $\mathcal{I}(E)$  measures the size of a small area  $\mathcal{N}(E)$  of  $(X, 0)$



Fine understanding of the inner metric structure of the germ

# Inner rates

I will focus on the case of surfaces.



The inner rate  $\mathcal{I}(E)$  of  $E$  is the contact order between the two curve  $\pi_*\gamma$  and  $\pi_*\gamma'$  on  $(X, 0)$ , with respect to the inner metric:

$$d_{\text{inner}}(\pi_*\gamma \cap S_{\mathbb{C}^n}(0, \varepsilon), \pi_*\gamma' \cap S_{\mathbb{C}^n}(0, \varepsilon)) \approx \varepsilon^{\mathcal{I}(E)}$$

Interpretation:


The inner rate  $\mathcal{I}(E)$  measures the size of a small area  $\mathcal{N}(E)$  of  $(X, 0)$

← Fine understanding of the inner metric structure of the germ



# Inner rates

I will focus on the case of surfaces.



The inner rate  $\mathcal{I}(E)$  of  $E$  is the contact order between the two curve  $\pi_*\gamma$  and  $\pi_*\gamma'$  on  $(X, 0)$ , with respect to the inner metric:

$$d_{\text{inner}}(\pi_*\gamma \cap S_{\mathbb{C}^n}(0, \varepsilon), \pi_*\gamma' \cap S_{\mathbb{C}^n}(0, \varepsilon)) \approx \varepsilon^{\mathcal{I}(E)}$$


Interpretation:

The inner rate  $\mathcal{I}(E)$  measures the size of a small area  $\mathcal{N}(E)$  of  $(X, 0)$

← Fine understanding of the inner metric structure of the germ

# Inner rates

I will focus on the case of surfaces.



The **inner rate**  $\mathcal{I}(E)$  of  $E$  is the contact order between the two curve  $\pi_*\gamma$  and  $\pi_*\gamma'$  on  $(X, 0)$ , with respect to the inner metric:

$$d_{\text{inner}}(\pi_*\gamma \cap S_{\mathbb{C}^n}(0, \varepsilon), \pi_*\gamma' \cap S_{\mathbb{C}^n}(0, \varepsilon)) \approx \varepsilon^{\mathcal{I}(E)}$$


Interpretation:

The inner rate  $\mathcal{I}(E)$  measures the size of a small area  $\mathcal{N}(E)$  of  $(X, 0)$

← Fine understanding of the inner metric structure of the germ

# Inner rates

I will focus on the case of surfaces.



The **inner rate**  $\mathcal{I}(E)$  of  $E$  is the contact order between the two curve  $\pi_*\gamma$  and  $\pi_*\gamma'$  on  $(X, 0)$ , with respect to the inner metric:

$$d_{\text{inner}}(\pi_*\gamma \cap S_{\mathbb{C}^n}(0, \varepsilon), \pi_*\gamma' \cap S_{\mathbb{C}^n}(0, \varepsilon)) \approx \varepsilon^{\mathcal{I}(E)}$$


## Interpretation:

The inner rate  $\mathcal{I}(E)$  measures the size of a small area  $\mathcal{N}(E)$  of  $(X, 0)$

← Fine understanding of the inner metric structure of the germ

# Inner rates

I will focus on the case of surfaces.



The **inner rate**  $\mathcal{I}(E)$  of  $E$  is the contact order between the two curve  $\pi_*\gamma$  and  $\pi_*\gamma'$  on  $(X, 0)$ , with respect to the inner metric:

$$d_{\text{inner}}(\pi_*\gamma \cap S_{\mathbb{C}^n}(0, \varepsilon), \pi_*\gamma' \cap S_{\mathbb{C}^n}(0, \varepsilon)) \approx \varepsilon^{\mathcal{I}(E)}$$

## Interpretation:

The inner rate  $\mathcal{I}(E)$  measures the size of a small area  $\mathcal{N}(E)$  of  $(X, 0)$

← Fine understanding of the inner metric structure of the germ

Example:

$$E_8 = \{x^2 + y^3 + z^5 = 0\} \subset \mathbb{C}^3$$

$\downarrow (y, z)$

discriminant  $\{y^3 + z^5 = 0\} \subset \mathbb{C}^2$

Example:

$$E_8 = \{x^2 + y^3 + z^5 = 0\} \subset \mathbb{C}^3$$

$$\downarrow (y, z)$$


discriminant  $\{y^3 + z^5 = 0\} \subset \mathbb{C}^2$

Example:

$$E_8 = \{x^2 + y^3 + z^5 = 0\} \subset \mathbb{C}^3$$

$\downarrow (y, z)$

discriminant  $\{y^3 + z^5 = 0\} \subset \mathbb{C}^2$




Example:

$$E_8 = \{x^2 + y^3 + z^5 = 0\} \subset \mathbb{C}^3$$

$\downarrow (y, z)$


discriminant  $\{y^3 + z^5 = 0\} \subset \mathbb{C}^2$





Example:

$$E_8 = \{x^2 + y^3 + z^5 = 0\} \subset \mathbb{C}^3$$




Classical questions:

- How does the geometry  $(X, 0)$  influence the inner rates?
- How to compute them?

Example:

$$E_8 = \{x^2 + y^3 + z^5 = 0\} \subset \mathbb{C}^3$$



Classical questions:

- How does the geometry  $(X, 0)$  influence the inner rates?
- How to compute them?

## Théorème (Belotto–F–Pichon, 2019)

Let  $\pi: X_\pi \rightarrow X$  be a good resolution of  $(X, 0)$ . Then all the inner rates of  $(X, 0)$  are completely determined by:

- the topology of  $(X, 0)$ , i.e. the weighted dual graph  $\Gamma_\pi$ ;
- the **arrows** of a **generic** hyperplane section;
- the **arrows** of the polar curves of a **generic** projection  $(X, 0) \rightarrow (\mathbb{C}^2, 0)$ .

This is a consequence of an explicit formula that we will see later.

Analogous to the study of weight functions on curves (Baker–Nicaise 2016).

# The non-archimedean link of a singularity

## Definition (Boucksom–Favre–Jonsson, F)

$$\mathrm{NL}(X, 0) = \left\{ v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{m}_{X,0}} \{v(f)\} = 1 \right\}$$

e.g. **divisorial valuation**  $\frac{\mathrm{ord}_E}{m(E)}$



It's a nice topological space, compact.

Example:  $\mathrm{NL}(\mathbb{A}_{\mathbb{C}}^2, 0) \cong$  valuative tree  
(Favre–Jonsson).

Non-archimedean avatar of the usual link:

## Theorem (F–Favre)

$L(X, 0)$  degenerates towards  $\mathrm{NL}(X, 0)$ .

Moreover, we have:

$$H_{\mathrm{sing}}^i(\mathrm{NL}(X, 0), \mathbb{Q}) \cong W^0 H_{\mathrm{sing}}^i(L(X, 0), \mathbb{Q}).$$

# The non-archimedean link of a singularity

## Definition (Boucksom–Favre–Jonsson, F)


$$\mathrm{NL}(X, 0) = \left\{ v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{m}_{X,0}} \{v(f)\} = 1 \right\}$$

e.g. divisorial valuation  $\frac{\mathrm{ord}_E}{m(E)}$



It's a nice topological space, compact.

Example:  $\mathrm{NL}(\mathbb{A}_{\mathbb{C}}^2, 0) \cong$  **valuative tree**  
(Favre–Jonsson).



Non-archimedean avatar of the usual link:

## Theorem (F–Favre)

$L(X, 0)$  degenerates towards  $\mathrm{NL}(X, 0)$ .

Moreover, we have:

$$H_{\mathrm{sing}}^i(\mathrm{NL}(X, 0), \mathbb{Q}) \cong W^0 H_{\mathrm{sing}}^i(L(X, 0), \mathbb{Q}).$$

# The non-archimedean link of a singularity

## Definition (Boucksom–Favre–Jonsson, F)

$$\mathrm{NL}(X, 0) = \left\{ v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{m}_{X,0}} \{v(f)\} = 1 \right\}$$

e.g. divisorial valuation  $\frac{\mathrm{ord}_E}{m(E)}$

It's a nice topological space, compact.

Example:  $\mathrm{NL}(\mathbb{A}_{\mathbb{C}}^2, 0) \cong$  valuative tree  
(Favre–Jonsson).



Non-archimedean avatar of the usual link:

## Theorem (F–Favre)

$L(X, 0)$  degenerates towards  $\mathrm{NL}(X, 0)$ .

Moreover, we have:

$$H_{\mathrm{sing}}^i(\mathrm{NL}(X, 0), \mathbb{Q}) \cong W^0 H_{\mathrm{sing}}^i(L(X, 0), \mathbb{Q}).$$



# The non-archimedean link of a singularity

## Definition (Boucksom–Favre–Jonsson, F)

$$\mathrm{NL}(X, 0) = \left\{ v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{m}_{X,0}} \{v(f)\} = 1 \right\}$$

e.g. divisorial valuation  $\frac{\mathrm{ord}_E}{m(E)}$

It's a nice topological space, compact.

Example:  $\mathrm{NL}(\mathbb{A}_{\mathbb{C}}^2, 0) \cong$  valuative tree  
(Favre–Jonsson).



Non-archimedean avatar of the usual link:

## Theorem (F–Favre)

$L(X, 0)$  degenerates towards  $\mathrm{NL}(X, 0)$ .

Moreover, we have:

$$H_{\mathrm{sing}}^i(\mathrm{NL}(X, 0), \mathbb{Q}) \cong W^0 H_{\mathrm{sing}}^i(L(X, 0), \mathbb{Q}).$$




# Skeletons and combinatorics of $NL(X, 0)$

$$NL(X, 0) = \{v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{M}_{X,0}} \{v(f)\} = 1\}$$

If  $\pi: X_\pi \rightarrow X$  is a good resolution of  $(X, 0)$  with dual graph  $\Gamma_\pi$ , there exists a **natural embedding**:

$$\Gamma_\pi \hookrightarrow NL(X, 0)$$

It sends a vertex  $v$  of  $\Gamma_\pi$  to the divisorial valuation associated with the exceptional component  $E_v \subset \pi^{-1}(0)$  that corresponds to  $v$ .



This induces a canonical homeomorphism:

$$\varprojlim_{\pi} \Gamma_\pi \xrightarrow{\sim} NL(X, 0)$$


# Skeletons and combinatorics of $NL(X, 0)$

$$NL(X, 0) = \{v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{M}_{X,0}} \{v(f)\} = 1\}$$

If  $\pi: X_\pi \rightarrow X$  is a good resolution of  $(X, 0)$  with dual graph  $\Gamma_\pi$ , there exists a natural embedding:

$$\Gamma_\pi \hookrightarrow NL(X, 0)$$

It sends a vertex  $v$  of  $\Gamma_\pi$  to the divisorial valuation associated with the exceptional component  $E_v \subset \pi^{-1}(0)$  that corresponds to  $v$ .



This induces a canonical homeomorphism:

$$\varprojlim_{\pi} \Gamma_\pi \xrightarrow{\sim} NL(X, 0)$$




# Skeletons and combinatorics of $NL(X, 0)$

$$NL(X, 0) = \{v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{M}_{X,0}} \{v(f)\} = 1\}$$

If  $\pi: X_\pi \rightarrow X$  is a good resolution of  $(X, 0)$  with dual graph  $\Gamma_\pi$ , there exists a natural embedding:

$$\Gamma_\pi \hookrightarrow NL(X, 0)$$

It sends a vertex  $v$  of  $\Gamma_\pi$  to the divisorial valuation associated with the exceptional component  $E_v \subset \pi^{-1}(0)$  that corresponds to  $v$ .



This induces a **canonical homeomorphism**:

$$\varprojlim_{\pi} \Gamma_\pi \xrightarrow{\sim} NL(X, 0)$$

# The Laplacian formula

Natural metric on  $\Gamma_\pi$ :

$$l([v, v']) = \frac{1}{m(v)m(v')}$$

where  $m(v)$  is the multiplicity of  $E_v$  in  $\pi^{-1}(0)$ .

→ Metric on  $\text{NL}(X, 0)$

The inner rates  $\mathcal{I}(E)$  extend to a continuous and piecewise linear map:



$$\mathcal{I}: \text{NL}(X, 0) \longrightarrow \mathbb{R}_{\geq 1}$$

Laplacian of  $\mathcal{I}$  on  $\Gamma_\pi$ :  $\Delta_{\Gamma_\pi}(\mathcal{I})(v) = \text{sum of the outgoing slopes of } \mathcal{I}|_{\Gamma_\pi} \text{ at } v$

Canonical divisor of  $\Gamma_\pi$ :  $K_{\Gamma_\pi}(v) = \text{val}_{\Gamma_\pi}(v) + 2g(v) - 2 = -\chi(\check{E}_v)$

Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperplane arrows at } v\} - \#\{\text{polar arrows at } v\})$$



# The Laplacian formula

Natural metric on  $\Gamma_\pi$ :


$$l([v, v']) = \frac{1}{m(v)m(v')}$$

where  $m(v)$  is the multiplicity of  $E_v$  in  $\pi^{-1}(0)$ .

→ Metric on  $\text{NL}(X, 0)$

The inner rates  $\mathcal{I}(E)$  extend to a **continuous and piecewise linear** map:

$$\mathcal{I}: \text{NL}(X, 0) \longrightarrow \mathbb{R}_{\geq 1}$$



Laplacian of  $\mathcal{I}$  on  $\Gamma_\pi$ :  $\Delta_{\Gamma_\pi}(\mathcal{I})(v) = \text{sum of the outgoing slopes of } \mathcal{I}|_{\Gamma_\pi} \text{ at } v$

Canonical divisor of  $\Gamma_\pi$ :  $K_{\Gamma_\pi}(v) = \text{val}_{\Gamma_\pi}(v) + 2g(v) - 2 = -\chi(\check{E}_v)$

Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperplane arrows at } v\} - \#\{\text{polar arrows at } v\})$$

# The Laplacian formula

Natural metric on  $\Gamma_\pi$ :



$$l([v, v']) = \frac{1}{m(v)m(v')}$$

where  $m(v)$  is the multiplicity of  $E_v$  in  $\pi^{-1}(0)$ .

→ Metric on  $\text{NL}(X, 0)$

The inner rates  $\mathcal{I}(E)$  extend to a continuous and piecewise linear map:

$$\mathcal{I}: \text{NL}(X, 0) \longrightarrow \mathbb{R}_{\geq 1}$$



**Laplacian** of  $\mathcal{I}$  on  $\Gamma_\pi$ :  $\Delta_{\Gamma_\pi}(\mathcal{I})(v) = \text{sum of the outgoing slopes of } \mathcal{I}|_{\Gamma_\pi} \text{ at } v$

Canonical divisor of  $\Gamma_\pi$ :  $K_{\Gamma_\pi}(v) = \text{val}_{\Gamma_\pi}(v) + 2g(v) - 2 = -\chi(\check{E}_v)$

Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperplane arrows at } v\} - \#\{\text{polar arrows at } v\})$$

# The Laplacian formula

Natural metric on  $\Gamma_\pi$ :

$$l([v, v']) = \frac{1}{m(v)m(v')}$$

where  $m(v)$  is the multiplicity of  $E_v$  in  $\pi^{-1}(0)$ .

→ Metric on  $\text{NL}(X, 0)$

The inner rates  $\mathcal{I}(E)$  extend to a continuous and piecewise linear map:



$$\mathcal{I}: \text{NL}(X, 0) \longrightarrow \mathbb{R}_{\geq 1}$$

Laplacian of  $\mathcal{I}$  on  $\Gamma_\pi$ :  $\Delta_{\Gamma_\pi}(\mathcal{I})(v) = \text{sum of the outgoing slopes of } \mathcal{I}|_{\Gamma_\pi} \text{ at } v$

**Canonical divisor** of  $\Gamma_\pi$ :  $K_{\Gamma_\pi}(v) = \text{val}_{\Gamma_\pi}(v) + 2g(v) - 2 = -\chi(\check{E}_v)$

Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperplane arrows at } v\} - \#\{\text{polar arrows at } v\})$$



# The Laplacian formula

Natural metric on  $\Gamma_\pi$ :



$$l([v, v']) = \frac{1}{m(v)m(v')}$$

where  $m(v)$  is the multiplicity of  $E_v$  in  $\pi^{-1}(0)$ .

→ Metric on  $\text{NL}(X, 0)$

The inner rates  $\mathcal{I}(E)$  extend to a continuous and piecewise linear map:

$$\mathcal{I}: \text{NL}(X, 0) \longrightarrow \mathbb{R}_{\geq 1}$$



Laplacian of  $\mathcal{I}$  on  $\Gamma_\pi$ :  $\Delta_{\Gamma_\pi}(\mathcal{I})(v) = \text{sum of the outgoing slopes of } \mathcal{I}|_{\Gamma_\pi} \text{ at } v$

Canonical divisor of  $\Gamma_\pi$ :  $K_{\Gamma_\pi}(v) = \text{val}_{\Gamma_\pi}(v) + 2g(v) - 2 = -\chi(\check{E}_v)$

Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperplane arrows at } v\} - \#\{\text{polar arrows at } v\})$$

# The Laplacian formula

## Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperpl. arrows at } v\} - \#\{\text{polar arrows at } v\})$$


Applications:

- Simple explicit computation of the inner rates
- Lê–Greuel–Teissier Formula
- We obtain strong restrictions on the relative positions of arrows.

Two possible proofs:

- Lifting the formula for  $\text{NL}(\mathbb{C}^2, 0)$  to the singular case: topology and monodromy of the Milnor fiber of a generic linear form, Dehn twists.
- Birational interpretation of the inner rates as (normalized) logarithmic Mather discrepancies: Fitting ideals, study of the zeroes and poles of some differential forms on resolutions.

$$\{(zx^2 + y^3)(x^3 + zy^2) + z^7 = 0\} \subset \mathbb{C}^3$$



# The Laplacian formula

## Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_{\pi}}(\mathcal{I})(v) = m(v)(K_{\Gamma_{\pi}}(v) + 2\#\{\text{hyperpl. arrows at } v\} - \#\{\text{polar arrows at } v\})$$


Applications:

- Simple explicit computation of the inner rates
- **Lê–Greuel–Teissier Formula**
- We obtain strong restrictions on the relative positions of arrows.

Two possible proofs:

- Lifting the formula for  $\text{NL}(\mathbb{C}^2, 0)$  to the singular case: topology and monodromy of the Milnor fiber of a generic linear form, Dehn twists.
- Birational interpretation of the inner rates as (normalized) logarithmic Mather discrepancies: Fitting ideals, study of the zeroes and poles of some differential forms on resolutions.

$$\{(zx^2 + y^3)(x^3 + zy^2) + z^7 = 0\} \subset \mathbb{C}^3$$





# The Laplacian formula

## Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperpl. arrows at } v\} - \#\{\text{polar arrows at } v\})$$


Applications:

- Simple explicit computation of the inner rates
- Lê–Greuel–Teissier Formula
- We obtain strong **restrictions on the relative positions of arrows**.

Two possible proofs:

- Lifting the formula for  $\text{NL}(\mathbb{C}^2, 0)$  to the singular case: topology and monodromy of the Milnor fiber of a generic linear form, Dehn twists.
- Birational interpretation of the inner rates as (normalized) logarithmic Mather discrepancies: Fitting ideals, study of the zeroes and poles of some differential forms on resolutions.

$$\{(zx^2 + y^3)(x^3 + zy^2) + z^7 = 0\} \subset \mathbb{C}^3$$



# The Laplacian formula

## Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperpl. arrows at } v\} - \#\{\text{polar arrows at } v\})$$


Applications:

- Simple explicit computation of the inner rates
- Lê–Greuel–Teissier Formula
- We obtain strong restrictions on the relative positions of arrows.

Two possible proofs:

- Lifting the formula for  $\text{NL}(\mathbb{C}^2, 0)$  to the singular case: **topology and monodromy of the Milnor fiber** of a generic linear form, Dehn twists.
- Birational interpretation of the inner rates as (normalized) logarithmic Mather discrepancies: Fitting ideals, study of the zeroes and poles of some differential forms on resolutions.

$$\{(zx^2 + y^3)(x^3 + zy^2) + z^7 = 0\} \subset \mathbb{C}^3$$



# The Laplacian formula

## Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperpl. arrows at } v\} - \#\{\text{polar arrows at } v\})$$

Applications:

- Simple explicit computation of the inner rates
- Lê–Greuel–Teissier Formula
- We obtain strong restrictions on the relative positions of arrows.

Two possible proofs:

- Lifting the formula for  $\text{NL}(\mathbb{C}^2, 0)$  to the singular case: topology and monodromy of the Milnor fiber of a generic linear form, Dehn twists.
- Birational interpretation of the inner rates as (normalized) logarithmic **Mather discrepancies**: Fitting ideals, study of the zeroes and poles of some differential forms on resolutions.

$$\{(zx^2 + y^3)(x^3 + zy^2) + z^7 = 0\} \subset \mathbb{C}^3$$

