

# Valuation spaces and metric properties of surface singularities

Lorenzo Fantini

Goethe-Universität Frankfurt am Main

August 25th, 2020

Seminario de Singularidades, Fortaleza

Includes joint works with C. Favre and M. Ruggiero, A. Belotto da Silva and A. Pichon.

Slides: <https://lorenzofantini.eu/fantini-fortaleza.pdf>

# Local structure of a singularity


A long history: Wirtinger 1895, Milnor 1968...

$X$  complex variety,  
 $0 \in X$  isolated singularity

$$(X, 0) \hookrightarrow (\mathbb{C}^N, 0)$$

Topology: the Conical Structure Theorem

$$0 < \varepsilon \ll 1 \implies X \cap B(0, \varepsilon) \stackrel{\text{homeo}}{\sim} \text{Cone}(X \cap S(0, \varepsilon))$$



Metrics on  $(X, 0)$

$$d_{\text{outer}}(x, y) = \|x - y\|_{\mathbb{C}^N}$$

$$d_{\text{inner}}(x, y) = \inf_{\substack{\gamma: [0,1] \rightarrow X, \\ \gamma(0)=x, \gamma(1)=y}} \{\text{length}(\gamma)\}$$


- Belotto–F–Pichon 2019: inner metric germs, not bi-Lipschitz classes!
- Belotto–F–Pichon 2020: we study LNE surfaces, i.e.  $d_{\text{outer}} \stackrel{\text{bi-Lip}}{\approx} d_{\text{inner}}$   
(Birbrair–Mostowski, 2010)

# Local structure of a singularity

A long history: Wirtinger 1895, Milnor 1968...

$X$  complex variety,  
 $0 \in X$  isolated singularity

$$(X, 0) \hookrightarrow (\mathbb{C}^N, 0)$$



Topology: the Conical Structure Theorem

$$0 < \varepsilon \ll 1 \implies X \cap B(0, \varepsilon) \stackrel{\text{homeo}}{\sim} \text{Cone}(X \cap S(0, \varepsilon))$$

Metrics on  $(X, 0)$

$$d_{\text{outer}}(x, y) = \|x - y\|_{\mathbb{C}^N}$$

$$d_{\text{inner}}(x, y) = \inf_{\substack{\gamma: [0,1] \rightarrow X, \\ \gamma(0)=x, \gamma(1)=y}} \{\text{length}(\gamma)\}$$

- Belotto–F–Pichon 2019: inner metric germs, not bi-Lipschitz classes!
- Belotto–F–Pichon 2020: we study LNE surfaces, i.e.  $d_{\text{outer}} \stackrel{\text{bi-Lip}}{\approx} d_{\text{inner}}$   
(Birbrair–Mostowski, 2010)

# Local structure of a singularity


A long history: Wirtinger 1895, Milnor 1968...

$X$  complex variety,  
 $0 \in X$  isolated singularity

$$(X, 0) \hookrightarrow (\mathbb{C}^N, 0)$$

## Topology: the Conical Structure Theorem

$$0 < \varepsilon \ll 1 \implies X \cap B(0, \varepsilon) \stackrel{\text{homeo}}{\sim} \text{Cone}(X \cap S(0, \varepsilon))$$



## Metrics on $(X, 0)$

$$d_{\text{outer}}(x, y) = \|x - y\|_{\mathbb{C}^N}$$

$$d_{\text{inner}}(x, y) = \inf_{\substack{\gamma: [0,1] \rightarrow X, \\ \gamma(0)=x, \gamma(1)=y}} \{\text{length}(\gamma)\}$$

- Belotto–F–Pichon 2019: inner metric germs, not bi-Lipschitz classes!
- Belotto–F–Pichon 2020: we study LNE surfaces, i.e.  $d_{\text{outer}} \stackrel{\text{bi-Lip}}{\approx} d_{\text{inner}}$   
(Birbrair–Mostowski, 2010)

# Local structure of a singularity


A long history: Wirtinger 1895, Milnor 1968...

$X$  complex variety,  
 $0 \in X$  isolated singularity

$$(X, 0) \hookrightarrow (\mathbb{C}^N, 0)$$

## Topology: the Conical Structure Theorem

$$0 < \varepsilon \ll 1 \implies X \cap B(0, \varepsilon) \stackrel{\text{homeo}}{\sim} \text{Cone}(X \cap S(0, \varepsilon))$$



## Metrics on $(X, 0)$

$$d_{\text{outer}}(x, y) = \|x - y\|_{\mathbb{C}^N}$$

$$d_{\text{inner}}(x, y) = \inf_{\substack{\gamma: [0,1] \rightarrow X, \\ \gamma(0)=x, \gamma(1)=y}} \{\text{length}(\gamma)\}$$

- Belotto–F–Pichon 2019: inner metric germs, not bi-Lipschitz classes!
- Belotto–F–Pichon 2020: we study LNE surfaces, i.e.  $d_{\text{outer}} \stackrel{\text{bi-Lip}}{\approx} d_{\text{inner}}$   
(Birbrair–Mostowski, 2010)

# Local structure of a singularity


A long history: Wirtinger 1895, Milnor 1968...

$X$  complex variety,  
 $0 \in X$  isolated singularity

$$(X, 0) \hookrightarrow (\mathbb{C}^N, 0)$$

## Topology: the Conical Structure Theorem

$$0 < \varepsilon \ll 1 \implies X \cap B(0, \varepsilon) \stackrel{\text{homeo}}{\sim} \text{Cone}(X \cap S(0, \varepsilon))$$



## Metrics on $(X, 0)$

$$d_{\text{outer}}(x, y) = \|x - y\|_{\mathbb{C}^N}$$

$$d_{\text{inner}}(x, y) = \inf_{\substack{\gamma: [0,1] \rightarrow X, \\ \gamma(0)=x, \gamma(1)=y}} \{\text{length}(\gamma)\}$$

- Belotto–F–Pichon 2019: inner metric germs, not bi-Lipschitz classes!
- Belotto–F–Pichon 2020: we study LNE surfaces, i.e.  $d_{\text{outer}} \stackrel{\text{bi-Lip}}{\approx} d_{\text{inner}}$   
(Birbrair–Mostowski, 2010)

# Local structure of a singularity


A long history: Wirtinger 1895, Milnor 1968...

$X$  complex variety,  
 $0 \in X$  isolated singularity

$$(X, 0) \hookrightarrow (\mathbb{C}^N, 0)$$

## Topology: the Conical Structure Theorem

$$0 < \varepsilon \ll 1 \implies X \cap B(0, \varepsilon) \stackrel{\text{homeo}}{\sim} \text{Cone}(X \cap S(0, \varepsilon))$$



## Metrics on $(X, 0)$


$$d_{\text{outer}}(x, y) = \|x - y\|_{\mathbb{C}^N}$$

$$d_{\text{inner}}(x, y) = \inf_{\substack{\gamma: [0,1] \rightarrow X, \\ \gamma(0)=x, \gamma(1)=y}} \{\text{length}(\gamma)\}$$

- Belotto–F–Pichon 2019: inner metric germs, not bi-Lipschitz classes!
- Belotto–F–Pichon 2020: we study LNE surfaces, i.e.  $d_{\text{outer}} \stackrel{\text{bi-Lip}}{\approx} d_{\text{inner}}$   
(Birbrair–Mostowski, 2010)

# Inner rates

I will focus on the case of surfaces.



The inner rate  $\mathcal{I}(E)$  of  $E$  is the contact order between the two curves  $\pi_*\gamma$  and  $\pi_*\gamma'$  on  $(X, 0)$ , with respect to the inner metric:

$$d_{\text{inner}}(\pi_*\gamma \cap S_{\mathbb{C}^n}(0, \varepsilon), \pi_*\gamma' \cap S_{\mathbb{C}^n}(0, \varepsilon)) \approx \varepsilon^{\mathcal{I}(E)}$$

Interpretation:

The inner rate  $\mathcal{I}(E)$  measures the size of a small area  $\mathcal{N}(E)$  of  $(X, 0)$




Fine understanding of the inner metric structure of the germ



# Inner rates

I will focus on the case of surfaces.



The inner rate  $\mathcal{I}(E)$  of  $E$  is the contact order between the two curves  $\pi_*\gamma$  and  $\pi_*\gamma'$  on  $(X, 0)$ , with respect to the inner metric:

$$d_{\text{inner}}(\pi_*\gamma \cap S_{\mathbb{C}^n}(0, \varepsilon), \pi_*\gamma' \cap S_{\mathbb{C}^n}(0, \varepsilon)) \approx \varepsilon^{\mathcal{I}(E)}$$

Interpretation:


The inner rate  $\mathcal{I}(E)$  measures the size of a small area  $\mathcal{N}(E)$  of  $(X, 0)$



Fine understanding of the inner metric structure of the germ

# Inner rates

I will focus on the case of surfaces.



The inner rate  $\mathcal{I}(E)$  of  $E$  is the contact order between the two curves  $\pi_*\gamma$  and  $\pi_*\gamma'$  on  $(X, 0)$ , with respect to the inner metric:

$$d_{\text{inner}}(\pi_*\gamma \cap S_{\mathbb{C}^n}(0, \varepsilon), \pi_*\gamma' \cap S_{\mathbb{C}^n}(0, \varepsilon)) \approx \varepsilon^{\mathcal{I}(E)}$$

Interpretation:


The inner rate  $\mathcal{I}(E)$  measures the size of a small area  $\mathcal{N}(E)$  of  $(X, 0)$



Fine understanding of the inner metric structure of the germ

# Inner rates

I will focus on the case of surfaces.



The **inner rate**  $\mathcal{I}(E)$  of  $E$  is the contact order between the two curves  $\pi_*\gamma$  and  $\pi_*\gamma'$  on  $(X, 0)$ , with respect to the inner metric:

$$d_{\text{inner}}(\pi_*\gamma \cap S_{\mathbb{C}^n}(0, \varepsilon), \pi_*\gamma' \cap S_{\mathbb{C}^n}(0, \varepsilon)) \approx \varepsilon^{\mathcal{I}(E)}$$

Interpretation:


The inner rate  $\mathcal{I}(E)$  measures the size of a small area  $\mathcal{N}(E)$  of  $(X, 0)$



Fine understanding of the inner metric structure of the germ

# Inner rates

I will focus on the case of surfaces.



The **inner rate**  $\mathcal{I}(E)$  of  $E$  is the contact order between the two curves  $\pi_*\gamma$  and  $\pi_*\gamma'$  on  $(X, 0)$ , with respect to the inner metric:

$$d_{\text{inner}}(\pi_*\gamma \cap S_{\mathbb{C}^n}(0, \varepsilon), \pi_*\gamma' \cap S_{\mathbb{C}^n}(0, \varepsilon)) \approx \varepsilon^{\mathcal{I}(E)}$$

## Interpretation:


The inner rate  $\mathcal{I}(E)$  measures the size of a small area  $\mathcal{N}(E)$  of  $(X, 0)$



Fine understanding of the inner metric structure of the germ

# Inner rates

I will focus on the case of surfaces.



The **inner rate**  $\mathcal{I}(E)$  of  $E$  is the contact order between the two curves  $\pi_*\gamma$  and  $\pi_*\gamma'$  on  $(X, 0)$ , with respect to the inner metric:

$$d_{\text{inner}}(\pi_*\gamma \cap S_{\mathbb{C}^n}(0, \varepsilon), \pi_*\gamma' \cap S_{\mathbb{C}^n}(0, \varepsilon)) \approx \varepsilon^{\mathcal{I}(E)}$$

## Interpretation:

The inner rate  $\mathcal{I}(E)$  measures the size of a small area  $\mathcal{N}(E)$  of  $(X, 0)$



Fine understanding of the inner metric structure of the germ

Example:

$$E_8 = \{x^2 + y^3 + z^5 = 0\} \subset \mathbb{C}^3$$

$$\downarrow (y, z)$$

discriminant  $\{y^3 + z^5 = 0\} \subset \mathbb{C}^2$

Example:

$$E_8 = \{x^2 + y^3 + z^5 = 0\} \subset \mathbb{C}^3$$

$$\downarrow (y, z)$$


discriminant  $\{y^3 + z^5 = 0\} \subset \mathbb{C}^2$

Example:

$$E_8 = \{x^2 + y^3 + z^5 = 0\} \subset \mathbb{C}^3$$

$\downarrow (y, z)$

discriminant  $\{y^3 + z^5 = 0\} \subset \mathbb{C}^2$






Example:

$$E_8 = \{x^2 + y^3 + z^5 = 0\} \subset \mathbb{C}^3$$


$\downarrow (y, z)$

discriminant  $\{y^3 + z^5 = 0\} \subset \mathbb{C}^2$



Example:

$$E_8 = \{x^2 + y^3 + z^5 = 0\} \subset \mathbb{C}^3$$




Fundamental questions (going back to Birbrair–Neumann–Pichon 2014):

- How does the geometry  $(X, 0)$  influence the inner rates?
- How to compute them?

Example:

$$E_8 = \{x^2 + y^3 + z^5 = 0\} \subset \mathbb{C}^3$$



Fundamental questions (going back to Birbrair–Neumann–Pichon 2014):

- How does the geometry  $(X, 0)$  influence the inner rates?
- How to compute them?

## Theorem (Belotto–F–Pichon, 2019)

Let  $\pi: X_\pi \rightarrow X$  be a good resolution of  $(X, 0)$ . Then all the inner rates of  $(X, 0)$  are completely determined by:

- the topology of  $(X, 0)$ , i.e. the weighted dual graph  $\Gamma_\pi$ ;
- the **arrows** of a **generic** hyperplane section;
- the **arrows** of the polar curves of a **generic** projection  $(X, 0) \rightarrow (\mathbb{C}^2, 0)$ .

This is a consequence of an explicit formula that we will see later.

To get this, we use a new tool: the **non-archimedean link** of the singularity  $(X, 0)$ .

# The non-archimedean link of a singularity

## Definition (Boucksom–Favre–Jonsson, F)

$$\mathrm{NL}(X, 0) = \{v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{m}_{X,0}} \{v(f)\} = 1\}$$

$$v(0) = +\infty, v(fg) = v(f) + v(g), v(f+g) \geq \min\{v(f), v(g)\}$$

- if  $(\gamma, 0) \subset (X, 0)$  curve germ, the order of vanishing at 0 along  $\gamma$  is a semivaluation
- if  $\pi: (Y, D) \rightarrow (X, 0)$  is a modification ( $Y$  normal,  $D$  snc divisor), then  $\mathrm{ord}_E$  is a semivaluation (“divisorial valuation”)

It’s a nice topological space, compact.

Example:  $\mathrm{NL}(\mathbb{A}_{\mathbb{C}}^2, 0) \cong$  valutive tree  
(Favre–Jonsson).

# The non-archimedean link of a singularity

## Definition (Boucksom–Favre–Jonsson, F)

$$\mathrm{NL}(X, 0) = \{v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{m}_{X,0}} \{v(f)\} = 1\}$$

$$v(0) = +\infty, v(fg) = v(f) + v(g), v(f+g) \geq \min\{v(f), v(g)\}$$

- if  $(\gamma, 0) \subset (X, 0)$  curve germ, the **order of vanishing** at 0 along  $\gamma$  is a semivaluation
- if  $\pi: (Y, D) \rightarrow (X, 0)$  is a modification ( $Y$  normal,  $D$  snc divisor), then  $\mathrm{ord}_E$  is a semivaluation (“**divisorial valuation**”)

It's a nice topological space, compact.

Example:  $\mathrm{NL}(\mathbb{A}_{\mathbb{C}}^2, 0) \cong$  valutive tree  
(Favre–Jonsson).

# The non-archimedean link of a singularity

## Definition (Bouxsom–Favre–Jonsson, F)


$$\text{NL}(X, 0) = \{v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{m}_{X,0}} \{v(f)\} = 1\}$$

$$v(0) = +\infty, v(fg) = v(f) + v(g), v(f+g) \geq \min\{v(f), v(g)\}$$

- if  $(\gamma, 0) \subset (X, 0)$  curve germ, the order of vanishing at 0 along  $\gamma$  is a semivaluation
- if  $\pi: (Y, D) \rightarrow (X, 0)$  is a modification ( $Y$  normal,  $D$  snc divisor), then  $\text{ord}_E$  is a semivaluation (“divisorial valuation”)

It’s a nice topological space, compact.

Example:  $\text{NL}(\mathbb{A}_{\mathbb{C}}^2, 0) \cong$  **valuative tree**  
(Favre–Jonsson).



# The non-archimedean link of a singularity

## Intuition

$NL(X, 0)$  is a non-archimedean version of the usual link  $L(X, 0)$

Indeed, denote by  $(z_1, \dots, z_N)$  a set of local coordinates of  $\mathbb{C}^N$  at 0. Then:

$$NL(X, 0) = \left\{ x \in X^{\text{an}} \mid \max_i |z_i(x)| = \varepsilon \right\}$$

Berkovich space of  $X$  w.r.t. the trivial absolute value

## Theorem (F–Favre)

$L(X, 0)$  degenerates towards  $NL(X, 0)$ .

Moreover, we have:

$$H_{\text{sing}}^i(NL(X, 0), \mathbb{Q}) \cong W^0 H_{\text{sing}}^i(L(X, 0), \mathbb{Q}).$$

# The non-archimedean link of a singularity

## Intuition

$NL(X, 0)$  is a non-archimedean version of the usual link  $L(X, 0)$

Indeed, denote by  $(z_1, \dots, z_N)$  a set of local coordinates of  $\mathbb{C}^N$  at 0. Then:

$$NL(X, 0) = \left\{ x \in X^{\text{an}} \mid \max_i |z_i(x)| = \varepsilon \right\}$$

Berkovich space of  $X$  w.r.t. the trivial absolute value

## Theorem (F–Favre)

$L(X, 0)$  degenerates towards  $NL(X, 0)$ .

Moreover, we have:

$$H_{\text{sing}}^i(NL(X, 0), \mathbb{Q}) \cong W^0 H_{\text{sing}}^i(L(X, 0), \mathbb{Q}).$$



# The non-archimedean link of a singularity

## Intuition

$NL(X, 0)$  is a non-archimedean version of the usual link  $L(X, 0)$

Indeed, denote by  $(z_1, \dots, z_N)$  a set of local coordinates of  $\mathbb{C}^N$  at 0. Then:

$$NL(X, 0) = \left\{ x \in X^{\text{an}} \mid \max_i |z_i(x)| = \varepsilon \right\}$$


Berkovich space of  $X$  w.r.t. the trivial absolute value

## Theorem (F–Favre)

$L(X, 0)$  degenerates towards  $NL(X, 0)$ .

Moreover, we have:

$$H_{\text{sing}}^i(NL(X, 0), \mathbb{Q}) \cong W^0 H_{\text{sing}}^i(L(X, 0), \mathbb{Q}).$$



# The non-archimedean link of a singularity

## Intuition

$NL(X, 0)$  is a non-archimedean version of the usual link  $L(X, 0)$

Indeed, denote by  $(z_1, \dots, z_N)$  a set of local coordinates of  $\mathbb{C}^N$  at 0. Then:

$$NL(X, 0) = \left\{ x \in X^{\text{an}} \mid \max_i |z_i(x)| = \varepsilon \right\}$$


Berkovich space of  $X$  w.r.t. the trivial absolute value

## Theorem (F–Favre)

$L(X, 0)$  degenerates towards  $NL(X, 0)$ .

Moreover, we have:

$$H_{\text{sing}}^i(NL(X, 0), \mathbb{Q}) \cong W^0 H_{\text{sing}}^i(L(X, 0), \mathbb{Q}).$$




# Skeletons and combinatorics of $NL(X, 0)$

$$NL(X, 0) = \{v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{M}_{X,0}} \{v(f)\} = 1\}$$

If  $\pi: X_\pi \rightarrow X$  is a good resolution of  $(X, 0)$  with dual graph  $\Gamma_\pi$ , there exists a **natural embedding**:

$$\Gamma_\pi \hookrightarrow NL(X, 0)$$

It sends a vertex  $v$  of  $\Gamma_\pi$  to the divisorial valuation associated with the exceptional component  $E_v \subset \pi^{-1}(0)$  that corresponds to  $v$ .



This induces a canonical homeomorphism:

$$\varprojlim_{\pi} \Gamma_\pi \xrightarrow{\sim} NL(X, 0)$$


# Skeletons and combinatorics of $NL(X, 0)$

$$NL(X, 0) = \{v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{M}_{X,0}} \{v(f)\} = 1\}$$

If  $\pi: X_\pi \rightarrow X$  is a good resolution of  $(X, 0)$  with dual graph  $\Gamma_\pi$ , there exists a natural embedding:

$$\Gamma_\pi \hookrightarrow NL(X, 0)$$

It sends a vertex  $v$  of  $\Gamma_\pi$  to the divisorial valuation associated with the exceptional component  $E_v \subset \pi^{-1}(0)$  that corresponds to  $v$ .



This induces a canonical homeomorphism:

$$\varprojlim_{\pi} \Gamma_\pi \xrightarrow{\sim} NL(X, 0)$$


# Skeletons and combinatorics of $NL(X, 0)$

$$NL(X, 0) = \{v: \widehat{\mathcal{O}_{X,0}} \rightarrow \mathbb{R}_+ \cup \{+\infty\} \text{ semivaluation} \mid \min_{f \in \mathfrak{M}_{X,0}} \{v(f)\} = 1\}$$

If  $\pi: X_\pi \rightarrow X$  is a good resolution of  $(X, 0)$  with dual graph  $\Gamma_\pi$ , there exists a natural embedding:


$$\Gamma_\pi \hookrightarrow NL(X, 0)$$

It sends a vertex  $v$  of  $\Gamma_\pi$  to the divisorial valuation associated with the exceptional component  $E_v \subset \pi^{-1}(0)$  that corresponds to  $v$ .



This induces a **canonical homeomorphism**:

$$\varprojlim_{\pi} \Gamma_\pi \xrightarrow{\sim} NL(X, 0)$$



Moreover, any proper birational map  $f: (Y, D) \rightarrow (X, 0)$  induces an isomorphism  $NL(Y, D) \cong NL(X, 0)$ .

### Theorem (Berkovich, Thuillier)

If  $\pi: (Y, D) \rightarrow (X, 0)$  is a good resolution of  $(X, 0)$ , then the retraction  $NL(X, 0) \rightarrow \Gamma_\pi = \text{Dual}(D)$  extends to a strong deformation retraction.


In particular,  $NL(X, 0)$  has the homotopy type of  $\Gamma_\pi$ .

NB: this works over any perfect field  $k$ , in any dimension.

### Corollary (Stepanov 2006, Thuillier 2007)

The homotopy type of the dual graph of a good resolution of  $(X, 0)$  does not depend on the choice of the resolution.

Remark:  $\text{Dual}(D)$  (and hence  $NL(X, 0)$ ) can be essentially arbitrary (Kollár), but it is contractible for isolated log terminal singularities (de Fernex–Kollár–Xu).



Moreover, any proper birational map  $f: (Y, D) \rightarrow (X, 0)$  induces an isomorphism  $NL(Y, D) \cong NL(X, 0)$ .

### Theorem (Berkovich, Thuillier)

If  $\pi: (Y, D) \rightarrow (X, 0)$  is a good resolution of  $(X, 0)$ , then the retraction  $NL(X, 0) \rightarrow \Gamma_\pi = \text{Dual}(D)$  extends to a strong deformation retraction.


In particular,  $NL(X, 0)$  has the homotopy type of  $\Gamma_\pi$ .

NB: this works over any perfect field  $k$ , in any dimension.

### Corollary (Stepanov 2006, Thuillier 2007)

The homotopy type of the dual graph of a good resolution of  $(X, 0)$  does not depend on the choice of the resolution.

Remark:  $\text{Dual}(D)$  (and hence  $NL(X, 0)$ ) can be essentially arbitrary (Kollár), but it is contractible for isolated log terminal singularities (de Fernex–Kollár–Xu).



Moreover, any proper birational map  $f: (Y, D) \rightarrow (X, 0)$  induces an isomorphism  $NL(Y, D) \cong NL(X, 0)$ .

### Theorem (Berkovich, Thuillier)

If  $\pi: (Y, D) \rightarrow (X, 0)$  is a good resolution of  $(X, 0)$ , then the retraction  $NL(X, 0) \rightarrow \Gamma_\pi = \text{Dual}(D)$  extends to a strong deformation retraction.

In particular,  $NL(X, 0)$  has the homotopy type of  $\Gamma_\pi$ .

NB: this works over any perfect field  $k$ , in any dimension.

### Corollary (Stepanov 2006, Thuillier 2007)

The homotopy type of the dual graph of a good resolution of  $(X, 0)$  does not depend on the choice of the resolution.

Remark:  $\text{Dual}(D)$  (and hence  $NL(X, 0)$ ) can be essentially arbitrary (Kollár), but it is contractible for isolated log terminal singularities (de Fernex–Kollár–Xu).



# Berkovich space structure

$NL(X, 0)$  inherits from the Berkovich spaces  $X^{\text{an}}$  a **non-archimedean analytic structure**.

Moreover, any good resolution  $\pi$  induces a decomposition of  $NL(X, 0)$  into (non-archimedean) discs and annuli. (Similar to the topological link!)

## Application 1 (F 2014)

Non-archimedean characterization of Nash's essential valuations of a  $k$ -surface.

$NL(X, 0)$  looks like a fractal :


## Application 2 (F–Favre–Ruggiero 2018)

Let  $(X, 0)$  normal  $k$ -surface singularity.

$NL(X, 0)$  is  
self-similar



$(X, 0)$  est is  
sandwich



# Berkovich space structure

$NL(X, 0)$  inherits from the Berkovich spaces  $X^{\text{an}}$  a non-archimedean analytic structure.

Moreover, any good resolution  $\pi$  induces a decomposition of  $NL(X, 0)$  into (non-archimedean) discs and annuli. (Similar to the topological link!)

## Application 1 (F 2014)

Non-archimedean characterization of Nash's **essential valuations** of a  $k$ -surface.

divisorial valuations of the minimal resolution

$NL(X, 0)$  looks like a fractal :


## Application 2 (F–Favre–Ruggiero 2018)

Let  $(X, 0)$  normal  $k$ -surface singularity.

$NL(X, 0)$  is  
self-similar



$(X, 0)$  est is  
sandwich



# Berkovich space structure

$NL(X, 0)$  inherits from the Berkovich spaces  $X^{\text{an}}$  a non-archimedean analytic structure.

Moreover, any good resolution  $\pi$  induces a decomposition of  $NL(X, 0)$  into (non-archimedean) discs and annuli. (Similar to the topological link!)

## Application 1 (F 2014)

Non-archimedean characterization of Nash's essential valuations of a  $k$ -surface.

divisorial valuations of the minimal resolution

$NL(X, 0)$  looks like a **fractal** :


## Application 2 (F–Favre–Ruggiero 2018)

Let  $(X, 0)$  normal  $k$ -surface singularity.

$NL(X, 0)$  is  
**self-similar**



$(X, 0)$  est is  
**sandwich**



# Back to the inner rates: the Laplacian formula

Natural metric on  $\Gamma_\pi$ :

$$l([v, v']) = \frac{1}{m(v)m(v')}$$

where  $m(v)$  is the multiplicity of  $E_v$  in  $\pi^{-1}(0)$ .

→ Metric on  $\text{NL}(X, 0)$

The inner rates  $\mathcal{I}(E)$  extend to a continuous and piecewise linear map:



$$\mathcal{I}: \text{NL}(X, 0) \longrightarrow \mathbb{R}_{\geq 1}$$

Laplacian of  $\mathcal{I}$  on  $\Gamma_\pi$ :  $\Delta_{\Gamma_\pi}(\mathcal{I})(v) = \text{sum of the outgoing slopes of } \mathcal{I}|_{\Gamma_\pi} \text{ at } v$

Canonical divisor of  $\Gamma_\pi$ :  $K_{\Gamma_\pi}(v) = \text{val}_{\Gamma_\pi}(v) + 2g(v) - 2 = -\chi(\check{E}_v)$

Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperplane arrows at } v\} - \#\{\text{polar arrows at } v\})$$



# Back to the inner rates: the Laplacian formula

Natural metric on  $\Gamma_\pi$ :


$$l([v, v']) = \frac{1}{m(v)m(v')}$$

where  $m(v)$  is the multiplicity of  $E_v$  in  $\pi^{-1}(0)$ .

→ Metric on  $\text{NL}(X, 0)$

The inner rates  $\mathcal{I}(E)$  extend to a **continuous and piecewise linear** map:

$$\mathcal{I}: \text{NL}(X, 0) \longrightarrow \mathbb{R}_{\geq 1}$$



Laplacian of  $\mathcal{I}$  on  $\Gamma_\pi$ :  $\Delta_{\Gamma_\pi}(\mathcal{I})(v) = \text{sum of the outgoing slopes of } \mathcal{I}|_{\Gamma_\pi} \text{ at } v$

Canonical divisor of  $\Gamma_\pi$ :  $K_{\Gamma_\pi}(v) = \text{val}_{\Gamma_\pi}(v) + 2g(v) - 2 = -\chi(\check{E}_v)$

Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperplane arrows at } v\} - \#\{\text{polar arrows at } v\})$$

# Back to the inner rates: the Laplacian formula

Natural metric on  $\Gamma_\pi$ :



$$l([v, v']) = \frac{1}{m(v)m(v')}$$

where  $m(v)$  is the multiplicity of  $E_v$  in  $\pi^{-1}(0)$ .

→ Metric on  $\text{NL}(X, 0)$

The inner rates  $\mathcal{I}(E)$  extend to a continuous and piecewise linear map:

$$\mathcal{I}: \text{NL}(X, 0) \longrightarrow \mathbb{R}_{\geq 1}$$



**Laplacian** of  $\mathcal{I}$  on  $\Gamma_\pi$ :  $\Delta_{\Gamma_\pi}(\mathcal{I})(v) = \text{sum of the outgoing slopes of } \mathcal{I}|_{\Gamma_\pi} \text{ at } v$

Canonical divisor of  $\Gamma_\pi$ :  $K_{\Gamma_\pi}(v) = \text{val}_{\Gamma_\pi}(v) + 2g(v) - 2 = -\chi(\check{E}_v)$

Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperplane arrows at } v\} - \#\{\text{polar arrows at } v\})$$

# Back to the inner rates: the Laplacian formula

Natural metric on  $\Gamma_\pi$ :

$$l([v, v']) = \frac{1}{m(v)m(v')}$$

where  $m(v)$  is the multiplicity of  $E_v$  in  $\pi^{-1}(0)$ .

→ Metric on  $\text{NL}(X, 0)$

The inner rates  $\mathcal{I}(E)$  extend to a continuous and piecewise linear map:



$$\mathcal{I}: \text{NL}(X, 0) \longrightarrow \mathbb{R}_{\geq 1}$$

Laplacian of  $\mathcal{I}$  on  $\Gamma_\pi$ :  $\Delta_{\Gamma_\pi}(\mathcal{I})(v) = \text{sum of the outgoing slopes of } \mathcal{I}|_{\Gamma_\pi} \text{ at } v$

**Canonical divisor** of  $\Gamma_\pi$ :  $K_{\Gamma_\pi}(v) = \text{val}_{\Gamma_\pi}(v) + 2g(v) - 2 = -\chi(\check{E}_v)$

Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperplane arrows at } v\} - \#\{\text{polar arrows at } v\})$$



# Back to the inner rates: the Laplacian formula

Natural metric on  $\Gamma_\pi$ :

$$l([v, v']) = \frac{1}{m(v)m(v')}$$

where  $m(v)$  is the multiplicity of  $E_v$  in  $\pi^{-1}(0)$ .

→ Metric on  $\text{NL}(X, 0)$

The inner rates  $\mathcal{I}(E)$  extend to a continuous and piecewise linear map:



$$\mathcal{I}: \text{NL}(X, 0) \longrightarrow \mathbb{R}_{\geq 1}$$

Laplacian of  $\mathcal{I}$  on  $\Gamma_\pi$ :  $\Delta_{\Gamma_\pi}(\mathcal{I})(v) = \text{sum of the outgoing slopes of } \mathcal{I}|_{\Gamma_\pi} \text{ at } v$

Canonical divisor of  $\Gamma_\pi$ :  $K_{\Gamma_\pi}(v) = \text{val}_{\Gamma_\pi}(v) + 2g(v) - 2 = -\chi(\check{E}_v)$

Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperplane arrows at } v\} - \#\{\text{polar arrows at } v\})$$





# Back to the inner rates: the Laplacian formula

## Theorem (Belotto–F–Pichon, 2019)

$$\Delta_{\Gamma_{\pi}}(\mathcal{I})(v) = m(v)(K_{\Gamma_{\pi}}(v) + 2\#\{\text{hyperpl. arrows at } v\} - \#\{\text{polar arrows at } v\})$$

Proof: lifting the formula for  $\text{NL}(\mathbb{C}^2, 0)$  to the singular case, studying the topology and monodromy of the Milnor fiber of a generic linear form, Dehn twists.

Applications:

- Simple explicit computation of the inner rates
- Lê–Greuel–Teissier Formula
- We obtain strong restrictions on the relative positions of arrows.

# Back to the inner rates: the Laplacian formula

## Theorem (Belotto–F–Pichon, 2019)


$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperpl. arrows at } v\} - \#\{\text{polar arrows at } v\})$$

Proof: lifting the formula for  $\text{NL}(\mathbb{C}^2, 0)$  to the singular case, studying the topology and monodromy of the Milnor fiber of a generic linear form, Dehn twists.

Applications:

- Simple explicit computation of the inner rates
- Lê–Greuel–Teissier Formula
- We obtain strong restrictions on the relative positions of arrows.

$$\{(zx^2 + y^3)(x^3 + zy^2) + z^7 = 0\} \subset \mathbb{C}^3$$



# Back to the inner rates: the Laplacian formula

## Theorem (Belotto–F–Pichon, 2019)


$$\Delta_{\Gamma_\pi}(\mathcal{I})(v) = m(v)(K_{\Gamma_\pi}(v) + 2\#\{\text{hyperpl. arrows at } v\} - \#\{\text{polar arrows at } v\})$$

Proof: lifting the formula for  $\text{NL}(\mathbb{C}^2, 0)$  to the singular case, studying the topology and monodromy of the Milnor fiber of a generic linear form, Dehn twists.

Applications:

- Simple explicit computation of the inner rates
- **Lê–Greuel–Teissier Formula**
- We obtain strong restrictions on the relative positions of arrows.

$$\{(zx^2 + y^3)(x^3 + zy^2) + z^7 = 0\} \subset \mathbb{C}^3$$



# Back to the inner rates: the Laplacian formula

## Theorem (Belotto–F–Pichon, 2019)


$$\Delta_{\Gamma_{\pi}}(\mathcal{I})(v) = m(v)(K_{\Gamma_{\pi}}(v) + 2\#\{\text{hyperpl. arrows at } v\} - \#\{\text{polar arrows at } v\})$$

Proof: lifting the formula for  $\text{NL}(\mathbb{C}^2, 0)$  to the singular case, studying the topology and monodromy of the Milnor fiber of a generic linear form, Dehn twists.

Applications:

- Simple explicit computation of the inner rates
- Lê–Greuel–Teissier Formula
- We obtain strong **restrictions on the relative positions of arrows.**

$$\{(zx^2 + y^3)(x^3 + zy^2) + z^7 = 0\} \subset \mathbb{C}^3$$



# Polar exploration

## Question

Given a resolution graph  $\Gamma$ , which configurations of arrows (both **hyperplane** and **polar**) on  $\Gamma$  can be realized by a surface singularity  $(X, 0)$ ?

## First answer (Belotto–F–Pichon 2020)

There are only finitely many possible configurations of arrows on  $\Gamma$ .

In fact, very few possible solutions:  
given the hyperplane arrows, the polar  
arrows are limited by the Laplacian  
formula.

$n = \#V(\Gamma_\pi)$  equations in  $2n$  unknown  
+ geometry of Lipman cones

Example: only one possibility for  $E_8$ .

# Polar exploration

## Question

Given a resolution graph  $\Gamma$ , which configurations of arrows (both **hyperplane** and **polar**) on  $\Gamma$  can be realized by a surface singularity  $(X, 0)$ ?

## First answer (Belotto–F–Pichon 2020)

There are only finitely many possible configurations of arrows on  $\Gamma$ .

In fact, very few possible solutions:  
given the hyperplane arrows, the polar  
arrows are limited by the Laplacian  
formula.

$n = \#V(\Gamma_\pi)$  equations in  $2n$  unknown  
+ geometry of Lipman cones

Example: only one possibility for  $E_8$ .

# Polar exploration

## Question

Given a resolution graph  $\Gamma$ , which configurations of arrows (both **hyperplane** and **polar**) on  $\Gamma$  can be realized by a surface singularity  $(X, 0)$ ?


## First answer (Belotto–F–Pichon 2020)

There are only finitely many possible configurations of arrows on  $\Gamma$ .

In fact, very few possible solutions:  
given the hyperplane arrows, the polar  
arrows are limited by the Laplacian  
formula.

$n = \#V(\Gamma_\pi)$  equations in  $2n$  unknown  
+ geometry of Lipman cones

Example: only one possibility for  $E_8$ .



Motivation: there are two ways to resolve the singularities of surfaces:

- via point blowups (Zariski 1939)  $\longrightarrow$  hyperplane sections
- via Nash transformations (Spivakovsky 1990)  $\longrightarrow$  polar curves

Lê Dũng Tráng (~2000): Are these two methods dual?

Duality between hyperplane sections and polar curves?

In both methods a crucial role is played by minimal singularities.

### Theorem (Spivakovsky 1990)

Let  $(X, 0)$  be a surface germ with a minimal singularity.

Then the topological type of  $(X, 0)$  determines the weighted dual graph of the minimal resolution of  $(X, 0)$  that factors through the blowup at 0 and by its Nash transform, together with the arrows of its hyperplane sections and those of its polar curves.

We have an analogous result for the discriminant curve of a generic projection.  
(Bondil 2003 in the minimal case, Belotto–F–Pichon 2020 in the LNE case)



Motivation: there are two ways to resolve the singularities of surfaces:

- via point blowups (Zariski 1939)  $\longrightarrow$  hyperplane sections
- via Nash transformations (Spivakovsky 1990)  $\longrightarrow$  polar curves

Lê Dũng Tráng ( $\sim 2000$ ): Are these two methods dual?

Duality between hyperplane sections and polar curves?

In both methods a crucial role is played by minimal singularities.

Theorem (Spivakovsky 1990)

Let  $(X, 0)$  be a surface germ with a minimal singularity.

Then the topological type of  $(X, 0)$  determines the weighted dual graph of the minimal resolution of  $(X, 0)$  that factors through the blowup at 0 and by its Nash transform, together with the arrows of its hyperplane sections and those of its polar curves.

We have an analogous result for the discriminant curve of a generic projection.  
(Bondil 2003 in the minimal case, Belotto–F–Pichon 2020 in the LNE case)

Motivation: there are two ways to resolve the singularities of surfaces:

- via point blowups (Zariski 1939)  $\longrightarrow$  hyperplane sections
- via Nash transformations (Spivakovsky 1990)  $\longrightarrow$  polar curves

Lê Dũng Tráng ( $\sim 2000$ ): Are these two methods dual?

Duality between hyperplane sections and polar curves?

In both methods a crucial role is played by minimal singularities.

### Theorem (Spivakovsky 1990)

Let  $(X, 0)$  be a surface germ with a minimal singularity.

Then the topological type of  $(X, 0)$  determines the weighted dual graph of the minimal resolution of  $(X, 0)$  that factors through the blowup at 0 and by its Nash transform, together with the arrows of its hyperplane sections and those of its polar curves.

We have an analogous result for the discriminant curve of a generic projection.  
(Bondil 2003 in the minimal case, Belotto–F–Pichon 2020 in the LNE case)

Motivation: there are two ways to resolve the singularities of surfaces:

- via point blowups (Zariski 1939)  $\longrightarrow$  hyperplane sections
- via Nash transformations (Spivakovsky 1990)  $\longrightarrow$  polar curves

Lê Dũng Tráng ( $\sim 2000$ ): Are these two methods dual?

Duality between hyperplane sections and polar curves?

In both methods a crucial role is played by minimal singularities.

**Theorem (Spivakovsky 1990, Belotto–F–Pichon 2020)**

Let  $(X, 0)$  be a surface germ with a ~~minimal~~ LNE singularity.

Then the topological type of  $(X, 0)$  determines the weighted dual graph of the minimal resolution of  $(X, 0)$  that factors through the blowup at 0 and by its Nash transform, together with the arrows of its hyperplane sections and those of its polar curves and all its inner rates.

We have an analogous result for the discriminant curve of a generic projection. (Bondil 2003 in the minimal case, Belotto–F–Pichon 2020 in the LNE case)