

Please write your proofs neatly in the space provided. Remember that searching the internet and discussing these questions with anyone, except your instructor or fellow students in this class through the general discussion board, is considered academic dishonesty.

(1) The unique positive real number solution to the equation  $x^5 + x = 10$  is irrational.

Hints: You may use the following without justification.

- $\forall n \in \mathbb{N}, \forall a_0, a_1, \dots, a_n \in \mathbb{R}$ , the poly.  $p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$  satisfies  $\forall a, b, y \in \mathbb{R}$  if  $y$  is in between  $p(a)$  and  $p(b)$  then for some  $c$  in between  $a$  and  $b$ ,  $y = p(c)$ .
- Any positive rational number can be represented by a fraction  $\frac{p}{q}$  such that  $p, q \in \mathbb{N}$  and the greatest common divisor of  $p, q$  is 1.
- The function  $f(x) = x^5 + x$  is an increasing function for  $x > 0$ .

*Proof:* There exists a unique positive real number solution  $x$  to the equation  $x^5 + x = 10$ , where  $f(x) = x^5 + x$  is an increasing function for  $x > 0$ , such that  $x$  must satisfy  $1 < x < 2$  because for  $x = 1$ ,  $(1)^5 + (1) = 2 < 10$ , and for  $x = 2$ ,  $(2)^5 + (2) = 34 > 10$ . Suppose that  $x$  is rational. Then  $x$  can be represented by a fraction  $p/q$  such that  $p, q \in \mathbb{N}$  and the greatest common divisor of  $p, q$  is 1. Substitution into the equation yields  $\frac{p^5}{q^5} + \frac{p}{q} = 10$ ,  $p^5 + pq^4 = 10q^5$ ,  $p(p^4 + q^4) = 10q^5$ ,  $pr = 10q^5$  where  $r$  is the integer  $p^4 + q^4$ , so  $p$  divides  $10q^5$ . Since  $p$  and  $q$  do not share any prime divisors,  $p$  must divide 10 so the possible values for  $p$  are 1, 2, 5, and 10.

In the case  $p = 1$ ,  $1 < \frac{1}{q} < 2$ ,  $\frac{1}{2} < q < 1$ . There are no integers in  $(\frac{1}{2}, 1)$ .

In the case  $p = 2$ ,  $1 < \frac{2}{q} < 2$ ,  $1 < q < 2$ . Again, there are no integers in  $(1, 2)$ .

In the case  $p = 5$ ,  $1 < \frac{5}{q} < 2$ ,  $2.5 < q < 5$ . The integers in  $(2.5, 5)$  are 3 and 4. If  $q = 3$ ,  $(\frac{5}{3})^5 + (\frac{5}{3}) = \frac{3530}{243} \neq 10$  and if  $q = 4$ ,  $(\frac{5}{4})^5 + (\frac{5}{4}) = \frac{4405}{1024} \neq 10$ , so  $q$  cannot be 3 or 4.

In the case  $p = 10$ ,  $1 < \frac{10}{q} < 2$ ,  $5 < q < 10$ . The integers in  $(5, 10)$  are 6, 7, 8, and 9, but  $q$  cannot be 6 or 8 since it would share the common divisor 2 with  $p$ , so that leaves 7 and 9. If  $q = 7$ ,  $(\frac{10}{7})^5 + (\frac{10}{7}) = \frac{124010}{16807} \neq 10$  and if  $q = 9$ ,  $(\frac{10}{9})^5 + (\frac{10}{9}) = \frac{165610}{59049} \neq 10$ , so  $q$  cannot be 7 or 9.

Therefore there are no integers  $p$  and  $q$  for  $\frac{p^5}{q^5} + \frac{p}{q} = 10$ , so  $x$  is not rational. That completes the proof  $\square$

(2) For all sets  $S, T$ ,  $S \subseteq T$  if and only if  $\mathcal{P}(S) \subseteq \mathcal{P}(T)$ .

*Proof:* Let  $S, T$  be sets. First let's prove the forward direction. Assume  $S \subseteq T$ . Let  $X$  be an arbitrary element of  $\mathcal{P}(S)$ , then  $X \subseteq S$  so  $X \subseteq T$ . Since  $\mathcal{P}(T)$  is the set of all subsets of  $T$ , then  $X \in \mathcal{P}(T)$ . Since  $X$  was arbitrary, it follows that all elements of  $\mathcal{P}(S)$  are in  $\mathcal{P}(T)$ , so  $\mathcal{P}(S) \subseteq \mathcal{P}(T)$ .

Next let's prove the reverse direction. Assume  $\mathcal{P}(S) \subseteq \mathcal{P}(T)$ . Since  $S \in \mathcal{P}(S)$ , it follows that  $S \in \mathcal{P}(T)$ . Therefore  $S \subseteq T$ . That completes the proof  $\square$

(3) For all integers  $n \geq 0$ ,  $5^{5n+1} + 4^{5n+2} + 3^{5n}$  is a multiple of 11.

*Proof:* Let's prove this by induction. First we must show that  $5^{5n+1} + 4^{5n+2} + 3^{5n}$  is a multiple of 11 when  $n = 0$ . So  $5^{5(0)+1} + 4^{5(0)+2} + 3^{5(0)} = 22$ , which is a multiple of 11. That establishes the base case. Let  $k \geq 0$  be an integer, and assume that  $5^{5k+1} + 4^{5k+2} + 3^{5k}$  is a multiple of 11. Then  $\exists j \in \mathbb{Z}$  such that  $5^{5k+1} + 4^{5k+2} + 3^{5k} = 11j$ . This is the inductive hypothesis.

We must show that  $5^{5(k+1)+1} + 4^{5(k+1)+2} + 3^{5(k+1)} = 11z$  for some integer  $z$ . Then

$$\begin{aligned}
& 5^{5(k+1)+1} + 4^{5(k+1)+2} + 3^{5(k+1)} = 5^{(5k+1)+5} + 4^{(5k+2)+5} + 3^{5k+5} \\
&= 5^5 \cdot 5^{5k+1} + 4^5 \cdot 4^{5k+2} + 3^5 \cdot 3^{5k} \\
&= 3125 \cdot 5^{5k+1} + 1024 \cdot 4^{5k+2} + 243 \cdot 3^{5k} \\
&= 3124 \cdot 5^{5k+1} + 1023 \cdot 4^{5k+2} + 242 \cdot 3^{5k} + (5^{5k+1} + 4^{5k+2} + 3^{5k}) \\
&= 3124 \cdot 5^{5k+1} + 1023 \cdot 4^{5k+2} + 242 \cdot 3^{5k} + 11j \quad (\text{by the inductive hypothesis}) \\
&= 11(248 \cdot 5^{5k+1} + 93 \cdot 4^{5k+2} + 22 \cdot 3^{5k} + j) \\
&= 11z, \text{ where } z = 248 \cdot 5^{5k+1} + 93 \cdot 4^{5k+2} + 22 \cdot 3^{5k} + j
\end{aligned}$$

and  $z \in \mathbb{Z}$  since it is the sum of products of integers. Since we have proved the basis step and the inductive step, we conclude that for all integers  $n \geq 0$ ,  $5^{5n+1} + 4^{5n+2} + 3^{5n}$  is a multiple of 11. That completes the proof by induction  $\square$

(4) For all positive integers  $n$ ,  $\sum_{k=1}^n k(C(n, k))^2 = nC(2n - 1, n - 1)$ .

Hint: Consider forming committees of  $n$  people (with a chairperson) that are chosen from a group of  $n$  Oregonians and  $n$  Washingtonians such that the chairperson is an Oregonian.

*Proof:* Let  $A$  and  $B$  be disjoint sets of  $n$  distinct elements where  $n \geq 1$  is an integer. The number of ways to form subsets of  $n$  elements where at least 1 element must be from set  $A$  is  $nC(2n - 1, n - 1)$  since there are  $n$  ways to choose 1 element from  $A$  and  $2n - 1$  elements remaining to choose  $n - 1$  elements.

Another way to count this is to count the number that involve  $k$  elements from set  $A$  and  $n - k$  elements from set  $B$ , and sum over the possible values of  $k$ , which are  $k = 1$  to  $k = n$ .

There are  $k$  ways to choose the 1 required element from set  $A$ ,  $C(n, k)$  ways to select  $k$  elements from set  $A$ , and  $C(n, n - k)$  ways to select  $n - k$  elements from set  $B$ , for  $k = 1, 2, \dots, n$ .

So the number of ways is  $\sum_{k=1}^n kC(n, k)C(n, n - k)$ . We know that  $C(n, k) = C(n, n - k)$  because the number of ways to choose  $k$  elements from  $n$  objects is the same as choosing the  $n - k$  elements to be left out. Therefore, the summation can be written as  $\sum_{k=1}^n k(C(n, k))^2$ , which we have shown to be equal to  $nC(2n - 1, n - 1)$   $\square$