



Genomic Selection in the UF Strawberry Breeding Program

Salvador A. Gezan, Luis Osorio, Sujeet Verma and Vance Whitaker
University of Florida

Rios - Advanced Plant Breeding, Nov 1, 2018




RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

Breeding in Florida since 1948





Florida Acreage 2018-19





'Florida Radiance'




'Florida Beauty'





'Florida Brilliance'








Breeder ← Nursery Grower ← Fruit Grower ← Shipper ← Wholesaler/
Retailer ← Consumer

Clonal multiplication	Disease resistance	Firmness	Fruit Size	Sweetness
Breakage resistance	Yield	Size consistency	Gloss	Internal color
Low chilling requirement	Rain resistance	Shape	Color uniformity	Freshness
Virus tolerance	Low cull rate	Color after cooling	Shelf life	
	Skin toughness			

$$r_a = -0.76$$

DR QTL on UF Genetic Map



UF Strawberry Breeding Cycle



2014-15

Cross

MASS



2015-16

Seedlings

2016-17

Adv. Sel.

2017-18


Cross



RosBREED

DISEASE RESISTANCE × HORTICULTURAL QUALITY

Genomic Selection Process



(From Heffner *et al.*, (2009))



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

Parent GS Research Objectives

- Explore GS models in true validations
- Effects of multiple training cycles on predictive ability (PA)
- Effects of marker density on PA
- Integrating GS into the breeding strategy



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

GS Field Trials

Experimental Design

- Two advanced selection trials (Yr. 1, Yr. 2)
RCBD, 5 blocks, 300-400 cloned genotypes

Phenotyping


Trait	Unit
Average Fruit Wt (AWT)	grams
Soluble Solids (Brix) (SSC)	%
Early Marketable Yield (EMY)	grams
Total Marketable Yield (TMY)	grams
Total Cull Fruit (TC)	%



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

GS Methods

Genotyping



90K IStraw90 Axiom® SNP array

Affimetrix & RosBREED

(Bassil and Davis et al. 2015)

- Marker Quality Control
- ~9000 SNPs



RosBREED

DISEASE RESISTANCE × HORTICULTURAL QUALITY

Analyses

- Models GBLUP, Bayes B, Bayes C, and RKHS
- BGLR (Perez P., de los Campos, G. 2010)

Comparison of GS Models

- Predictive Ability: $\text{corr}(y, \hat{g})$
- PBLUP: traditional pedigree-based estimates (no markers)
- Genetic Gain Efficiency from Parent Selection (5%, 10%):
 $\hat{g}_{inc}/\hat{g}_{com}$


GS Results

Predictive Ability - Year 1 to predict Year 2-

Trait	PBLUP	GS Models			
		GBLUP	Bayes B	Bayes C	RKHS
AWT	0.44	0.49	0.49	0.49	0.52
SSC	0.37	0.43	0.44	0.44	0.45
TMY	0.24	0.31	0.35	0.34	0.33
TC	0.14	0.32	0.35	0.35	0.32




GS Genetic Gain Example



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

Parental GS Genetic Gains




■ 5% ■ 10%

RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY




Multiple Years of Training

Bayes B ~ 9,000 SNPs




RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

Random Marker Selection



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

GS for Parent Selection



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY


How This Works Practically

- **Sept - Oct:** Predict BVs for *Objectively Measured Traits* for Advanced Selections for which no data is available (selected as seedlings previous year)
- **Nov - Dec:** Early-season field observations for all *Visually Evaluated Traits*
- **Jan - Feb:** Select 5-7 new parents for immediate crossing (typically 35-45 total parents per year)
- **Outcome:** for a portion of the parents, breeding cycle reduced from 3 years (minimum) to 2 years



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

GS for Parent Selection



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

Can we use GS for seedling selection?

- Explore GS within families to predict seedling performance
- How do we evaluate the performance of this method?
- How would we integrate GS with MASS for major loci?



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

Within-Family GS Experiment

Trial Design

Planted: 10/2018
RCBD: 3 Replications
Families: 6
Seedlings: 75/Fam
Total N: 450 seedlings

Females	Males			
	A	B	C	D
A	X		X	X
B			X	X
C				X
D				

Model Evaluation (02/2018)

TNT

$$\underline{A/* + B/* = 300}$$

TST


$$\underline{A^*B = 75}$$

TRN = Families with Common Parents (4)



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

Within-Family GS Experiment



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

Within-Family GS Experiment

	AWT	EMY	SSC	TC	FRM
Seedling Selection (PA)	0.28	0.16	0.13	0.28	0.24
Gain Over Family Mean (top 10%)	4.4%	14.1%	9.3%	14.1%	15.6%



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

Within-Family GS Experiment

	AWT	EMY	SSC	TC	FRM
Seedling Selection (PA)	0.28	0.16	0.13	0.28	0.24
Gain Over Family Mean (10%)	4.4%	14.1%	9.3%	14.1%	15.6%
Yearly Breeding Gain Last 5 Years (%)	2.7%	5.7%	-1.2%	0.5%	--



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

Within-Family GS Experiment


	AWT	EMY	SSC	TC	FRM
Seedling Selection (PA)	0.28	0.16	0.13	0.28	0.24
Gain Over Family Mean (10%)	4.4%	14.1%	9.3%	14.1%	15.6%
Yearly Breeding Gain Last 5 Years (%)	2.7%	5.7%	-1.2%	0.5%	--

For Seedling GS in UF strawberry program only 500 SNPs needed




RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY

Hypothetical WF GS for Cross



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY



RosBREED

DISEASE RESISTANCE × HORTICULTURAL QUALITY

Summary

- **GS for Parent Selection** allows accelerated use of new parents in crossing, reducing length of cycle
- **GS for Seedling Selection** could increase gains for complex traits if cost-effective, low-density genotyping available



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY



RosBREED
DISEASE RESISTANCE × HORTICULTURAL QUALITY