#### 1

# EE5600 - Introduction to AI & ML Assignment 1

## Virendra Patil AI20MTECH01003

Abstract—Given the vector coordinates of the angular points of the triangle, we need to find the area of the triangle. In vector space, the area of parallelogram formed by any two vectors is given by the magnitude of the cross product of those two vectors. The area of triangle whose two sides are represented by vectors is nothing but half the area of parallelogram formed by the respective vectors. In essence, area of triangle whose two sides are represented by two vectors u and v is given by  $\frac{1}{2} \times ||(u \times v)||$ , where " $u \times v$ " represents cross product of two vectors. The python implementation of the said method is available at https://github.com/virendra-patil/EE5600-Intro-to-AI-ML/tree/main/Assignment 1.

#### 1 Problem Statement

Vector2, Example II, Problem 4: Find the area of the triangle whose coordinates of angular points are respectively,

$$\begin{pmatrix} a \\ b+c \end{pmatrix}$$
,  $\begin{pmatrix} a \\ b-c \end{pmatrix}$  and  $\begin{pmatrix} -a \\ c \end{pmatrix}$ .

#### 2 Solution

We solve the above stated problem in two steps. First, we find two vectors representing sides of the given triangle. In second step we calculate area of parallelogram and halve it to obtain the area of triangle. Initially we propose a generalise solution and extend it by solving the same problem using example coordinates.

Step I:

Let,

$$\mathbf{X} = \begin{pmatrix} a \\ b+c \end{pmatrix}, \ \mathbf{Y} = \begin{pmatrix} a \\ b-c \end{pmatrix} \text{ and } \mathbf{Z} = \begin{pmatrix} -a \\ c \end{pmatrix}$$
 (1)

Rewriting X, Y and Z as product of a matrix and a vector, we get,

$$\mathbf{X} = \begin{bmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} \end{bmatrix}$$

$$\mathbf{Y} = \begin{bmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} \end{bmatrix}$$

$$\mathbf{Z} = \begin{bmatrix} \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{bmatrix} \end{bmatrix}$$
(2)

As X, Y and Z represent coordinates of angular points of the triangle, by vector calculus X-Y, Y-Z and X-Z would represent the sides of the triangle. To calculate area of triangle we need any two sides. Here we consider X-Y and Y-Z.

$$\mathbf{X} - \mathbf{Y} = \begin{bmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} \end{bmatrix}$$
(3)

$$\mathbf{X} - \mathbf{Y} = \begin{bmatrix} \begin{pmatrix} 1 - 1 & 0 - 0 & 0 - 0 \\ 0 - 0 & 1 - 1 & 1 - (-1) \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} \end{bmatrix}$$
(4)

$$\mathbf{X} - \mathbf{Y} = \begin{bmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{bmatrix} \end{bmatrix} \tag{5}$$

$$\mathbf{X} - \mathbf{Y} = \begin{pmatrix} 0 \\ 2c \end{pmatrix} \tag{6}$$

$$\mathbf{Y} - \mathbf{Z} = \begin{bmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} - \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{bmatrix} \end{bmatrix} \quad (7)$$

$$\mathbf{Y} - \mathbf{Z} = \begin{bmatrix} \begin{pmatrix} 1 - (-1) & 0 - 0 & 0 - 0 \\ 0 - 0 & 1 - 0 & -1 - 1 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} \end{bmatrix}$$
(8)

$$\mathbf{Y} - \mathbf{Z} = \begin{bmatrix} \begin{pmatrix} 2 & 0 & 0 \\ 0 & 1 & -2 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{bmatrix} \end{bmatrix} \tag{9}$$

$$\mathbf{Y} - \mathbf{Z} = \begin{pmatrix} 2a \\ b - 2c \end{pmatrix} \tag{10}$$

### Step II:

Now we calculate area of triangle formed by the two sides of the triangle. The area is given by half the area of parallelogram. Area of parallelogram is the magnitude of cross product of two vectors. Hence, from 6 and 10,

Area of Triangle =  $\frac{1}{2} \times$  Area of Parallelogram

(11)

$$= \frac{1}{2} \times ||(\mathbf{X} - \mathbf{Y}) \times (\mathbf{Y} - \mathbf{Z})|| \quad (12)$$

$$= \frac{1}{2} \times \left\| \begin{pmatrix} 0 \\ 2c \end{pmatrix} \times \begin{pmatrix} 2a \\ b - 2c \end{pmatrix} \right\| \tag{13}$$

$$= \frac{1}{2} \times \left\| \begin{pmatrix} 0 \\ 0 \\ 4ac \end{pmatrix} \right\| \tag{14}$$

$$= \frac{1}{2} \times \sqrt{0^2 + 0^2 + (4ac)^2}$$
 (15)

$$=\frac{1}{2}\times\sqrt{16a^2c^2}\tag{16}$$

$$= \frac{1}{2} \times 4ac \text{ unit}^2 \tag{17}$$

$$= 2ac \text{ unit}^2 \tag{18}$$




Fig. 1: Plot representing the triangle formed by angular coordinates

$$\mathbf{X} - \mathbf{Y} = \begin{pmatrix} 0 \\ 2c \end{pmatrix}$$

$$= \begin{pmatrix} 0 \\ 16 \end{pmatrix}$$
(19)

$$\mathbf{Y} - \mathbf{Z} = \begin{pmatrix} 2a \\ b - 2c \end{pmatrix}$$

$$= \begin{pmatrix} 4 \\ -12 \end{pmatrix}$$
(20)

Area of Triangle = 
$$\frac{1}{2} \times 4ac$$
 (21)

$$= \frac{1}{2} \times 4 \times 2 \times 8 \tag{22}$$

$$= 32 \text{ unit}^2$$
 (23)

#### 3 Numerical Example

Lets consider a example coordinate system where a = 2, b = 4 and c = 8. Given a, b and c we can calculate X-Y and Y-Z using equations 6 and 10.




Fig. 2: Plot representing the triangle formed by angular coordinates