


Física Estelar

Lluís Galbany, Ed. Mecenas (#16)

Inma Domínguez, Ed. Mecenas (#17)


Antonio García, Ed. Mecenas (#16)



Curso 2019-2020


Syllabus

Aim: to understand the structure and evolution of stars, and their observational properties, using known laws of physics.



Stellar structure and evolution
(O. R. Pols)

Understanding Stellar Evolution
(Henny J. G. L. M. Lamers & Emily M. Levesque)



Goals

In this course we will:

- understand the global properties of stars: energetics and timescales
- study the micro-physics relevant for stars: the equation of state, nuclear reactions, energy transport and opacity
- derive the equations necessary to model the internal structure of stars
- examine (quantitatively) the properties of simplified stellar models
- survey (mostly qualitatively) how stars of different masses evolve, and the endpoints of stellar evolution (white dwarfs, neutron stars)
- discuss a few ongoing research areas in stellar evolution

Syllabus

Tema 1: Estructura Estelar

Principios de conservación y ecuaciones de estructura estelar. Condiciones de contorno y métodos de resolución. Ecuación de estado y opacidad. Transporte de energía: Radiación, convección y conducción. Criterios de estabilidad convectiva.

Tema 2: Fuentes de Energía Estelar

Teorema del Virial. Reacciones termonucleares y ritmos de reacción. Principales cadenas y ciclos de combustión nuclear. Otros procesos nucleares de interés astrofísico.

Tema 3: Evolución Estelar

Formación estelar y pre-secuencia principal. Límites de masa estelar: enanas marrones y planetas. Edad cero y secuencia principal. Estimación de edades de cúmulos estelares. Evolución en la rama de las gigantes: estrellas RGB y AGB. Formación de enanas blancas. Estrellas masivas y supernovas de colapso gravitatorio.

Tema 4: Objetos compactos

Evolución de enanas blancas. Ecuación de Volkov-Openheimer. Estrellas de neutrones: ecuación de estado. Agujeros negros. Aplicaciones.

Tema 5: Evolución Estelar en Sistemas Binarios.

Supernovas termonucleares: Aplicaciones cosmológicas. Binarias cataclísmicas. Novas. Erupciones de rayos X. Estrellas gigantes binarias y anomalías químicas.

Tema 6: Pulsaciones Estelares.

Pulsaciones esféricas adiabáticas y no adiabáticas. Mecanismos de pulsación. La banda de inestabilidad en el diagrama HR. Oscilaciones no radiales. Astroismología

1. Introduction

What is a star?

A star is...

Una masa de gas que se encuentra en equilibrio entre la presión hidrostática y la de radiación

Cuerpo gaseoso que emite radiación

Grandes bolas de plasma que emiten energía y permiten la vida en la Tierra

Reactor nuclear que eventualmente explota o se apaga

Cuerpo celeste alrededor del cual orbitan planetas

Causante de las quemaduras solares

Objeto al que se le piden deseos

What is a star?

A star is an object that:

- (1) radiates energy from an internal source; and
- (2) is bound by its own gravity.

Stars must evolve: have *life* and *death*.



Stars can have only a limited range of masses: $0.1 \lesssim M_{\odot} \lesssim 100$



Fundamental properties of stars

The **Sun**, our nearest neighbor

Standard parameters of the Sun:

$$\boxed{\begin{aligned}M_{\odot} &\sim 1.99 \times 10^{33} \text{ g} \\R_{\odot} &\sim 6.96 \times 10^{10} \text{ cm} \\L_{\odot} &\sim 3.84 \times 10^{33} \text{ ergs}^{-1}\end{aligned}}$$

$$L = 4\pi R^2 \sigma T_{\text{eff}}^4 \longrightarrow T_{\odot, \text{eff}} \sim 5777 \text{ K}$$


Solar **abundances** ($X + Y + Z = 1$) *(...also v_{rotation})*

Nr	Element	Z	m (AMU)	Abund	Nr	Element	Z	m (AMU)	Abund
1	H	1	1.0079	0.00	6	N	7	14.007	-3.95
2	He	2	4.0026	-1.01	7	Mg	12	24.305	-4.42
3	O	8	15.999	-3.07	8	Si	14	28.086	-4.45
4	C	6	12.011	-3.44	9	S	16	32.066	-4.79
5	Ne	10	20.180	-3.91	10	Fe	26	55.847	-4.46

$$H : He : C + N + O + Ne : \text{rest} = 0.70 : 0.28 : 0.016 : 0.003$$


Effective temperature

Radiation curves of Blackbodies: $F = \sigma T^4 = \sigma(T_{\text{eff}}^4)$



Effective temperature

Radiation curves of Blackbodies: $F = \sigma T^4 = \sigma(T_{\text{eff}}^4)$



Fundamental properties of stars

The **Sun**, our nearest neighbor

Standard parameters of the Sun:

$$\boxed{M_{\odot} \sim 1.99 \times 10^{33} \text{ g}}$$
$$R_{\odot} \sim 6.96 \times 10^{10} \text{ cm}$$
$$L_{\odot} \sim 3.84 \times 10^{33} \text{ ergs}^{-1}$$


$$L = 4\pi R^2 \sigma T_{\text{eff}}^4 \longrightarrow T_{\odot, \text{eff}} \sim 5777 \text{ K}$$

Solar **abundances** ($X + Y + Z = 1$) *(...also v_{rotation})*

Nr	Element	Z	m (AMU)	Abund	Nr	Element	Z	m (AMU)	Abund
1	H	1	1.0079	0.00	6	N	7	14.007	-3.95
2	He	2	4.0026	-1.01	7	Mg	12	24.305	-4.42
3	O	8	15.999	-3.07	8	Si	14	28.086	-4.45
4	C	6	12.011	-3.44	9	S	16	32.066	-4.79
5	Ne	10	20.180	-3.91	10	Fe	26	55.847	-4.46


$$\text{H : He : C + N + O + Ne : rest} = 0.70 : 0.28 : 0.016 : 0.003$$

Sun abundance




$A = \log(n_x/n_H) + 12$ (log of number of atoms per 10^{12} H atoms)
also used number of atoms per 10^6 Si atoms

Sun abundance



$A = \log(n_x/n_H) + 12$ (log of number of atoms per 10^{12} H atoms)
also used number of atoms per 10^6 Si atoms


Abundances of nuclei on the chart of nuclides:



Observations of stellar parameters

Astronomical observations can yield information about stellar fundamental quantities:

1. Distance - parallax method



Star


$$d \sim \frac{R}{\theta}$$



Observations of stellar parameters

Astronomical observations can yield information about stellar fundamental quantities:

1. Distance - parallax method




$$d \sim \frac{R}{\theta}$$



Observations of stellar parameters

Astronomical observations can yield information about stellar fundamental quantities:

1. Distance - parallax method




$$d \sim \frac{R}{\theta}$$



Observations of stellar parameters

Astronomical observations can yield information about stellar fundamental quantities:


1. Distance - parallax method



Observations of stellar parameters

Astronomical observations can yield information about stellar fundamental quantities:


1. Distance - parallax method



Observations of stellar parameters

Astronomical observations can yield information about stellar fundamental quantities:


1. Distance - parallax method



Observations of stellar parameters

Astronomical observations can yield information about stellar fundamental quantities:

1. Distance - parallax method




$$d \sim \frac{R}{\theta} \quad (\theta \text{ in radians})$$



Observations of stellar parameters

Astronomical observations can yield information about stellar fundamental quantities:

1. Distance - parallax method



$$d \sim \frac{R}{\theta} \quad (\theta \text{ in radians})$$



$$R = 1.5 \times 10^{13} \text{ cm}, 1'' = 4.85 \times 10^{-6} \text{ radians}$$


so a parallax of $1''$ = distance of $3.09 \times 10^{18} \text{ cm} = 1 \text{ parsec}$

Observations of stellar parameters

2. Angular radius


With interferometry, for stars sufficiently extended in the sky

$$\theta = \frac{R}{d} \quad \xrightarrow{L = 4\pi R^2 \sigma T_{\text{eff}}^4} \quad \sigma T_{\text{eff}}^4 = \frac{f_{\text{bol}}}{\theta^2}$$



Observations of stellar parameters

VLT interferometer, Cerro Paranal Observatory, Chile



Observations of stellar parameters

3. Mass

The most accurate mass measurements for stars come from binaries.

$$M_1 a_1 = M_2 a_2$$


Center of mass

and

$$\left(\frac{2\pi}{P}\right)^2 = G \frac{(M_1 + M_2)}{a^3}$$

$$= G \frac{(M_1 + M_2)}{(a_1 + a_2)^3}$$

Kepler's 3rd law



(requires orbits and radial velocities for both stars...)

Observations of stellar parameters

4. Luminosity

The luminosity (L) of a star can be derived from its distance and its radiative flux measured at Earth using the relations:

$$L = 4\pi d^2 f_{\text{bol}} = 4\pi R^2 \sigma T_{\text{eff}}^4 \quad \text{or} \quad \frac{L}{L_{\odot}} = \left(\frac{R}{R_{\odot}} \right)^2 \left(\frac{T_{\text{eff}}}{5777\text{K}} \right)^4$$

where


f_{bol} = flux integrated over all wavelengths and corrected for ISM and atmospheric attenuation

T_{eff} = approximation of the gas temperature in the photosphere of the star at optical depth $\tau \approx 1$

Observations of stellar parameters

5. Magnitude and color

Magnitudes reflect the energy flux received on Earth in different wavelength bands.



Color indices are defined as the ratio of the flux in two wavelength bands, and often expressed as the difference between two apparent magnitudes (m_λ) in different filters

Observations of stellar parameters

$$m_\lambda = -2.5 \times \log \left(\frac{f_\lambda}{f_{0,\lambda}} \right)$$

Note: a *lower* value of m_λ corresponds to a *larger* flux

where f_λ is in $\text{erg cm}^{-2} \text{ s}^{-1}$ and $f_{0,\lambda}$ is the reference flux at that λ

Magnitude systems (ref. fluxes) typically used are :

Vega, star

$$f_0(\text{U|B|V}) = 4.2|6.4|3.8 \times 10^{-8} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ nm}^{-1}$$


AB, flat $3.63 \times 10^{-20} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1} = 3631 \text{ Jy}$

$$AB_{\text{mag}} = -2.5 \times \log F_\nu - 48.6$$


ST, flat $3.63 \times 10^{-9} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$

$$ST_{\text{mag}} = -2.5 \times \log F_\lambda - 21.1$$

Observations of stellar parameters



Observations of stellar parameters



Observations of stellar parameters

Adopting the Johnson UBVRI filters, $m_V = V = -2.5 \times \log \left(\frac{f_V}{f_{0,V}} \right)$

B-V color index is $B - V = -2.5 \times \log \left(\frac{f_B}{f_V} \right)$ [+ difference in $f_{0,\lambda}$]
Ex. Calculate term
for Vega and AB.

Colors are defined as BLUE-RED mags. Following BB radiation, a **larger** (redder) value for color corresponds to a **lower** (cooler) T_{eff}

If a star's distance is known we can calculate its **absolute magnitude** M_λ (the m_λ the star would have at a distance of 10pc):

$$M_\lambda = m_\lambda + 5 - 5 \times \log(d)$$

The **absolute bolometric magnitude** M_{bol} describes the flux of the star integrated over all wavelengths at a distance of 10pc:


$$M_{\text{bol}} = -2.5 \times \log \left(\frac{L}{L_\odot} \right) + 4.74$$

and $M_{\text{bol}} - M_\lambda = BC_\lambda$, **the bolometric correction**

Observations of stellar parameters

6... Other spectral parameters

gravity, chemical abundances, Teff, rotational velocity...



Spectra of Stars with Different Metal Content

Observations of stellar parameters

So far, we listed *surface* stellar parameters

We need a theory of stellar structure to derive the internal properties of a star.

However, direct windows on the *interior* of a star exist:

Neutrinos that escape without interaction. Only detected from the Sun

Seismology, the frequency of different stellar oscillations contains information about the speed of sound waves inside the star, so do density and temperature.

The Structure of Stars

To get started...

Let's define some standard regions in a star...



The Structure of Stars

To get started...

Let's define some standard regions in a star...

a) **core** - site of current/former fusion



The Structure of Stars

To get started...

Let's define some standard regions in a star...

- a) **core** - site of current/former fusion
- b) **shell zone** - fusion occurring
in shell(s) around core




The Structure of Stars

To get started...

Let's define some standard regions in a star...

- a) **core** - site of current/former fusion
- b) **shell zone** - fusion occurring
in shell(s) around core
- c) **envelope** - region between
shell and atmosphere




The Structure of Stars

To get started...

Let's define some standard regions in a star...

- a) **core** - site of current/former fusion
- b) **shell zone** - fusion occurring in shell(s) around core
- c) **envelope** - region between shell and atmosphere
- d) **atmosphere** - optical depth is $\tau \lesssim 10$;
radiation can escape




The Structure of Stars

To get started...

Let's define some standard regions in a star...

- a) **core** - site of current/former fusion
- b) **shell zone** - fusion occurring in shell(s) around core
- c) **envelope** - region between shell and atmosphere
- d) **atmosphere** - optical depth is $\tau \lesssim 10$; radiation can escape
- e) **chromosphere/corona** - region in cool star above atmosphere where T rises due to shocks




The Structure of Stars

To get started...

Let's define some standard regions in a star...

- a) **core** - site of current/former fusion
- b) **shell zone** - fusion occurring in shell(s) around core
- c) **envelope** - region between shell and atmosphere
- d) **atmosphere** - optical depth is $\tau \lesssim 10$;
radiation can escape
- e) **chromosphere/corona** - region in cool star above atmosphere where T rises due to shocks
- f) **wind** - region where gas escapes at $>> v_{\text{esc}}$



Stellar Evolution in a Nutshell

— stars are spheres of gas held together by their own gravitational attraction

Stellar Evolution in a Nutshell

- stars are spheres of gas held together by their own gravitational attraction
- they stay stable for a remarkably long time; gravity is balanced by gas pressure + radiation pressure

Stellar Evolution in a Nutshell

- stars are spheres of gas held together by their own gravitational attraction
- they stay stable for a remarkably long time; gravity is balanced by gas pressure + radiation pressure
- this implies a strong T and density gradient, with both much higher in the center; energy flows outward, leaves the star at the photosphere, and is radiated into space

Stellar Evolution in a Nutshell

- stars are spheres of gas held together by their own gravitational attraction
- they stay stable for a remarkably long time; gravity is balanced by gas pressure + radiation pressure
- this implies a strong T and density gradient, with both much higher in the center; energy flows outward, leaves the star at the photosphere, and is radiated into space
- stars must replenish this energy loss, demanding an internal energy source: nuclear fusion in the center of the star, where T and ρ are sufficiently high

Stellar Evolution in a Nutshell

- stars are spheres of gas held together by their own gravitational attraction
- they stay stable for a remarkably long time; gravity is balanced by gas pressure + radiation pressure
- this implies a strong T and density gradient, with both much higher in the center; energy flows outward, leaves the star at the photosphere, and is radiated into space
- stars must replenish this energy loss, demanding an internal energy source: nuclear fusion in the center of the star, where T and ρ are sufficiently high
- if fusion stops the star contracts, releasing E_{pot} . Some is used to continue the energy flow, while the rest raises T in the center of the star. Once T is high enough the next fusion reaction starts.

Stellar Evolution in a Nutshell

- stars are spheres of gas held together by their own gravitational attraction
- they stay stable for a remarkably long time; gravity is balanced by gas pressure + radiation pressure
- this implies a strong T and density gradient, with both much higher in the center; energy flows outward, leaves the star at the photosphere, and is radiated into space
- stars must replenish this energy loss, demanding an internal energy source: nuclear fusion in the center of the star, where T and ρ are sufficiently high
- if fusion stops the star contracts, releasing E_{pot} . Some is used to continue the energy flow, while the rest raises T in the center of the star. Once T is high enough the next fusion reaction starts.
- this cycle continues until fusion fuel is exhausted. Most stars stay in a perfect hydrostatic balance overall during this evolution.

Stellar Evolution in a Nutshell



- stars are spheres of gas held together by their own gravitational attraction
- they stay stable for a remarkably long time; gravity is balanced by gas pressure + radiation pressure
- this implies a strong T and density gradient, with both much higher in the center; energy flows outward, leaves the star at the photosphere, and is radiated into space
- stars must replenish this energy loss, demanding an internal energy source: nuclear fusion in the center of the star, where T and ρ are sufficiently high
- if fusion stops the star contracts, releasing E_{pot} . Some is used to continue the energy flow, while the rest raises T in the center of the star. Once T is high enough the next fusion reaction starts.
- this cycle continues until fusion fuel is exhausted. Most stars stay in a perfect hydrostatic balance overall during this evolution.
- once the nuclear energy source is gone, gravity wins. In low-mass stars the core compresses and becomes a white dwarf. In high-mass stars the core collapses, producing a neutron star or black hole, and ejects the outer layers, producing a supernova.

Stellar properties relations

Mass-luminosity & Mass-radius relations


All 3 parameters are needed, so only possible for *main sequence stars** from double-lined eclipsing binaries

*stars that fuse H in their center



Stellar Evolution in a Nutshell


Mass-luminosity relation



Relation is only valid for stars during core H fusion (red giants/supergiants, degenerate stars, etc., do not follow the relation)

Stellar Evolution in a Nutshell

Mass-luminosity relation




Relation is only valid for stars during core H fusion (red giants/supergiants, degenerate stars, etc., do not follow the relation)

For $M > 6M_{\odot}$, slope flattens due to increasing role of radiation pressure

Stellar Evolution in a Nutshell

Mass-luminosity relation




Relation is only valid for stars during core H fusion (red giants/supergiants, degenerate stars, etc., do not follow the relation)


For $M > 6M_{\odot}$, slope flattens due to increasing role of radiation pressure

For $M < 0.6M_{\odot}$, slope flattens due to increasing role of convection throughout the star and changes in opacity


The Hertzsprung-Russell Diagram




The Hertzsprung-Russell Diagram




The Hertzsprung-Russell Diagram




The Hertzsprung-Russell Diagram




The Hertzsprung-Russell Diagram




The Hertzsprung-Russell Diagram




The Hertzsprung-Russell Diagram




The Hertzsprung-Russell Diagram




The Hertzsprung-Russell Diagram




The Hertzsprung-Russell Diagram



The Hertzsprung-Russell Diagram



The Color-Magnitude Diagram



The Color-Magnitude Diagram






Figure courtesy of ESA.

The Color-Magnitude Diagram

Gaia





CMD and HRD



Left: Reproduced from Buonanno, R et al, A&A, Vol. 290, p.69-103 (1994), reproduced with permission. © ESO.

Right: Reproduced from Johnson, H. L. and Sandage, A. R., 'Three-Color Photometry in the Globular Cluster M3', Astrophysical Journal, vol. 124, p.379, 1956.



CMD and HRD



Left: Reproduced from Buonanno, R et al, A&A, Vol. 290, p.69-103 (1994), reproduced with permission. © ESO.

Right: Reproduced from Johnson, H. L. and Sandage, A. R., 'Three-Color Photometry in the Globular Cluster M3', Astrophysical Journal, vol. 124, p.379, 1956.

CMD and HRD






1. on the blue/hot side stars cover a small $(B-V)$, large T_{eff} range
2. on the red/cool side, stars cover a small T_{eff} , large $(B-V)$ range
3. horizontal branch curves down strongly in the CMD

Left: Reproduced from Buonanno, R et al, A&A, Vol. 290, p.69-103 (1994), reproduced with permission. © ESO.

Right: Reproduced from Johnson, H. L. and Sandage, A. R., 'Three-Color Photometry in the Globular Cluster M3', Astrophysical Journal, vol. 124, p.379, 1956.


The Color-Magnitude Diagram



As the clusters increase in age:

- the MS gets shorter
- the ratio of red giants to MS stars increases

The Color-Magnitude Diagram



Stellar populations



Population I: in the galactic disk, spiral arms and open clusters. $t < 10^9$ yr, $Z \sim 0.5 - 1 Z_\odot$

Population II: in the galactic disk, halo, globular clusters. $t \sim 10^{10}$ yr, $Z \sim 0.01 - 0.1 Z_\odot$

Population III: halo? $Z \sim 0?$

Basic assumptions: Th. Stell. Evo.

Basic assumptions: Th. Stell. Evo.

1. Isolated in space

- Their structure and evolution depend only on intrinsic properties (mass and composition)
- For binaries and dense clusters, evolution can be influenced by interaction

Basic assumptions: Th. Stell. Evo.

1. Isolated in space

- Their structure and evolution depend only on intrinsic properties (mass and composition)
- For binaries and dense clusters, evolution can be influenced by interaction

2. Formed with a homogeneous composition

- Because molecular clouds are well-mixed
- Assuming a quasi-solar composition: $X=0.706$ $Y=0.28$, $Z=0.014$

Basic assumptions: Th. Stell. Evo.

1. Isolated in space

- Their structure and evolution depend only on intrinsic properties (mass and composition)
- For binaries and dense clusters, evolution can be influenced by interaction

2. Formed with a homogeneous composition

- Because molecular clouds are well-mixed
- Assuming a quasi-solar composition: $X=0.706$ $Y=0.28$, $Z=0.014$

3. Have spherical symmetry

- Promoted by self-gravity.
- Possible deviation from non-central forces: No rotation, no magnetic fields

Exercises

- 1.1** Stars are assumed to be isolated in space. The star closest to the sun, Proxima Centauri, is 4.3 light-years away. How many solar radii is that? How it affects the Sun gravitationally? How this force compares to the attractive gravitational force the Sun suffers from Jupiter?
- 1.2** Stars are assumed to form with a uniform composition. What elements is the Sun made of?
- 1.3** The Sun rotates around its axis every 27 days. Calculate the ratio of the centrifugal acceleration a over the gravitational acceleration g for a mass element on the surface of the Sun.
- 1.4** The masses of stars are approximately in the range $0.08 M_{\odot} \leq M \leq 100 M_{\odot}$. Why is there an upper limit? Why is there a lower limit?
- 1.5** Can you think of methods to measure (1) the mass, (2) the radius, and (3) the luminosity of a star?
- 1.6** Think of a method to estimate the age of the clusters, discuss with your fellow students. Estimate the ages and compare with the results of your fellow students. Can you give an error range on your age estimates?

Exercises (2)

- 1.7** (a) Calculate the radius of an M5 I supergiant with $\log(L/L_\odot) = 5.50$ and $T_{\text{eff}} = 2700$ K.
(b) Assume a mass of approximately $20M_\odot$ and calculate the mean density of the star.
(c) Calculate the escape velocity.
(d) Compare these values with those for the Sun.
- 1.8** (a) Calculate the luminosities of the horizontal branch stars with $B-V \approx +0.30$ and $BC = 0.11$ in the cluster M3 (NGC 5272). The distance to M3 is 10.4 kpc and its interstellar extinction is negligible.
(b) What is the bolometric correction of the two stars at $B-V \approx 0.25$?
- 1.9** The star τ Sco has an apparent visual magnitude of $V = +2.8$ and a spectral type of approximately B0V. Parallax measurements indicate a distance of 470 ly.
(a) Calculate the absolute visual magnitude.
(b) Adopt the bolometric correction from Table 2.1 and calculate the luminosity L .
(c) Adopt the value of T_{eff} from Table 2.1 and calculate the radius.
(d) Estimate the mass.
(e) Calculate the acceleration of gravity at the stellar surface and the escape velocity.
(f) Calculate the mean density of the star.
(g) Compare these values with those of the Sun.
- 1.10** The Gaia satellite will measure parallaxes with an accuracy of $2 \times 10^{-5}''$ for stars with $V < 15$
(a) What is the distance d of an M5 I supergiant with $\log(L/L_\odot) = 5.50$ and $BC = 3.70$ that has $V = 12$, if the effect of interstellar extinction is ignored?
(b) What is the relative distance accuracy $\sigma(d)/d$ of such a star?
(c) Assuming that T_{eff} is known, what is the relative accuracy in radius $\sigma(R)/R$?

Questions

Q1: Note that the B-V color hardly changes with Teff for temperatures above about 30,000 K. Explain this in physical terms, using the properties of Planck's formula for blackbody radiation.

Q2: Explain in physical terms why the red giant branch curves strongly to high values of B-V for a relatively small change in Teff. (Hint: use the properties of the Planck curve for blackbody radiation)

Q3: Explain why the distribution of stars in the CMD of M3 is completely different from the Milky Way.