


Física Estelar

Lluís Galbany, Ed. Mecenas (#16)

Inma Domínguez, Ed. Mecenas (#16)

Antonio García, Ed. Mecenas (#16)



Curso 2019-2020

5. Energy transport in stellar interiors

Energy Transport

We have seen that in stellar interiors:

LTE is a good approach

Thermal equilibrium

Internal energy sources, nuclear reactions, compensate energy loss from the surface,

$$L = L_{\text{nuc}} = - \frac{dE_{\text{nuc}}}{dt}$$

The total energy is conserved, and the virial theorem states that the E_{int} and E_{pot} are conserved as well,

$$\dot{E}_{\text{tot}} = \dot{E}_{\text{int}} = \dot{E}_{\text{pot}} = 0$$

This stationary state is known as **Thermal Equilibrium (TE)**

Energy is radiated at the surface at the same rate at which it is produced by nuclear reactions in the interior.

Energy Transport

We have seen that in stellar interiors:

LTE is a good approach

Mean free path is extremely small ($\lambda \ll R$)

The time radiation takes to escape from the center of the Sun by the random walk process is the K-H timescale.

Thermal (Kelvin-Helmholtz) timescale, (changes in the thermal structure; *from V. Th.*)

$$\tau_{KH} = \frac{E_{int}}{L} \simeq \frac{|E_{pot}|}{2L} \simeq \frac{GM^2}{2RL} \simeq 1.5 \times 10^7 \left(\frac{M}{M_\odot} \right)^2 \frac{R_\odot}{R} \frac{L_\odot}{L} \text{yr}$$

describes how fast changes in thermal structure of a star can occur

Changes in the Sun luminosity would occur after millions of years, on the timescale for radiative energy transport: K-H timescale for thermal readjustment.

Energy Transport

In this situation, energy can be transported (hot to cold):

DIFFUSION: random thermal motion of particles.

- *Radiative diffusion* in the case of photons
- *Heat conduction* for gas particles

CONVECTION: collective ordered motion of gas particles
Very efficient & rapid mixing

This leads to 2 new equations for stellar structure

Local energy conservation

Virial Th, regulates the global energy budget.
(conservation)

In local scale the internal energy can be changed by two forms,

$$\delta u = \delta q + \frac{P}{\rho^2} \delta \rho$$


— Heat — Work

Changes in the heat content can occur due to:

- Heat is added by the release of nuclear energy (ϵ_{nuc})
- Heat is removed by the release of neutrinos ($\epsilon_{\text{nuc}}, \epsilon_\nu$)
- Heat is absorbed/emitted according to the balance of heat fluxes (*local luminosity*; $l = 4\pi r^2 F$)
where $l(0) = 0$ and $l(R) = L$

Local energy conservation

Consider a spherical shell. The heat content of the shell is:



$$\delta Q = \delta q \Delta m$$

so,

$$\delta Q = \epsilon_{\text{nuc}} \Delta m \delta t - \epsilon_{\nu} \Delta m \delta t + l(m) \delta t - l(m + \Delta m) \delta t$$

$$l(m + \Delta m) \delta t = l(m) + \frac{\partial l}{\partial m} \Delta m$$

rearranging,

$$\delta q = \left(\epsilon_{\text{nuc}} - \epsilon_{\nu} - \frac{\partial l}{\partial m} \right) \delta t$$


And combined with the change of internal energy eq.,

$$\boxed{\frac{\partial l}{\partial m} = \epsilon_{\text{nuc}} - \epsilon_{\nu} - \frac{\partial u}{\partial t} + \frac{P}{\rho^2} \frac{\partial \rho}{\partial t}}$$

LEC eq.

Local energy conservation

Consider a spherical shell. The heat content of the shell is:



$$\delta Q = \delta q \Delta m$$

so,

$$\delta Q = \epsilon_{\text{nuc}} \Delta m \delta t - \epsilon_{\nu} \Delta m \delta t + l(m) \delta t - l(m + \Delta m) \delta t$$

$$l(m + \Delta m) \delta t = l(m) + \frac{\partial l}{\partial m} \Delta m$$

rearranging,

$$\delta q = \left(\epsilon_{\text{nuc}} - \epsilon_{\nu} - \frac{\partial l}{\partial m} \right) \delta t$$

And combined with the change of internal energy eq.,

$$\frac{\partial l}{\partial m} = \epsilon_{\text{nuc}} - \epsilon_{\nu} - \frac{\partial u}{\partial t} + \frac{P}{\rho^2} \frac{\partial \rho}{\partial t}$$

LEC eq.

$$\Rightarrow = -T \frac{\partial s}{\partial t} = \epsilon_{\text{gr}}$$

Local energy conservation

$$\frac{\partial l}{\partial m} = \epsilon_{\text{nuc}} - \epsilon_\nu + \epsilon_{\text{gr}}$$

$\epsilon_{\text{gr}} > 0$ Energy released (contraction)

$\epsilon_{\text{gr}} < 0$ Energy absorbed (expansion)

Thermal equilibrium achieved when $\epsilon_{\text{gr}} = 0$, so

$$\frac{\partial l}{\partial m} = \epsilon_{\text{nuc}} - \epsilon_\nu$$

Andd integratinng over the mass (Lagr. coord.)

$$L = \int_0^M \epsilon_{\text{nuc}} dm - \int_0^M \epsilon_\nu dm \equiv L_{\text{nuc}} - L_\nu$$

Neglecting $L_\nu \rightarrow$

Thermal equilibrium

Internal energy sources, nuclear reactions, compensate energy loss from the surface,

$$L = L_{\text{nuc}} = -\frac{dE_{\text{nuc}}}{dt}$$

The total energy is conserved, and the viral theorem states that the E_{int} and E_{tot} are conserved as well,

$$\dot{E}_{\text{tot}} = \dot{E}_{\text{int}} = \dot{E}_{\text{pot}} = 0$$

This stationary state is known as **Thermal Equilibrium (TE)**

Energy is radiated at the surface at the same rate at which it is produced by nuclear reactions in the interior.

Diffusion

When gradient of energy, diffusion is given by Fick's diffusion law:

$$F = -D \nabla U \quad \text{where}$$

$$D = \frac{1}{3} \bar{v} l \quad \text{Diffusion coefficient}$$

$$\nabla U = C_V \nabla T$$

$$F = -\frac{1}{3} \bar{v} l C_V \nabla T$$

For photons, $\bar{v} = c$

$$U = aT^4 \quad C_V = \frac{dU}{dT} = 4aT^3$$

And l can be taken from
the eq. of radiative transfer

$$\frac{dI_\nu}{ds} = -\overline{\kappa}_\nu \rho I_\nu \xrightarrow{\text{Opacity}} l_{\text{ph}} = \frac{1}{\kappa_\nu \rho}$$

$$\text{so, } F_{\text{rad}} = -\frac{4}{3} \frac{acT^3}{\kappa\rho} \nabla T \sim \frac{l}{4\pi r^2}$$

in spherical Symmetry

Radiative conductivity

Radiative diffusion (photons)

rearranging,
$$\frac{\partial T}{\partial r} = - \frac{3\kappa\rho}{16\pi acT^3} \frac{1}{r^2}$$

or combined with mass continuity eq.
$$\frac{\partial T}{\partial m} = - \frac{3}{64\pi^2 ac} \frac{\kappa l}{r^4 T^3}$$

which describes the T gradient in stellar interiors when E is transported by radiation

Only valid when $l_{\text{ph}} \ll R$, when LTE holds (e.g. in stellar surface $l_{\text{ph}} \geq R$)

In HE, we can combine this with eq. Motion
$$\frac{dP}{dm} = - \frac{Gm}{4\pi r^4}$$

$$\frac{dT}{dm} = \frac{dP}{dm} \frac{dT}{dP} = - \frac{Gm}{4\pi r^4} \frac{T}{P} \frac{d \log T}{d \log P}$$

so then, we can define the radiative T gradient

$$\nabla_{\text{rad}} = \left(\frac{d \log T}{d \log P} \right)_{\text{rad}} = \frac{3}{16\pi acG} \frac{\kappa l P}{m T^4}$$

Roseland mean opacity

So far, we have assumed radiative diffusion is independent of frequency
However, in practice, opacity depends on frequency

$$F_\nu = -D_\nu \nabla U_\nu = -D_\nu \frac{\partial U_\nu}{\partial T} \nabla T$$

where $D_\nu = \frac{1}{3} c l_\nu = \frac{c}{3\kappa_\nu \rho}$ and $U_\nu = \frac{8\pi h}{c^3} \frac{\nu^3}{e^{\frac{h\nu}{kT}} - 1}$

Planck function for BB radiation (Ch. 3)

the total flux is integrated over all frequencies

$$F = - \left[\frac{c}{3\rho} \int_0^\infty \frac{1}{\kappa_\nu} \frac{\partial U}{\partial T} d\nu \right] \nabla T = - [K_{\text{rad}}] \nabla T$$

From both expressions of conductivity,

$$\frac{1}{\kappa} = \frac{1}{4aT^3} \int_0^\infty \frac{1}{\kappa_\nu} \frac{\partial U}{\partial T} d\nu$$

is the Rosseland mean opacity
(the average transparency of the star)

Heat conduction (gas particles)

Collisions between gas particles (e- or ions) can also transport heat

However, their cross sections are typically $\sim 10^{-19} \text{ cm}^2$ and $l_{\text{gas}} \ll l_{\text{ph}}$

Also $\bar{v} \ll c$, so in general this contribution can be ignored

Only important when particles (e.g. e-) are degenerate

- v increase (momenta approach the Fermi momentum)
- Mean free path $l_e \gg l_{\text{ph}}$, and e- conduction becomes more efficient

Similarly, $F_{cd} = -K_{cd}\nabla T = -\frac{4acT^3}{3\kappa_{cd}\rho}\nabla T$

And in general,

$$F = F_{\text{rad}} + F_{cd} = -(K_{\text{rad}} + K_{cd})\nabla T = -\frac{4acT^3}{3\kappa\rho}\nabla T$$

where $\frac{1}{\kappa} = \frac{1}{\kappa_{\text{rad}}} + \frac{1}{\kappa_{cd}}$

LOONEY TUNES



That's all folks.