





Studying Core Collapsed SNe environment with CALIFA



Lluís Galbany, CENTRA-IST Lisboa
Vallery Stanishev, CENTRA-IST Lisboa
Ana Mourão, CENTRA-IST Lisboa
Myriam Rodrigues, ESO Chile
Hector Flores, Observatoire de Paris



Studying Core Collapsed **and Type Ia** SNe environment with CALIFA



Lluís Galbany, CENTRA-IST Lisboa
Vallery Stanishev, CENTRA-IST Lisboa
Ana Mourão, CENTRA-IST Lisboa
Myriam Rodrigues, ESO Chile
Hector Flores, Observatoire de Paris



Motivation

CCSNe

SNela

Stellar evolution
(progenitors)

Anderson, Modjaz, Prieto, Leloudas,
Smartt, Soderberg, Follatelli, Pignata...

overall galaxy properties
slits o imaging - local properties

CALIFA - IFS

Cosmology

Improve standardization
adding host/environment properties
Reduce systematic errors

Sullivan, Kelly, Howell, Lampeitl,
Hamuy, D'Andrea, Gallagher...

most based on global
characteristics (or gradients)

previous study (IFS)
Stanishev+12 (7 SNe)

+

CALIFA - IFS

properties of one single pixel

Sample selection

- Cross-Check for SNe in both NED database (by name) and IAU list (by coordinates)

266 galaxies observed (at least with one grating)

56 hosted **67** SNe (9 with 2 SNe, 1 with 3 SNe) (**35** COMB, **21** only V500)

44 in the field of view: **12** SNIa, **9** SNIbc, **17** SNII (2 b, 1 P, 4 n), **6** untyped

5 SNIa hosts with no emission lines:



NGC4784

UGC10097

NGC6146

NGC6166

NGC6173

Sample selection


- Cross-Check for SNe in both NED database (by name) and IAU list (by coordinates)

266 galaxies observed (at least with one grating)

56 hosted **67** SNe (9 with 2 SNe, 1 with 3 SNe) (**35** COMB, **21** only V500)

44 in the field of view: **12** SNIa, **9** SNIbc, **17** SNII (2 b, 1 P, 4 n), **6** untyped

5 SNIa hosts with no emission lines:



NGC4784

UGC10097

NGC6146

NGC6166

NGC6173

Sample selection

- Cross-Check for SNe in both NED database (by name) and IAU list (by coordinates)


266 galaxies observed (at least with one grating)

56 hosted 67 SNe (9 with 2 SNe, 1 with 3 SNe) (35 COMB, 21 only V500)


44 in the field of view: 12 SNIa, 9 SNIbc, 17 SNII (2 b, 1 P, 4 n), 6 untyped

5 SNIa hosts with no emission lines: Sample, 33 SNe: 17 II, 9 Ibc, 7 Ia

- Search for images with the SN (visually check)




- Fit galaxy center (Σ_{rows} and Σ_{columns})



- Determine SN position from offset (IAU, literature...)

Reduction

- Use Rubén (et al.) **masks** when available
- **Resample** to 1Å, **remove** sky lines and **shift** to the rest-frame
- **Fit** all spectra with STARLIGHT




CB07: 17 Ages 10^6 to $1.8 \cdot 10^{10} M_{\odot}$
 4 metallicities 0.004, 0.05, 0.2, $2.5 Z_{\odot}$ ($Z_{\odot}=0.02$)
 S/N window: [4580:4640] Å
 Fit range: [3660:7000] Å

MILES stellar library
 + Padova (1994) evol. tracks
 + Chabrier (2003) IMF
 + Marigo (2007)

- **Subtract** continuum and **fit** emission lines with MPFIT


[O II] $\lambda 3727$	[Ne I] $\lambda 4931$	[O I] $\lambda 6364$
[Ne III] $\lambda 3868$	[O III] $\lambda 4959$	[N II] $\lambda 6548$
H ϵ $\lambda 3970$	[O III] $\lambda 5007$	H α $\lambda 6563$
H δ $\lambda 4102$	[O I] $\lambda 5577$	[N II] $\lambda 6583$
H γ $\lambda 4340$	[He I] $\lambda 5875$	[S II] $\lambda 6716$
[O III] $\lambda 4363$	[O I] $\lambda 6300$	[S II] $\lambda 6731$
H β $\lambda 4861$	[S III] $\lambda 6312$	

H α velocity to shift whole spectrum
 Balmer decrement to correct for extinction $\rightarrow E(B-V)$
 Line flux, line velocity, gaussian fit parameters
 Produce 2D maps for each line




Reduction

- Use Rubén (et al.) **masks** when available
- **Resample** to 1Å, **remove** sky lines and **shift** to the rest-frame
- **Fit** all spectra with STARLIGHT



CB07: 17 Ages 10^6 to $1.8 \cdot 10^{10} M_{\odot}$
 4 metallicities 0.004, 0.05, 0.2, $2.5 Z_{\odot}$ ($Z_{\odot}=0.02$)
 S/N window: [4580:4640] Å
 Fit range: [3660:7000] Å



- **Subtract** continuum and **fit** emission lines with MPFIT


[O II] $\lambda 3727$	[Ne I] $\lambda 4931$	[O I] $\lambda 6364$
[Ne III] $\lambda 3868$	[O III] $\lambda 4959$	[N II] $\lambda 6548$
H ϵ $\lambda 3970$	[O III] $\lambda 5007$	H α $\lambda 6563$
H δ $\lambda 4102$	[O I] $\lambda 5577$	[N II] $\lambda 6583$
H γ $\lambda 4340$	[He I] $\lambda 5875$	[S II] $\lambda 6716$
[O III] $\lambda 4363$	[O I] $\lambda 6300$	[S II] $\lambda 6731$
H β $\lambda 4861$	[S III] $\lambda 6312$	

H α velocity to shift whole spectrum

Balmer decrement to correct for extinction $\rightarrow E(B-V)$

Line flux, line velocity, gaussian fit parameters

Produce 2D maps for each line



Measurements

- Electronic density ($N_e = \text{SIIa/SIIb}$)
- Diagnostic diagram (BPT)
- Several metallicity calibrations

Pettini & Pagel 04 (PP04): [N2](#), [O3N2](#)

Perez-Montero & Contini 09 (PC09): [O3N2_NO](#)



Kobulnicky & Kewley 04 (KK04): [R23](#)

Tremonti+04 (T04): [R23_T](#)

Pilyugin+10: [P10](#)

Pilyugin+11: [P11](#)

- Ionization parameter
- Star formation rate (sqarcsec, sqkpc, Σ)
- Stellar populations parameters from STARLIGHT



Kinemetry

Deprojection

Azimuthal average

Voronoi binning

Integrated spectrum

3" aperture spectrum

Kinematics

Fit ellipses using Krajnovic et al. 2006



Deprojection

Azimuthal average

Voronoi binning

Integrated spectrum

3" aperture spectrum



Kinematics

Deprojection

Fit ellipses using Krajnovic et al. 2006



↪ Measure distances in the galactic plane

Azimuthal average

Voronoi binning

Integrated spectrum

3" aperture spectrum



Kinematics

Fit ellipses using Krajnovic et al. 2006

Deprojection

Measure distances in the galactic plane



Azimuthal average

Co-add ell. rings centered in the core

Voronoi binning

Integrated spectrum

3" aperture spectrum



Kinematics

Fit ellipses using Krajnovic et al. 2006

Deprojection

Measure distances in the galactic plane

Azimuthal average



Co-add ell. rings centered in the core

Voronoi binning

Requiring $S/N > 20$

Integrated spectrum

3" aperture spectrum



Kinematics

Fit ellipses using Krajnovic et al. 2006

Deprojection

Measure distances in the galactic plane

Azimuthal average

Co-add ell. rings centered in the core


Voronoi binning

Requiring $S/N > 20$

Integrated spectrum

For high- z (aperture effects)

3" aperture spectrum



Kinematics

Fit ellipses using Krajnovic et al. 2006

Deprojection

Measure distances in the galactic plane

Azimuthal average

Co-add ell. rings centered in the core

Voronoi binning


Requiring $S/N > 20$

Integrated spectrum

For high- z (aperture effects)

3" aperture spectrum


Allow comparisons (SDSS + fiber)



Results...


$$Z_{\text{Ia}} > Z_{\text{Ibc}} > Z_{\text{II}}$$

long slit @SN




Anderson+10

$$Z_{\text{Ia}} > Z_{\text{Ibc}} > Z_{\text{II}}$$






long slit @SN




Anderson+10


$$Z_{\text{Ia}} > Z_{\text{Ibc}} > Z_{\text{II}}$$



H α emission (SF)




H α imaging




Anderson+12


NCR result



H α emission (SF)




H α imaging




Anderson+12


NCR result



H α emission (SF)




H α imaging




Anderson+12


NCR result




H α emission (SF)




@SN/MAX






@SN/TOTAL



H α emission (SF)




HIexplorer



global Z Vs Local Z


long slit @SN






II -0.034
Ibc -0.011
Ia 0.111

II -0.034
Ibc -0.009
Ia 0.107

II -0.013
Ibc 0.004
Ia 0.027




Modjaz+11




global Z Vs Local Z

long slit @SN



Modjaz+11



Conclusions and TO DOs

- We found some indications on differences between type II, Ibc and Ia SNe
 - mass, age, metallicity
 - progenitors
- we found differences between global and local metallicities
 - ...cosmology (aperture, gradients...)
- Need of more statistics to make strong conclusions
- Need to correlate other parameters
- Suggestions are welcome

Conclusions and TO DOs

- We found some indications on differences between type II, Ibc and Ia SNe
 - mass, age, metallicity
 - progenitors
- we found differences between global and local metallicities
 - ...cosmology (aperture, gradients...)
- Need of more statistics to make strong conclusions
- Need to correlate other parameters
- Suggestions are welcome

Obrigacias!

more...

