



IFU observations of SN environments



Lluís Galbany, CENTRA-IST Lisboa
Vlery Stanishev, CENTRA-IST Lisboa
Ana Mourão, CENTRA-IST Lisboa
Myriam Rodrigues, ESO, Chile
Hector Flores, Obs. Paris, France



~~Conclusions~~ Summary

- We found some indications on differences between type II, Ibc and Ia SNe
 - GCD
 - Star formation
 - Metallicity
- We found differences between global and local measurements
- work in progress... (gradients, any other idea?)

Astronomy & Astrophysics manuscript no. ccsne
April 10, 2013

© ESO 2013

Integral Field Unit spectroscopy of supernova host galaxies

Lluís Galbany¹, Vallery Stanishev¹, Ana M. Mourão¹, Myriam Rodrigues², Hector Flores³, Luke Skywalker⁴, Bilbo Baggins⁵, Robb Stark⁶, and other heroes.

¹ CENTRA - Centro Multidisciplinar de Astrofísica, Instituto Superior Técnico, Av. Rovisco Pais 1, 1049-001 Lisbon, Portugal.

² European Southern Observatory, Alonso de Cordova 3107 Casilla 19001 - Vitacura -Santiago, Chile

³ GEPI, Observatoire de Paris, UMR 8111, CNRS, Université Paris Diderot, 5 place Jules Janssen, 92190 Meudon, France

⁴ IRF - Institute of Research on the Force, Jedi University, 08187 Galactic City, Crousant.

⁵ Center for Research in Astronomy (CRA), University of the Shire, 08402 Hobbiton, Eriador.

⁶ Institute of Advanced Sciences, University of the North, 17007 Winterfell, Westeros.

Received ——— / Accepted ———

Abstract

We aim to study the properties of the environment where supernovae (SNe) exploded, using Integral Field Unit (IFU) spectroscopic information of nearby galaxies ($0.005 < z < 0.03$) provided by the Calar Alto Legacy Integral Field Area (CALIFA) Survey and previous observations. With this approach we are going further than using simply an aperture spectrum centered at the galaxy core, or a spectrum from a slit positioned at a SN explosion, since we have 2-dimensional spectroscopic information of the whole field of each host galaxy. We find differences in the location distributions among different SN types, being type Ibc more centered and occurring closer to H-II regions than type Ia, being type II in the middle of those. We also found differences although in the local elemental abundances of type Ibc compared to type II SNe. ...

Key words. supernovae: general – supernovae: host galaxies – techniques: spectroscopic

1. Introduction

tbw... Integral Field Unit (IFU)

IMPORTANCE OF SUPERNOVAE: Supernovae... ANDERSON: A key goal in current SN search is to provide links between progenitor systems and observed transients.

CCSNE: Core-collapsed supernovae are the final stage in the evolution of stars with large initial masses ($> 8 M_{\odot}$). They are thought to arise from massive stars that collapse once they have spent their nuclear fuel. The exact understanding of their progenitor systems, and explosion mechanism remains elusive. HAKOBYAN08: Type II SNe bursts because of the collapse of the cores of young massive stars - the reds supergiants, whose envelopes contain a large amount of hydrogen. type Ibc SNe also bursts during the collapse of the cores of massive stars, probably Wolf-Rayet stars, but a large portion of the hydrogen envelopes of these stars are lost during the evolution of a close binary system or by some other means, which is probably why their progenitors have higher metallicity. TURKU: Core-collapse supernovae (CCSNe) mark the end-point of the evolution of massive stars, producing neutron stars and black holes, and in exceptional cases, long-duration gamma-ray bursts (GRBs). These phenomena play a vital role in our understanding of stellar evolution, the synthesis of heavy elements, and through feedback processes also in galaxy evolution. Furthermore, as CCSNe come from massive short-lived stars, their explosion rate directly reflects the on-

going rate of star formation in their host galaxies, and thus CCSNe are beginning to be used as probes of the cosmic star formation history.

DIFFERENCES IBC/II: VANDERBERG97 showed that Ibc appeared to be more concentrated than II within host galaxies. EXPLANATION OF THE DIFFERENCES: Higher metallicity stars have higher line driven winds (Puls et al 1996, Hadrizti and Puls 2000, Mokiemi et al. 2007), thus the progenitors lose more mass through these winds prior to SN, hence removing their H and He envelopes.

SNEIA: Type-Ia SNe...

...as Anderson et al 2009, Prieto et al. 2008 conclude, metallicity should be directly derived at the SN locations... ..lack of CCSNe and presence of thermonuclear SNIa in early type elliptical galaxies...

GLOBAL VS LOCAL PARAMETERS: GLOBAL PARAMETERS IA: Usually global parameters of the host galaxies have been used to try to correlate host and SN properties... Hamuy et al. (2000), bright Type Ia SNe occur preferentially in young stellar environments (AGE). Hamuy et al. (2000), bright SNe occur in low luminosity hosts. Gallagher et al. (2005), faint SNe occur in metal-rich hosts (METALLICITY). Sullivan et al. (2010); Kelly et al. (2010); Lampeitl et al. (2010), bright SNe in massive hosts (MASS) and in hosts with low SFR per stellar mass (specific SFR), after LC correction.

Stanishev et al. (2012): many studies, bright SNe occur in star-forming hosts, faint in passive hosts. Gallagher et al. (2008) - GLOBAL METALLICITY. Howell et al. (2009) - no metallicity, PROGENITOR AGE. Sullivan et al.

Send offprint requests to: L. Galbany, e-mail: lluís.galbany@ist.utl.pt

Motivation

CCSNe

SN Ia

Stellar evolution
(progenitors)

Anderson, Modjaz, Prieto, Leloudas,
Smartt, Soderberg, Follatelli, Pignata...

overall galaxy properties
slits o imaging - local properties

CALIFA - IFS

Cosmology

Improve standardization
adding host/environment properties
Reduce systematic errors

Sullivan, Kelly, Howell, Lampeitl,
Hamuy, D'Andrea, Gallagher...

most based on global
characteristics (or gradients)


previous study (IFS)
Stanishev+12 (7 SNe)

+

CALIFA - IFS

properties of one single pixel

Second order corrections: Environment



Look for dependences of the SN properties on the host galaxy properties (focused on global characteristics of the host)

As they evolve with redshift, such dependences would impact the cosmological parameters

Hamuy et al. (1996)

Hamuy et al. (2000)

Gallagher et al. (2005)

Sullivan et al. (2006)

Gallagher et al. (2008)

Hicken et al. (2009)

Howell et al. (2009)

Neil et al. (2009)

Brandt et al. (2010)

Cooper et al. (2010)

Sullivan et al. (2010)

Kelly et al. (2010)

Lampeitl et al. (2010)

D'Andrea et al. (2011)

Gupta et al. (2011)

Nordin et al. (2011)

Konishi et al. (2011)

Smith et al. (2012)

...

Bright events occur preferentially in **young** stellar environments.

Luminous SNe are produced in **metal-poor** neighborhoods

Age is more likely to be the source of LC variability than **metallicity**

Brighter events are found in systems with ongoing **star-formation**

Progenitor age primarily determines the peak luminosity

SN Ia in **spiral** hosts are intrinsically fainter (*after LC-corr*)

more massive progenitors give rise to less luminous explosions

Older hosts produce less-extincted SNe Ia

Luminous SNe associated with recent **star-formation** and **young** prog.

SN Ia are more luminous or more numerous in **metal-poor** galaxies

SN Ia are brighter in **massive** hosts (metal-rich) and with low **SFR** (*after LC-corr*)

SN Ia in physically **larger**, more **massive** hosts are ~10% brighter

introduce the stellar **mass** of the host in the parametrization


SNe are 0.1 mag brighter in **high-metallicity** hosts after corr.

older galaxies host SNe Ia that are brighter

passive and **massive** galaxies host faint SNe

SNe in **metal-rich** hosts become brighter after corrections

SNe rate is higher in **star-forming** galaxies



Anderson+12

Sample selection

- Cross-Check SNe IAU list with CALIFA galaxies (by coord.)

~300 galaxies observed (at least with one grating)

55 hosted 65 SNe (8 with 2 SNe, 1 with 3 SNe)

47 in the field of view: 13 SNIa, 9 SNIbc, 19 SNII (2b, 2P, 5n), 6 untyped

4 SNIa hosts with no emission lines:

37 SNe: 19 II, 9 Ibc, 9 Ia

+ 23 SNe: 8 II, 4 Ibc, 11 Ia

60 SNe: 27 II, 13 Ibc, 20 Ia


- Previous observations



feasibility study for CALIFA, *Sánchez+12*

PINGS Survey, *Rosales-Ortega+10*

SNIa hosts, *Stanishev+12*

NGC5668, *Marino+12*





Kinometry

Deprojection

Azimuthal average

Voronoi binning

Integrated spectrum

3" aperture spectrum

Kinematics

Fit ellipses using *Krajnovic et al. 2006*



Deprojection

Azimuthal average

Voronoi binning

Integrated spectrum

3" aperture spectrum



Kinematics

Deprojection

Fit ellipses using *Krajnovic et al. 2006*



↳ Measure distances in the galactic plane

Azimuthal average

Voronoi binning

Integrated spectrum

3" aperture spectrum



Kinematics

Fit ellipses using *Krajnovic et al. 2006*

Deprojection

Measure distances in the galactic plane



Azimuthal average

Co-add ell. rings centered in the core

Voronoi binning

Integrated spectrum

3" aperture spectrum



Kinematics

Fit ellipses using *Krajnovic et al. 2006*

Deprojection

Measure distances in the galactic plane

Azimuthal average



Co-add ell. rings centered in the core

Voronoi binning

Requiring $S/N > 20$

Integrated spectrum

3" aperture spectrum



Kinematics

Fit ellipses using *Krajnovic et al. 2006*

Deprojection

Measure distances in the galactic plane

Azimuthal average

Co-add ell. rings centered in the core


Voronoi binning

Requiring $S/N > 20$

Integrated spectrum

For high- z (aperture effects)

3" aperture spectrum



Kinematics

Fit ellipses using *Krajnovic et al. 2006*

Deprojection

Measure distances in the galactic plane

Azimuthal average

Co-add ell. rings centered in the core

Voronoi binning


Requiring $S/N > 20$

Integrated spectrum

For high- z (aperture effects)


3" aperture spectrum

Allow comparisons (SDSS + fiber)




Results...

GCDs






GCDs





HII regions

HIIexplorer
Sánchez et al. 2012




Σ_{SFR}

Stellar age





$$Z_{\text{Ia}} \geq Z_{\text{Ibc}} > Z_{\text{II}}$$




Broad-band photometry Fiber slit spectroscopy


$z=0.016$




$z=0.25$



global Z Vs Local Z



metallicity gradients



Conclusions

- We found some indications on differences between type II, Ibc and Ia SNe
 - GCD
 - Star formation
 - Metallicity
- We found differences between global and local measurements
- work in progress... (gradients, any other idea?)

Integral Field Unit spectroscopy of supernova host galaxies

Lluís Galbany¹, Vallery Stanishev¹, Ana M. Mourão¹, Myriam Rodrigues², Hector Flores³, Luke Skywalker⁴, Bilbo Baggins⁵, Robb Stark⁶, and other heroes.

¹ CENTRA - Centro Multidisciplinar de Astrofísica, Instituto Superior Técnico, Av. Rovisco Pais 1, 1049-001 Lisbon, Portugal.

² European Southern Observatory, Alonso de Cordova 3107 Casilla 19001 - Vitacura -Santiago, Chile

³ GEPI, Observatoire de Paris, UMR 8111, CNRS, Université Paris Diderot, 5 place Jules Janssen, 92190 Meudon, France

⁴ IRF - Institute of Research on the Force, Jedi University, 08187 Galactic City, Crousant.

⁵ Center for Research in Astronomy (CRA), University of the Shire, 08402 Hobbiton, Eriador.

⁶ Institute of Advanced Sciences, University of the North, 17007 Winterfell, Westeros.

Received ——— / Accepted ———

Abstract

We aim to study the properties of the environment where supernovae (SNe) exploded, using Integral Field Unit (IFU) spectroscopic information of nearby galaxies ($0.005 < z < 0.03$) provided by the Calar Alto Legacy Integral Field Area (CALIFA) Survey and previous observations. With this approach we are going further than using simply an aperture spectrum centered at the galaxy core, or a spectrum from a slit positioned at a SN explosion, since we have 2-dimensional spectroscopic information of the whole field of each host galaxy. We find differences in the location distributions among different SN types, being type Ibc more centered and occurring closer to H-II regions than type Ia, being type II in the middle of those. We also found differences although in the local elemental abundances of type Ibc compared to type II SNe. ...

Key words. supernovae: general – supernovae: host galaxies – techniques: spectroscopic

1. Introduction

tbw... Integral Field Unit (IFU)

IMPORTANCE OF SUPERNOVAE: Supernovae... ANDERSON: A key goal in current SN search is to provide links between progenitor systems and observed transients.

CCSNE: Core-collapsed supernovae are the final stage in the evolution of stars with large initial masses ($> 8 M_{\odot}$). They are thought to arise from massive stars that collapse once they have spent their nuclear fuel. The exact understanding of their progenitor systems, and explosion mechanism remains elusive. HAKOBYAN08: Type II SNe bursts because of the collapse of the cores of young massive stars - the reds supergiants, whose envelopes contain a large amount of hydrogen. type Ibc SNe also bursts during the collapse of the cores of massive stars, probably Wolf-Rayet stars, but a large portion of the hydrogen envelopes of these stars are lost during the evolution of a close binary system or by some other means, which is probably why their progenitors have higher metallicity. TURKU: Core-collapse supernovae (CCSNe) mark the end-point of the evolution of massive stars, producing neutron stars and black holes, and in exceptional cases, long-duration gamma-ray bursts (GRBs). These phenomena play a vital role in our understanding of stellar evolution, the synthesis of heavy elements, and through feedback processes also in galaxy evolution. Furthermore, as CCSNe come from massive short-lived stars, their explosion rate directly reflects the on-

going rate of star formation in their host galaxies, and thus CCSNe are beginning to be used as probes of the cosmic star formation history.

DIFFERENCES IBC/II: VANDERBERG97 showed that Ibc appeared to be more concentrated than II within host galaxies. EXPLANATION OF THE DIFFERENCES: Higher metallicity stars have higher line driven winds (Puls et al 1996, Hadrizti and Puls 2000, Mokiemi et al. 2007), thus the progenitors lose more mass through these winds prior to SN, hence removing their H and He envelopes.

SNEIA: Type-Ia Sne...

...as Anderson et al 2009, Prieto et al. 2008 conclude, metallicity should be directly derived at the SN locations... ..lack of CCSNe and presence of thermonuclear SNIa in early type elliptical galaxies...

GLOBAL VS LOCAL PARAMETERS: GLOBAL PARAMETERS IA: Usually global parameters of the host galaxies have been used to try to correlate host and SN properties... Hamuy et al. (2000), bright Type Ia SNe occur preferentially in young stellar environments (AGE). Hamuy et al. (2000), bright SNe occur in low luminosity hosts. Gallagher et al. (2005), faint SNe occur in metal-rich hosts (METALLICITY). Sullivan et al. (2010); Kelly et al. (2010); Lampeitl et al. (2010), bright SNe in massive hosts (MASS) and in hosts with low SFR per stellar mass (specific SFR), after LC correction.

Stanishev et al. (2012): many studies, bright SNe occur in star-forming hosts, faint in passive hosts. Gallagher et al. (2008) - GLOBAL METALLICITY. Howell et al. (2009) - no metallicity, PROGENITOR AGE. Sullivan et al.