



Integral Field Spectroscopy of nearby supernova host galaxies




Lluís Galbany, CENTRA-IST
Vallery Stanishev, CENTRA-IST
Ana M. Mourão, CENTRA-IST
Myriam Rodrigues, ESO
Hector Flores, Obs. Paris



CALIFA Survey

Motivation




Core collapse SN

Massive stars (8 to 30 M_{sun})

Differences depending on progenitor
mass loss before explosion

Few (CCSNe) and no (SNe Ia) direct progenitor detection
(e.g. Smartt+09)

Alternative methods to constrain progenitor properties: **ENVIRONMENT**



Type Ia SN

CO White dwarfs in binary systems
accreting mass from a companion

Homogeneous brightness
---> Cosmology

SNe Ia cosmology

Several works have been looking for correlations between the Hubble residuals (HR) and global properties of the host galaxy:

Hamuy et al. (1996)

Hamuy et al. (2000)

Gallagher et al. (2005)

Sullivan et al. (2006)

Gallagher et al. (2008)

Hicken et al. (2009)

Howell et al. (2009)

Neil et al. (2009)

Brandt et al. (2010)

Cooper et al. (2010)

Sullivan et al. (2010)

Kelly et al. (2010)

Lampeitl et al. (2010)

D'Andrea et al. (2011)

Gupta et al. (2011)

Nordin et al. (2011)

Konishi et al. (2011)

Smith et al. (2012)

...

Bright events occur preferentially in **young** stellar environments.

Luminous SNe are produced in **metal-poor** neighborhoods

Age is more likely to be the source of LC variability than **metallicity**

Brighter events are found in systems with ongoing **star-formation**

Progenitor age primarily determines the peak luminosity

SN Ia in **spiral** hosts are intrinsically fainter *(after LC-corr)*

more massive progenitors give rise to less luminous explosions

Older hosts produce less-extincted SNe Ia

Luminous SNe associated with recent **star-formation** and **young** prog.

SN Ia are more luminous or more numerous in **metal-poor** galaxies

SN Ia are brighter in **massive** hosts (metal-rich) and with low **SFR** *(after LC-corr)*

SN Ia in physically **larger**, more **massive** hosts are ~10% brighter

introduce the stellar **mass** of the host in the parametrization

SNe are 0.1 mag brighter in **high-metallicity** hosts after corr.


older galaxies host SNe Ia that are brighter

passive and **massive** galaxies host faint SNe

SNe in **metal-rich** hosts become brighter after corrections

SNe rate is higher in **star-forming** galaxies

Progenitor constraints




Cumulative distributions of local star-formation at SN position for several SN types

More correlation to the Star-formation, means less time to migrate from the formation region (age), and to lose enveloping layers (mass)


Anderson+12

Environmental studies



- Global properties
 - Photometry/imaging (Sullivan+10, Lampeitl+10, Anderson+09, ...)
 - Single-aperture / long-slit spectroscopy (at host galaxy core) (Prieto+08, D'Andrea+12, ...)
- Local properties
 - Global values + gradients (Boissier+09, Galbany+12, ...)
 - Single-aperture / long-slit spectroscopy (at SN position) (Anderson+10&12, Modjaz+11, ...)
 - Integral field Spectroscopy (Stanishev+12, Kuncarayakti+13, ...)

Calar Alto Legacy Integral Field Area



CALIFA Survey

Sánchez+12


- Survey of ~600 galaxies of all types at $z=0.005$ to 0.03
- diameter selected from SDSSDR7, $45 < D_{25} < 80$, to fit in the IFU FOV
CALIFA mother sample: 939 galaxies
- IFS using PPAK @ 3.5m CAHA
2 setups: mid (V500) and high-res (V1200)
Spectral coverage [3700-7000 Å]
Spatial resolution ~1 arcsec
- 250 dark nights over 3 years
- ~3000 spectra per galaxy
- Data will freely distributed to the community.

DRI (100 galaxies), Huseman+13

Sample selection



- Cross-check SNe IAU list with CALIFA galaxies (by coord.)
 - ~350 galaxies observed so far
 - 42 hosted 50 SNe (after careful inspection)
- + previous observations (SAME instrument!):
 - Feasibility Study for CALIFA, Sanchez+12
 - PINGS Survey, Rosales-Ortega+10
 - SNe Ia hosts, Stanishev+12
 - NGC5668, Marino+12
 - Interacting galaxies project, Barrera-Ballesteros in prep.

75 SNe: 32 type II, 13 type Ibc, 30 type Ia



STARLIGHT *Cid Fernandes et al. 2005*

CB07: 17 Ages 10^6 to $1.8 \cdot 10^{10} M_{\odot}$
 4 metallicities 0.004, 0.05, 0.2, $2.5 Z_{\odot}$



Kinematics

Fit ellipses using *Krajnovic et al. 2006*



Deprojection

Azimuthal average

Voronoi binning

Integrated spectrum

3" aperture spectrum



Kinematics

Deprojection

Fit ellipses using *Krajnovic et al. 2006*



↳ Measure distances in the galactic plane

Azimuthal average

Voronoi binning

Integrated spectrum

3" aperture spectrum



Kinematics

Fit ellipses using *Krajnovic et al. 2006*

Deprojection

Measure distances in the galactic plane



Azimuthal average

Co-add ell. rings centered in the core


Voronoi binning

Integrated spectrum

3" aperture spectrum




Star-forming regions




- HIIexplorer (Sanchez+12) to select HII clumps from Ha emission maps
- Measure distances from the SN explosion site to the center of the nearest HII clump

Star-forming regions




- HIIexplorer (Sanchez+12) to select HII clumps from Ha emission maps
- Measure distances from the SN explosion site to the center of the nearest HII clump


HII clump
distance




SF density




$\langle \log t^* \rangle$




H α EW




HII clump
distance




SF density



$\langle \log t^* \rangle$




H α EW




Sequence on association to star-forming regions: from SNe Ibc to SNe Ia

Metallicity




From gas emission line ratios




From stellar population

Metallicity



From gas emission line ratios




From stellar population




Sequence in metallicity at SN position from **Sne Ia** to **Sne II**

Slight difference between mean values of **Ibc** and **II** (~ 0.01 dex)
larger differences in the median (~ 0.04 dex)

Aperture effects

4 different measurements:







- CEN - LOC ~ 0.07 dex CC SNe
- INT - LOC ~ -0.08 dex SNe Ia
- INT(AGN) $\sim$ INT(noAGN)
- GRAD - LOC ~ 0.01 dex CC SNe
 ~ -0.03 dex SNe Ia
 error when using gradients


Metallicity gradients

- Measure metallicity gradients normalizing distance to the disc effective radius (r_e)



- Show a universal gradient (~ -0.1 dex)
- study in preparation using
all CALIFA galaxies
(Sanchez+ in prep.)

Metallicity gradients



- Type Ia SNe do not show a decrease in metallicity at larger distances
- CC SNe local metallicity have lower values in the outskirts

Conclusions

- Differences found in the environment of different SN types
 - Association to star-formation
 - Local metallicity

that can help constrain the properties of each type of SN progenitor

- IFS allowed us to study aperture effects (spectroscopy at different redshift) and the use of metallicity gradients as an indirect approximation to the local values.
 - Central and integrated spectra
 - Integrated spectra including AGN

differences found depending on SN type: usefulness in host galaxy studies in Type Ia cosmology