


INTEGRAL FIELD SPECTROSCOPY OF SUPERNOVA HOST GALAXIES



CALIFA Survey

Lluís Galbany
(CENTRA, Portugal and MAS/DAS, Chile)

Vallery Stanishev (CENTRA, Portugal)

Ana Mourão (CENTRA, Portugal)

Myriam Rodrigues (ESO, Chile)


Hector Flores (GEPI, France)

and the *CALIFA* collaboration




Institut d'Estudis Espacials de Catalunya (IEEC)
21 Maig 2014, Bellaterra

SN zoo



D. Kasen

CO white dwarfs in binary systems accreting mass from a companion




Two scenarios: - single degenerate (WD + massive star)
- double degenerate (2 WD)

Homogeneous brightness → Cosmology

Massive stars (8 to 30 Msun)

Two paths: - single massive stars + winds
- binary systems transferring mass



Differences depending on progenitor mass loss before explosion


Mass loss ↑

- Ic lose both H and He envelopes
- Ib lose H envelope
- IIb intermediate between II and Ib
- II retain external layers (H and He)

Few CCSNe and no SNe Ia direct progenitor detection (e.g. Smartt+09)

Alternative methods to constrain progenitor properties: ENVIRONMENT

Progenitor constraints



Cumulative distributions of local star-formation at SN position for several SN types from H α imaging
Anderson+12

Assuming that all the stars in a cluster have similar ages, more association to the star-forming region can be understood as young and more massive star


More **massive** stars have shorter lifetimes (**age**) and therefore have less time to move away from the star-forming regions

SN Environmental studies


- Global properties
 - photometry / imaging
(Sullivan+10, Lampeitl+10, Anderson+09...)
 - single-aperture / long-slit spectroscopy (@galaxy core)
(Prieto+08, D'Andrea+12...)
- Local properties
 - central values + gradients
(Boissier+09, Galbany+12...)
 - single-aperture / long-slit spectroscopy (@SN position)
(Anderson+10&12, Modjaz+11...)
- Integral Field Spectroscopy
(Stanishev+12, Kuncarayakti+13ab)



Why nearby? The FoV issue




SNe Ia host galaxies with IFU



Stanishev+12

Calar Alto Legacy Integral Field Area

Sánchez+12



CALIFA Survey

- Survey of ~ 600 galaxies of all types at $z=0.005$ to 0.03
- diameter selected from SDSS DR7, $45 < D_{25} < 80$, to fit in the IFU FOV
CALIFA mother sample: 939 galaxies
- IFS using PPAK @ 3.5m CAHA
2 setups: mid (V500) and high-res (V1200)
Spectral coverage [3700-7000 Å]
Spatial resolution ~ 1 arcsec
- 250 dark nights over 3 years
- ~ 3000 spectra per galaxy
- Data will freely distributed to the community.

DR1 (100 galaxies), Husemann+13

DR2 (300 galaxies), García-Benito in prep.

Sample selection

- Cross-Check SNe IAU list with CALIFA galaxies (by coord.)

~400 galaxies observed so far (at least with one grating)

65 hosted 73 SNe (6 with 2 SNe, 1 with 3 SNe)

58 SNe in the field of view: 22 II, 13 Ibc/Ilb, 23 Ia

Previous studies (SAME INSTRUMENT!!)

4 feasibility study for CALIFA, Sánchez+12

8 PINGS Survey, Rosales-Ortega+10

5 SNIa hosts, Stanishev+12


NGC5668, Marino+12

NGC3982, Marino in prep.

12 mergers, Barrera-Ballesteros in prep.

37 SNe: 11 II, 7 Ibc/Ilb, 19 Ia

95 SNe: 33 II, 20 Ibc, 42 Ia





STARLIGHT

Cid Fernandes et al. 2005

CB07: 17 Ages 10^6 to $1.8 \cdot 10^{10} M_{\odot}$

4 metallicities $0.004, 0.05, 0.2, 2.5 Z_{\odot}$



Kinematics




Fit ellipses using [Krajnovic et al. 2006](#)

Deprojection

Measure distances in the galactic plane

Voronoi (spatial) binning

Increase S/N in the outskirts



Kinematics




Fit ellipses using [Krajnovic et al. 2006](#)

Deprojection

Measure distances in the galactic plane

Voronoi (spatial) binning

Increase S/N in the outskirts



GCDs

Distance as a proxy of local metallicity, assuming the presence of gradients


(Galbany+12)

Comparison with ASIAGO SN catalogue

Ibc/I Ib closer to the galaxy core

Ia found further




Ibc/I Ib would explode in metal-richer environments since they explode closer to the center of the galaxy



Global host galaxy properties

Construction of Integrated spectra

- on-going SFR from extinction-corrected H α emission line maps
- galaxy total mass
- specific SFR (sSFR=SFR/mass)






Global host galaxy properties

Construction of Integrated spectra

- on-going SFR from extinction-corrected H α emission line maps
- galaxy total mass
- specific SFR (sSFR=SFR/mass)

Brinchmann+2004






Global host galaxy properties

Construction of Integrated spectra


- on-going SFR from extinction-corrected H α emission line maps
- galaxy total mass
- specific SFR (sSFR=SFR/mass)

Elbaz+2007

Brinchmann+2004



Global host galaxy properties




Most of the stellar mass is locked in old stellar populations

- SFH: percent of stellar populations
- galaxy mean age
- H α equivalent width as a tracer of young stellar population

$$\sim \frac{\text{line}}{\text{cont.}}$$

gas
stars $\leadsto$ mass



50% Ia
30% II
20% Ibc/IIb


-> high-z

Local host galaxy properties

Star-formation rate density
($\Sigma\text{SRF}=\text{SFR}/\text{kpc}^2$)

SFR in the same
neighboring region

Accounting for inclination




Even in galaxies with similar SFR, **Ibc/IIb** tend to explode in higher ΣSRF regions than II and **Ia**

II	-2.05
Ibc/IIb	-1.80
Ia -SF	-2.44
Ia -all	-2.89

Local host galaxy properties

- Stellar age
- Percent of young stellar population
- H α equivalent width




II 8.551 (0.106)
 Ibc/Ilb 8.356 (0.168)
 Ia -SF 8.978 (0.084)
 Ia-all 9.121 (0.091)


II 35.8 (4.1)
 Ibc/Ilb 41.0 (5.8)
 Ia -SF 24.1 (3.8)
 Ia-all 21.1 (3.5)

II 36.7 (5.0)
 Ibc/Ilb 57.3 (14.3)
 Ia -SF 15.2 (2.2)

Local host galaxy properties - NCR




Local host galaxy properties - star forming regions




HII clumps selected from H α maps using HIIexplorer (Sánchez+12)

Measure distances from the SN explosion site to the nearest HII clump


Local host galaxy properties - star forming regions




Local host galaxy properties - oxygen abundance



Disk effective radius (r_e)





Determination of the disk effective radius by fitting the light profile contribution from the disk



Decoupling bulge and disk using the projection of the r-band brightness

Local host galaxy properties - metallicity gradients




Show a characteristic gradient (~ -0.1 dex)

No dependence on O/H estimator or SN type

Local host galaxy properties - local metallicity

- Oxygen abundance from the emission lines (with several indicators)
- Stellar metallicity from the fit to the continuum fit (Z_L , Z_M)




II 8.645 (0.029)
 Ibc/Ilb 8.662 (0.040)
 Ia -SF 8.734 (0.018)

II 0.023 (0.002)
 Ibc/Ilb 0.025 (0.003)
 Ia -SF 0.026 (0.002)
 Ia-all 0.027 (0.002)

II 0.024 (0.003)
 Ibc/Ilb 0.024 (0.004)
 Ia -SF 0.027 (0.003)
 Ia-all 0.028 (0.002)




Local host galaxy properties - metallicity gradients



CC SNe local metallicity have lower values in the outskirts

SNe Ia do not show a decrease in metallicity at larger distances

Aperture effects




Conclusions

- IFU is a powerful technique at low redshift
- Differences found in galactocentric distances. This can be understood as **differences in the progenitor metallicity**, in sequence from type Ibc/Ilb, type II, and type Ia SNe.
- Differences found in association to star-forming regions. This can be understood as **differences in the progenitor mass and age**, in the same SN type sequence.
- **No differences** found in type Ibc/Ilb and type II SNe **environmental metallicities**, giving support to the progenitor mass and age to determine the SN type. **SNe Ia** occur systematically in **metal-rich environments**.
- Differences between local and integrated values can be understood as uncertainties on the estimations of these parameters at high redshift. Although not so important for SNe Ia.



Present

SINFONI
IMACS
SNIFS
GMOS

...



Kuncarayakti+ 2013ab
Kuncarayakti+ in prep.






Future

New IFUs almost prepared or under construction

- Larger FoV
- Better spatial resolution

3.5 x 3.5 arcmin²





Gràcies