

Probabilistic modelling of protein sub-cellular localisation

Laurent Gatto

laurent.gatto@uclouvain.be

de Duve Institute – UCLouvain slides

<http://bit.ly/20190329ISBA>

29 March 2019 – ISBA


Abstract

In biology, **localisation is function** - understanding the sub-cellular localisation of proteins is paramount to comprehend the context of their functions. Mass spectrometry-based **spatial proteomics** and contemporary **machine learning** enable to build proteome-wide spatial maps, informing us on the location of thousands of proteins. Nevertheless, while some proteins can be found in a single location within a cell, up to half of proteins may reside in multiple locations, can dynamically re-localise, or reside within an unknown functional compartment, leading to considerable **uncertainty** in associating a protein to their sub-cellular location. Recent advances enable us to **probabilistically** model protein localisation as well as quantify the uncertainty in the location assignments, thus leading to better and more trustworthy biological interpretation of the data.

1. **Use case:** spatial proteomics.
2. Novel **computational biology research and developments** to acquire reliable biological knowledge.
3. **Behind the scenes:** software/data structures and open research practice.

Use case: spatial proteomics.

Cell organisation - regulation of protein localisation



Spatial proteomics is the systematic study of protein localisations.

Spatial proteomics - Why?

Localisation is function

- ▶ The cellular sub-division allows cells to establish a range of distinct micro-environments, each favouring different biochemical reactions and interactions and, therefore, allowing each compartment to fulfil a particular functional role.
- ▶ Localisation and sequestration of proteins within sub-cellular niches is a fundamental mechanism for the post-translational regulation of protein function.

Re-localisation in

- ▶ **Differentiation** stem cells.
- ▶ **Activation** of biological processes.

Spatial proteomics - Why?

Mis-localisation

Disruption of the targeting/trafficking process alters proper sub-cellular localisation, which in turn perturb the cellular functions of the proteins.

- ▶ Abnormal protein localisation leading to the **loss of functional** effects in diseases ([Laurila and Vihinen, 2009](#)).
- ▶ Disruption of the nuclear/cytoplasmic transport (nuclear pores) have been detected in many types of **carcinoma cells** ([Kau et al., 2004](#)).
- ▶ Sub-cellular localisation of MC4R with ADCY3 at neuronal primary cilia underlies a common pathway for genetic predisposition to **obesity** ([Siljee et al., 2018](#)).

Spatial proteomics - How, experimentally




Figure: Organelle proteomics approaches ([Gatto et al., 2010](#))

Fusion proteins and immunofluorescence




Figure: Targeted protein localisation. Example of discrepancies between IF and FPs as well as between FP tagging at the N and C termini (Stadler et al., 2013).


Spatial proteomics - How, experimentally

Single cell direct observation	Population level				
	Subcellular fractionation (number of fractions)				
	1 fraction	2 fractions (enriched and crude)	n discrete fractions	n continuous fractions (gradient approaches)	
	GFP Epitope Prot.-spec. antibody	Pure fraction catalogue	Subtractive proteomics (enrichment)	Invariant rich fraction (clustering)	PCP (χ^2)
Cataloguing			Relative abundance		
Tagging	Quantitative mass spectrometry				


Figure: Organelle proteomics approaches ([Gatto et al., 2010](#)).

Gradient approaches: [Dunkley et al. \(2006\)](#), [Foster et al. \(2006\)](#).

Explorative/discovery approaches, [steady-state global localisation maps](#).




Cell lysis




Fractionation/centrifugation

e.g. Mitochondrion



Quantitation/identification
by mass spectrometry

e.g. Mitochondrion



Quantitation data

	Fraction ₁	Fraction ₂	...	Fraction _m
p ₁	q _{1,1}	q _{1,2}	...	q _{1,m}
p ₂	q _{2,1}	q _{2,2}	...	q _{2,m}
p ₃	q _{3,1}	q _{3,2}	...	q _{3,m}
p ₄	q _{4,1}	q _{4,2}	...	q _{4,m}
⋮	⋮	⋮	⋮	⋮
p _j	q _{j,1}	q _{j,2}	...	q _{j, m}

Quantitation data and organelle markers

	Fraction ₁	Fraction ₂	...	Fraction _m	markers
p ₁	q _{1,1}	q _{1,2}	...	q _{1,m}	unknown
p ₂	q _{2,1}	q _{2,2}	...	q _{2,m}	<i>loc₁</i>
p ₃	q _{3,1}	q _{3,2}	...	q _{3,m}	unknown
p ₄	q _{4,1}	q _{4,2}	...	q _{4,m}	<i>loc_i</i>
⋮	⋮	⋮	⋮	⋮	⋮
p _j	q _{j,1}	q _{j,2}	...	q _{j, m}	unknown

Data analysis

Data analysis

- ▶ **Visualisation** (cluster, unsupervised learning)
- ▶ **Classification** (supervised learning)
- ▶ **Novelty detection** (semi-supervised learning)
- ▶ Data integration (transfer learning)
- ▶ **Multi-localisation (Bayesian spatial proteomics)**
- ▶ Spatial dynamics

To uncover and understand biology

Visualisation




Figure: From Gatto et al. (2010), *Arabidopsis thaliana* data from Dunkley et al. (2006)

Supervised Machine Learning






Figure: Support vector machines classifier (after 5% FDR classification cutoff) on the embryonic stem cell data from [Christoforou et al. \(2016\)](#).

Novel **computational biology research and developments** to acquire reliable biological knowledge.

Importance of annotation



Incomplete annotation, and therefore lack of training data, for many/most organelles. *Drosophila* data from [Tan et al. \(2009\)](#).

Semi-supervised learning: novelty detection




Figure: Left: Original *Drosophila* data from Tan et al. (2009). Right: After semi-supervised learning and classification, Breckels et al. (2013).




Figure: The effect of organelle discovery upon sub-cellular protein localisation Breckels et al. (2013).

Computational advances: Transfer learning



What about using **addition data**, such as annotations from the Gene Ontology (GO), sequence features (pseudo aminoacid composition), signal peptide, trans-membrane domains (length, number, ...), images (IF, FP), interaction data, prediction software, ...

- ▶ From a user perspective: "**free/cheap**" vs. expensive and time-consuming experiments.
- ▶ Abundant (all proteins, 100s of features) vs. (experimentally) limited/**targeted** (1000s of proteins, 6 – 20 of features)
- ▶ For localisation in system at hand: *low* vs. high **quality**
- ▶ **Static** vs. **dynamic**

Transfer learning

Support/complement the **primary** target domain (experimental data) with **auxiliary** data (annotation, imaging, PPI, ...) features without compromising the integrity of our primary data.

PRIMARY EXPERIMENTAL DATA



AUXILIARY DRY DATA

Transfer learning results





Figure: From Breckels et al. (2016) *Learning from heterogeneous data sources: an application in spatial proteomics*.

How much do we learn? How much do we miss?



RESEARCH ARTICLE

A Bayesian mixture modelling approach for spatial proteomics

Oliver M. Crook^{1,2,3}, Claire M. Mulvey², Paul D. W. Kirk³, Kathryn S. Lilley², Laurent Gatto^{1,2*}

1 Computational Proteomics Unit, Department of Biochemistry, University of Cambridge, Cambridge, UK, **2** Cambridge Centre for Proteomics, Department of Biochemistry, University of Cambridge, Cambridge, UK, **3** MRC Biostatistics Unit, Cambridge Institute for Public Health, Cambridge, UK

✉ Current address: de Duve Institute, UCLouvain, Brussels, Belgium

* laurent.gatto@uclouvain.be



OPEN ACCESS

Citation: Crook OM, Mulvey CM, Kirk PDW, Lilley KS, Gatto L (2018) A Bayesian mixture modelling approach for spatial proteomics. PLoS Comput Biol 14(11): e1006516. <https://doi.org/10.1371/journal.pcbi.1006516>

Editor: Christine Vogel, NYU, UNITED STATES

Received: May 23, 2018

Accepted: September 17, 2018

Published: November 27, 2018

Copyright: © 2018 Crook et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Analysis of the spatial sub-cellular distribution of proteins is of vital importance to fully understand context specific protein function. Some proteins can be found with a single location within a cell, but up to half of proteins may reside in multiple locations, can dynamically re-localise, or reside within an unknown functional compartment. These considerations lead to uncertainty in associating a protein to a single location. Currently, mass spectrometry (MS) based spatial proteomics relies on supervised machine learning algorithms to assign proteins to sub-cellular locations based on common gradient profiles. However, such methods fail to quantify uncertainty associated with sub-cellular class assignment. Here we reformulate the framework on which we perform statistical analysis. We propose a Bayesian generative classifier based on Gaussian mixture models to assign proteins probabilistically to sub-cellular niches, thus proteins have a probability distribution over sub-cellular locations, with Bayesian computation performed using the expectation-maximisation (EM) algorithm, as well as Markov-chain Monte-Carlo (MCMC). Our methodology allows proteome-wide uncertainty quantification, thus adding a further layer to the analysis of spatial proteomics. Our framework is flexible, allowing many different systems to be analysed and reveals new modelling opportunities for spatial proteomics. We find our methods perform competitively with current state-of-the-art machine learning methods, whilst simultaneously providing

A Bayesian Mixture Modelling Approach For Spatial Proteomics

- ▶ *T Augmented Gaussian Mixture model (TAGM)* is a **multivariate Gaussian generative model** for MS-based spatial proteomics data. It posits that each annotated sub-cellular niche can be modelled by a multivariate Gaussian distribution.

A Bayesian Mixture Modelling Approach For Spatial Proteomics

- ▶ *T Augmented Gaussian Mixture model (TAGM)* is a **multivariate Gaussian generative model** for MS-based spatial proteomics data. It posits that each annotated sub-cellular niche can be modelled by a multivariate Gaussian distribution.
- ▶ With the prior knowledge that many proteins are not captured by known sub-cellular niches, we augment our model with an **outlier component**. Outliers are often dispersed and thus this additional component is described by a heavy-tailed distribution: the multivariate Student's t-distribution, leading us to a *T Augmented Gaussian Mixture model*.

A Bayesian Mixture Modelling Approach For Spatial Proteomics

- ▶ *T Augmented Gaussian Mixture model (TAGM)* is a **multivariate Gaussian generative model** for MS-based spatial proteomics data. It posits that each annotated sub-cellular niche can be modelled by a multivariate Gaussian distribution.
- ▶ With the prior knowledge that many proteins are not captured by known sub-cellular niches, we augment our model with an **outlier component**. Outliers are often dispersed and thus this additional component is described by a heavy-tailed distribution: the multivariate Student's t-distribution, leading us to a *T Augmented Gaussian Mixture model*.
- ▶ This methodology allows proteome-wide **uncertainty quantification**, thus adding a further layer to the analysis of spatial proteomics.

We initially model the distribution of profiles associated with proteins that localise to the k -th component as multivariate normal with mean vector $\boldsymbol{\mu}_k$ and covariance matrix $\boldsymbol{\Sigma}_k$, so that:

$$\mathbf{x}_i | z_i = k \sim \mathcal{N}(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k). \quad (1)$$

We initially model the distribution of profiles associated with proteins that localise to the k -th component as multivariate normal with mean vector $\boldsymbol{\mu}_k$ and covariance matrix $\boldsymbol{\Sigma}_k$, so that:

$$\mathbf{x}_i | z_i = k \sim \mathcal{N}(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k). \quad (1)$$

We extend it by introducing an additional *outlier component*. To do this, we augment our model by introducing a further indicator latent variable ϕ . Each protein \mathbf{x}_i is now described by an additional variable ϕ_i , with $\phi_i = 1$ indicating that protein \mathbf{x}_i belongs to a organelle derived component and $\phi_i = 0$ indicating that protein \mathbf{x}_i is not well described by these known components. This outlier component is modelled as a multivariate T distribution with degrees of freedom κ , mean vector \mathbf{M} , and scale matrix V .

$$\mathbf{x}_i | z_i = k, \phi_i \sim \mathcal{N}(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)^{\phi_i} \mathcal{T}(\kappa, \mathbf{M}, V)^{1-\phi_i}. \quad (2)$$




Figure: Plate diagram specifying the conditional independencies and




Figure: Illustration of how the TAGM model describes the pluripotent mouse embryonic stem cell data. Each ellipse contains a proportion of total probability of a particular multivariate Gaussian density. The outer ellipse contains 99% of the total probability whilst the middle and inner ellipses contain 95% and 90% of the probability respectively.






Figure: Assignment of proteins of *unknown* location to one of the annotated classes. The dots are scaled according to the protein assignment probabilities.









Figure: Exportin 5 (Q924C1) forms part of the micro-RNA export machinery, transporting miRNA from the nucleus to the cytoplasm for further processing. It then translocates back through the nuclear pore complex to return to the nucleus to mediate further transport between nucleus and cytoplasm. The model correctly infers that it most likely localises to the cytosol but there is some uncertainty with this assignment. This uncertainty is reflected in possible assignment of Exportin 5 to the nucleus non-chromatin and reflects the multi-location of the protein.

Whole sub-cellular proteome uncertainty



Spatial dynamics

Trans-localisation event during monocyte to macrophage differentiation

Investigate the effect of lipopolysaccharides (LPS)-mediated inflammatory response in human monocytic cells (THP-1)

Data

- ▶ Triplicate **temporal** profiling (0, 2, 4, 6, 12, 24 hours).
- ▶ Triplicate **spatial** profiling (0 vs 12 hours) - early trafficking, before actual morphological differentiation at 24h.

Work lead by **Dr Claire Mulvey** at the Cambridge Centre for Proteomics.




Figure: Spatial maps of unstimulated and LPS-treated cells (combined triplicates).





Figure: Relocation of Protein Kinase C α and β from the cytosol to the plasma membrane, **driving maturation into a differentiated macrophage phenotype.**




Figure: Relocation of Signal transducer and activator of transcription 6 (STAT6) from the cytosol to the Nucleus, **activating anti-bacterial and anti-viral-like response**. Validated by microscopy and see also [Chen et al. \(2011\)](#).

Behind the scenes: software/data structures and open research practice.

Beyond the figures¹

- ▶ Software: **infrastructure** (MSnbase, Gatto and Lilley (2012)), **dedicated machine learning** (pRoloc, Gatto et al. (2014b)), **interactive visualisation**² (pRolocGUI, Breckels et al. (2017)) and **data** (pRolocdata, Gatto et al. (2014b)) for spatial proteomics.

¹... which are all reproducible, by the way.

²<https://lgatto.shinyapps.io/christoforou2015/>

³between and within domains/software

Beyond the figures¹

- ▶ Software: **infrastructure** (MSnbase, Gatto and Lilley (2012)), **dedicated machine learning** (pRoloc, Gatto et al. (2014b)), **interactive visualisation**² (pRolocGUI, Breckels et al. (2017)) and **data** (pRolocdata, Gatto et al. (2014b)) for spatial proteomics.
- ▶ The **Bioconductor** (Huber et al., 2015) ecosystem for high throughput biology data analysis and comprehension: **open source**, and **coordinated and collaborative**³ **open development**, enabling **reproducible research**, enables understanding of the data (not a black box) and **drive scientific innovation**.

¹... which are all reproducible, by the way.

²<https://lgatto.shinyapps.io/christoforou2015/>

³between and within domains/software

Open research: open source software

The screenshot shows the GitHub repository for **lgatto / pRoloC**. The repository is described as "A unifying bioinformatics framework for organelle proteomics" with a link to <http://lgatto.github.io/pRoloC/>. It has 2,051 commits, 10 branches, 25 releases, 1 environment, and 14 contributors. The repository is in the **master** branch. A table of files is shown below:

File	Description	Latest commit	Time
R	fix to make work with devel		6 day
data	Merge branch 'master' into devel		2 year
inst	updated documentation		6 month
man	new logPosterioris accessor		2 month
src	more C exporting fuss		10 month
tests	add test back		6 day
vignettes	automatic fix of indentation		6 day
_Rbuildignore	ignore .editorconfig when building		3 month
.editorconfig	fix tab with spaces and add editorconfig		4 month
.gitignore	ignore docs, update news		5 month
.travis.yml	Fix indentation		5 month
CONDUCT.md	Merge branch 'master' into devel		3 year
DESCRIPTION	bump devel version on gh		6 day
NAMESPACE	fix notes and warnings		7 day
NEWS	bump devel version on gh		6 day
NEWS.md	bump devel version on gh		6 day

The screenshot shows the Bioconductor website for the **pRoloC** package. The Bioconductor logo is at the top left, and the navigation bar includes **Home**, **Install**, and **Help**. The breadcrumb trail is **Home > Bioconductor 3.8 > Software Packages > pRoloC**. The package name **pRoloC** is displayed in green. Below it, there are statistics: **platforms** (all), **Rank** (254 / 1649), **posts** (1 / 2 / 2 / 0), and **in Bioc** (6 years). There are also buttons for **build**, **warnings**, and **updated < 1 week**. The DOI is [10.18129/BIOC.P0160C](https://doi.org/10.18129/BIOC.P0160C). The description states: "A unifying bioinformatics framework for spatial proteomics". The Bioconductor version is Release (3.8). The description of the package is: "The pRoloC package implements machine learning and visualisation methods for the analysis and interpretation of quantitative mass spectrometry data to reliably infer protein sub-cellular localisation." The authors are Laurent Gatto, Oliver Crook and Lisa M. Breckels, with contributions from Thomas Burger and Samuel Wleczorek. The maintainer is Laurent Gatto. The citation is: "Gatto L, Breckels LM, Wleczorek S, Burger T, Lilley KS (2014). 'Mass-spectrometry based spatial proteomics data analysis using pRoloC and pRoloCdata.' Bioinformatics." The package is also cited in other publications, such as "The effect of organelle discovery upon sub-cellular protein localisation." and "A foundation for reliable spatial proteomics data analysis."

Figure: Gatto et al. (2014a) Left: Public repository for the pRoloC software (<https://github.com/lgatto/pRoloC>). Right: official Bioconductor page.

Open and reproducible research

[Github](#) / [Q&Pep-manuscript](#)

[Download](#)

[Code](#)
[Issues](#)
[Pull requests](#)
[Insights](#)
[Settings](#)

Assessing sub-ocular resolution in spatial proteomics experiments

[problems](#)
[spatial proteomics](#)
[python](#)
[proteomics](#)
[Manage tags](#)

[195](#) commits
 [p1](#) branch
 [0](#) releases

[Branch: master](#)
[New pull request](#)
[Create new file](#)
[Upload files](#)

File	Commit	Message
figs	figs	Update README.md
figs	figs	add data
figs	figs	Update for bioRxiv
figs	figs	add cover letter 2
figs	figs	add cover letter 1
figs	figs	add Travis file
figs	figs	addressing more reviewers comments
figs	figs	add Travis file
figs	figs	Update README.md
figs	figs	add cover letter
figs	figs	add cover letter
figs	figs	add cover letter 2
figs	figs	add cover letter
figs	figs	group assessment section with rib-cluster sims
figs	figs	group assessment section with rib-cluster sims
figs	figs	Calculate group distribution metrics
figs	figs	incorporate Kathryn and Laura's comments
figs	figs	incorporate Kathryn and Laura's comments
figs	figs	minor updates and change marker transfer paragraph
figs	figs	fix table
figs	figs	Update for bioRxiv
figs	figs	changes to new part in col
figs	figs	Update for bioRxiv
figs	figs	Update for bioRxiv
figs	figs	incorporate Kathryn and Laura's comments
figs	figs	incorporate Kathryn and Laura's comments



bioRxiv
THE PREPRINT SERVER FOR BIOLOGY

[HOME](#) | [ABOUT](#) | [SUBMIT](#) | [ALERTS/RSS](#) | [CHANNELS](#)

Advanced Search

New Results

View current version of this article

Assessing sub-cellular resolution in spatial proteomics experiments

Comment on this paper

 Laurent Gatto, Lisa M Breckels, Kathryn S Lilley

doi: <https://doi.org/10.1101/077890>

New preprint in Current Opinion in Cellular Biology doi: [10.1016/j.cob.2018.11.015](https://doi.org/10.1016/j.cob.2018.11.015)

Abstract

Info/History

Metrics


 Download PDF


Abstract


The sub-cellular localisation of a protein is vital in defining its function, and a protein's mis-localisation is known to lead to adverse effect. As a result, numerous experimental techniques and datasets have been published, with the aim of deciphering the localisation of proteins at various scales and resolutions, including high profile mass spectrometry-based efforts. Here, we present a meta-analysis assessing and comparing the sub-cellular resolution of 29 such mass spectrometry-based spatial proteomics experiments using a newly developed tool termed QSub. Our goal is to provide a simple quantitative report of how well spatial proteomics resolve the sub-cellular niches they describe to inform data developers and users of such methods.


Copyright


The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.


 ScienceDirect

 Outline


 Download


 Share

 Export

 Current Opinion in Chemical Biology


Volume 48, February 2019, Pages 123–149

 ELSEVIER



Assessing sub-cellular resolution in spatial proteomics experiments


Laurent Gatto ^{1,2,3,7}*, Lisa M. Breckels ^{1,2}, Kathryn S. Lilley ²

 [Show more](#)

<https://doi.org/10.1016/j.cop.2018.11.015>

Get rights and content

Under a Creative Commons license

 open access

Abstract

The **sub-cellular localisation** of a protein is vital in defining its function, and a protein's mis-localisation is known to lead to adverse effect. As a result, numerous **experimental techniques** and datasets have been published, with the aim of deciphering the localisation of proteins at various scales and resolutions, including high profile mass spectrometry-based efforts. Here, we present a meta-analysis assessing and comparing the sub-cellular resolution of 29 such mass spectrometry-based spatial **proteomics** experiments using a newly developed tool termed **QSeq**. Our goal is to provide a simple quantitative report of how well spatial proteomics resolve the sub-cellular niches they describe to inform and guide developers and users of such methods.

Figure: Gatto et al. (2018) reproducible document
(<https://github.com/lgatto/QSep-manuscript>), preprint
(<https://doi.org/10.1101/377630>) and paper
(<https://doi.org/10.1016/j.cbpa.2018.11.015>).

Working with open and reproducible research in mind doesn't mean releasing everything prematurely, it means

- ▶ managing research in a way one can find data and results at every stage
- ▶ one can reproduce results, re-run/compare them with new data or different methods/parameters, and
- ▶ one can release data (or parts thereof) when/if appropriate.

Conclusions

- ▶ Protein sub-cellular localisation: technologies (hyperLOPIT) and opportunities.
- ▶ Reliance on computational biology and dedicated software (pRoLoc *et al.*) to interpret data and acquire biological knowledge.
- ▶ Rigorous computational infrastructure and sound data analysis and interpretation is a **long term investment**.

References I

- L M Breckels, S B Holden, D Wojnar, C M Mulvey, A Christoforou, A Groen, M W Trotter, O Kohlbacher, K S Lilley, and L Gatto. Learning from heterogeneous data sources: An application in spatial proteomics. *PLoS Comput Biol*, 12(5):e1004920, May 2016. doi: 10.1371/journal.pcbi.1004920.
- Lisa Breckels, Thomas Naake, and Laurent Gatto. *pRolocGUI: Interactive visualisation of spatial proteomics data*, 2017. URL <http://ComputationalProteomicsUnit.github.io/pRolocGUI/>. R package version 1.11.2.
- LM Breckels, L Gatto, A Christoforou, AJ Groen, KS Lilley, and MW Trotter. The effect of organelle discovery upon sub-cellular protein localisation. *J Proteomics*, 88:129–40, Aug 2013.
- H Chen, H Sun, F You, W Sun, X Zhou, L Chen, J Yang, Y Wang, H Tang, Y Guan, W Xia, J Gu, H Ishikawa, D Gutman, G Barber, Z Qin, and Z Jiang. Activation of stat6 by sting is critical for antiviral innate immunity. *Cell*, 147(2):436–46, Oct 2011. doi: 10.1016/j.cell.2011.09.022.
- A Christoforou, C M Mulvey, L M Breckels, A Geladaki, T Hurrell, P C Hayward, T Naake, L Gatto, R Viner, A Martinez Arias, and K S Lilley. A draft map of the mouse pluripotent stem cell spatial proteome. *Nat Commun*, 7:8992, Jan 2016. doi: 10.1038/ncomms9992.
- TPJ Dunkley, S Hester, IP Shadforth, J Runions, T Weimar, SL Hanton, JL Griffin, C Bessant, F Brandizzi, C Hawes, RB Watson, P Dupree, and KS Lilley. Mapping the Arabidopsis organelle proteome. *PNAS*, 103(17):6518–6523, Apr 2006.
- LJ Foster, CL de Hoog, Y Zhang, Y Zhang, X Xie, VK Mootha, and M Mann. A mammalian organelle map by protein correlation profiling. *Cell*, 125(1):187–199, Apr 2006.
- L Gatto and KS Lilley. MSnbase - an R/Bioconductor package for isobaric tagged mass spectrometry data visualization, processing and quantitation. *Bioinformatics*, 28(2):288–9, Jan 2012.
- L Gatto, JA Vizcaino, H Hermjakob, W Huber, and KS Lilley. Organelle proteomics experimental designs and analysis. *Proteomics*, 2010.
- L Gatto, L M Breckels, S Wiczorek, T Burger, and K S Lilley. Mass-spectrometry based spatial proteomics data analysis using pRoloc and pRolocdata. *Bioinformatics*, Jan 2014a.
- L Gatto, LM Breckels, T Burger, DJ Nightingale, AJ Groen, C Campbell, N Nikolovski, CM Mulvey, A Christoforou, M Ferro, and KS Lilley. A foundation for reliable spatial proteomics data analysis. *MCP*, 13(8):1937–52, Aug 2014b.

References II

- Laurent Gatto, Lisa M Breckels, and Kathryn S Lilley. Assessing sub-cellular resolution in spatial proteomics experiments. *bioRxiv*, 2018. doi: 10.1101/377630.
- W Huber, V J Carey, R Gentleman, S Anders, M Carlson, B S Carvalho, H C Bravo, S Davis, L Gatto, T Girke, R Gottardo, F Hahne, K D Hansen, R A Irizarry, M Lawrence, M I Love, J MacDonald, V Obenchain, A K Oleś, H Pagès, A Reyes, P Shannon, G K Smyth, D Tenenbaum, L Waldron, and M Morgan. Orchestrating high-throughput genomic analysis with Bioconductor. *Nat Methods*, 12(2):115–21, Jan 2015. doi: 10.1038/nmeth.3252.
- TR Kau, JC Way, and PA Silver. Nuclear transport and cancer: from mechanism to intervention. *Nat Rev Cancer*, 4(2):106–17, Feb 2004.
- K Laurila and M Vihinen. Prediction of disease-related mutations affecting protein localization. *BMC Genomics*, 10:122, 2009.
- J E Siljee, Y Wang, A A Bernard, B A Ersoy, S Zhang, A Marley, M Von Zastrow, J F Reiter, and C Vaisse. Subcellular localization of mc4r with adcy3 at neuronal primary cilia underlies a common pathway for genetic predisposition to obesity. *Nat Genet*, Jan 2018. doi: 10.1038/s41588-017-0020-9.
- C Stadler, E Rexhepaj, V R Singan, R F Murphy, R Pepperkok, M Uhlén, J C Simpson, and E Lundberg. Immunofluorescence and fluorescent-protein tagging show high correlation for protein localization in mammalian cells. *Nat Methods*, 10(4):315–23, Apr 2013.
- DJL Tan, H Dvinge, A Christoforou, P Bertone, A Arias Martinez, and KS Lilley. Mapping organelle proteins and protein complexes in *Drosophila melanogaster*. *J Proteome Res*, 8(6):2667–2678, Jun 2009.

Acknowledgements

- ▶ **Mr Oliver Crook and Dr Lisa Breckels**, (U of Cambridge): spatial proteomics, machine learning, software.
- ▶ **Dr Sebastian Gibb and Dr Johannes Rainer**: MS and proteomics software.
- ▶ Prof Kathryn Lilley (U of Cambridge), Dr Claire Mulvey, (CRUK Cambridge Institute): data.
- ▶ Funding: BBSRC, Wellcome Trust

Thank you for your attention