

# Mapping the sub-cellular proteome

Probabilistic modelling of protein sub-cellular localisation

Laurent Gatto

`laurent.gatto@uclouvain.be`


`http://lgatto.github.io/about`

de Duve Institute – UCLouvain

**Protein Folding and Stability**

30 August 2019 – Liège

# Cell organisation - localisation is function




**Spatial proteomics** is the systematic study of protein localisations.

**Localisation – re-localisation – mis-localisation**

**Explorative/discovery approaches, steady-state global localisation maps** (as opposed to microscopy-based targeted approaches).

**Density gradient:** PCP ([Dunkley et al., 2006](#)), LOPIT ([Foster et al., 2006](#)), hyperLOPIT ([Christoforou et al., 2016](#); [Mulvey et al., 2017](#)) and

**Differential centrifugation** [Itzhak et al. \(2016\)](#), LOPIT-DC ([Geladaki et al., 2018](#)).



## Quantitation data

	Fraction <sub>1</sub>	Fraction <sub>2</sub>	...	Fraction <sub>L</sub>
<b>x<sub>1</sub></b>	x <sub>1,1</sub>	x <sub>1,2</sub>	...	x <sub>1,L</sub>
<b>x<sub>2</sub></b>	x <sub>2,1</sub>	x <sub>2,2</sub>	...	x <sub>2,L</sub>
<b>x<sub>3</sub></b>	x <sub>3,1</sub>	x <sub>3,2</sub>	...	x <sub>3,L</sub>
⋮	⋮	⋮	⋮	⋮
<b>x<sub>i</sub></b>	x <sub>i,1</sub>	x <sub>i,2</sub>	...	x <sub>i,L</sub>
⋮	⋮	⋮	⋮	⋮
<b>x<sub>N</sub></b>	x <sub>N,1</sub>	x <sub>N,2</sub>	...	x <sub>N, L</sub>

## Quantitation data and organelle markers

	Fraction <sub>1</sub>	Fraction <sub>2</sub>	...	Fraction <sub>L</sub>	markers
$x_1$	$x_{1,1}$	$x_{1,2}$	...	$x_{1,L}$	unknown
$x_2$	$x_{2,1}$	$x_{2,2}$	...	$x_{2,L}$	<i>loc<sub>1</sub></i>
$x_3$	$x_{3,1}$	$x_{3,2}$	...	$x_{3,L}$	unknown
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$x_i$	$x_{i,1}$	$x_{i,2}$	...	$x_{i,L}$	<i>loc<sub>k</sub></i>
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$x_N$	$x_{N,1}$	$x_{N,2}$	...	$x_{N,K}$	unknown

# Visualisation






Figure: From Gatto et al. (2010), *Arabidopsis thaliana* data from Dunkley et al. (2006)

# Supervised Machine Learning



**Figure:** Support vector machines classifier (after 5% FDR classification cutoff) on the embryonic stem cell data from [Christoforou et al. \(2016\)](#).

# How much do we learn? How much do we miss?





# A Bayesian Mixture Modelling Approach For Spatial Proteomics

- ▶ *T* Augmented Gaussian Mixture model (TAGM) is a **multivariate Gaussian generative model** for MS-based spatial proteomics data. It posits that each annotated sub-cellular niche can be modelled by a multivariate Gaussian distribution.

# A Bayesian Mixture Modelling Approach For Spatial Proteomics

- ▶ *T Augmented Gaussian Mixture model (TAGM)* is a **multivariate Gaussian generative model** for MS-based spatial proteomics data. It posits that each annotated sub-cellular niche can be modelled by a multivariate Gaussian distribution.
- ▶ With the prior knowledge that many proteins are not captured by known sub-cellular niches, we augment our model with an **outlier component**. Outliers are often dispersed and thus this additional component is described by a heavy-tailed distribution: the multivariate Student's t-distribution, leading us to a *T Augmented Gaussian Mixture model* (Crook et al., 2018, 2019).

# A Bayesian Mixture Modelling Approach For Spatial Proteomics

- ▶ *T Augmented Gaussian Mixture model (TAGM)* is a **multivariate Gaussian generative model** for MS-based spatial proteomics data. It posits that each annotated sub-cellular niche can be modelled by a multivariate Gaussian distribution.
- ▶ With the prior knowledge that many proteins are not captured by known sub-cellular niches, we augment our model with an **outlier component**. Outliers are often dispersed and thus this additional component is described by a heavy-tailed distribution: the multivariate Student's t-distribution, leading us to a *T Augmented Gaussian Mixture model* (Crook et al., 2018, 2019).
- ▶ This methodology allows proteome-wide **uncertainty quantification**, thus adding a further layer to the analysis of spatial proteomics.

We initially model the distribution of profiles associated with proteins that localise to the  $k$ -th component as multivariate normal with mean vector  $\boldsymbol{\mu}_k$  and covariance matrix  $\boldsymbol{\Sigma}_k$ , so that:



$$\mathbf{x}_i | z_i = k \quad \sim \mathcal{N}(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \quad (1)$$

We initially model the distribution of profiles associated with proteins that localise to the  $k$ -th component as multivariate normal with mean vector  $\boldsymbol{\mu}_k$  and covariance matrix  $\boldsymbol{\Sigma}_k$ , so that:


$$\mathbf{x}_i | z_i = k \sim \mathcal{N}(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k) \quad (1)$$


We extend it by introducing an additional *outlier component*. To do this, we augment our model by introducing a further indicator latent variable  $\phi$ . Each protein  $\mathbf{x}_i$  is now described by an additional variable  $\phi_i$ , with  $\phi_i = 1$  indicating that protein  $\mathbf{x}_i$  belongs to a organelle derived component and  $\phi_i = 0$  indicating that protein  $\mathbf{x}_i$  is not well described by these known components. This outlier component is modelled as a multivariate T distribution with degrees of freedom  $\kappa$ , mean vector  $\mathbf{M}$ , and scale matrix  $V$ .

$$\mathbf{x}_i | z_i = k, \phi_i \sim \mathcal{N}(\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)^{\phi_i} \mathcal{T}(\kappa, \mathbf{M}, V)^{1-\phi_i} \quad (2)$$



**Figure:** Assignment of proteins of *unknown* location to one of the annotated classes. The dots are scaled according to the protein assignment probabilities.








**Figure:** Exportin 5 (Q924C1) forms part of the micro-RNA export machinery, transporting miRNA from the nucleus to the cytoplasm for further processing. It then translocates back through the nuclear pore complex to return to the nucleus to mediate further transport between nucleus and cytoplasm. The model correctly infers that it most likely localises to the cytosol but there is some uncertainty with this assignment. This uncertainty is reflected in possible assignment of Exportin 5 to the nucleus non-chromatin and reflects the multi-location of the protein.



## Whole sub-cellular proteome uncertainty



# Spatial dynamics


## Trans-localisation event during monocyte to macrophage differentiation

Investigate the effect of lipopolysaccharides (LPS)-mediated inflammatory response in human monocytic cells (THP-1)


### Data

- ▶ Triplicate **temporal** profiling (0, 2, 4, 6, 12, 24 hours).
- ▶ Triplicate **spatial** profiling (0 vs 12 hours) - early trafficking, before actual morphological differentiation at 24h.


Work lead by **Dr Claire Mulvey** at the Cambridge Centre for Proteomics.



**Figure:** Spatial maps of unstimulated and LPS-treated cells (combined triplicates).



**Figure:** Relocation of Protein Kinase C  $\alpha$  and  $\beta$  from the cytosol to the plasma membrane, **driving maturation into a differentiated macrophage phenotype.**



**Figure:** Relocation of Signal transducer and activator of transcription 6 (STAT6) from the cytosol to the Nucleus, **activating anti-bacterial and anti-viral-like response**. Validated by microscopy and see also [Chen et al. \(2011\)](#).

## Folding and stability

- ▶ Re-localisation upon protein post-translational modification (PTM).
- ▶ Effect of PTM on protein 3D structure.
- ▶ Link between 3D structure and localisation.

**Behind the scenes:** software/data structures and open research practice.

## Beyond the figures<sup>1</sup>

- ▶ Software: **infrastructure** (MSnbase, Gatto and Lilley (2012)), **dedicated machine learning** (pRoloc, Gatto et al. (2014b)), **interactive visualisation**<sup>2</sup> (pRolocGUI, Breckels et al. (2017)) and **data** (pRolocdata, Gatto et al. (2014b)) for spatial proteomics.

---

<sup>1</sup>... which are all reproducible, by the way.

<sup>2</sup><https://lgatto.shinyapps.io/christoforou2015/>

<sup>3</sup>between and within domains/software



## Beyond the figures<sup>1</sup>

- ▶ Software: **infrastructure** (MSnbase, Gatto and Lilley (2012)), **dedicated machine learning** (pRoloc, Gatto et al. (2014b)), **interactive visualisation**<sup>2</sup> (pRolocGUI, Breckels et al. (2017)) and **data** (pRolocdata, Gatto et al. (2014b)) for spatial proteomics.
- ▶ The **Bioconductor** (Huber et al., 2015) ecosystem for high throughput biology data analysis and comprehension: **open source**, and **coordinated and collaborative**<sup>3</sup> **open development**, enabling **reproducible research**, enables understanding of the data (not a black box) and **drive scientific innovation**.

---

<sup>1</sup>... which are all reproducible, by the way.

<sup>2</sup><https://lgatto.shinyapps.io/christoforou2015/>

<sup>3</sup>between and within domains/software

# Open research: open source software

The screenshot shows the GitHub repository for **lgatto / pRoloc**. The repository is described as "A unifying bioinformatics framework for organelle proteomics" with a link to <http://lgatto.github.io/pRoloc/>. It has 2,051 commits, 10 branches, 25 releases, 1 environment, and 14 contributors. The repository includes a table of files and their commit history:

File	Commit Message	Time
R	fix to make work with devel	6 day
data	Merge branch 'master' into devel	2 year
inst	updated documentation	6 month
man	new logPosterior accessor	2 month
src	more C exporting luss	10 month
tests	add test back	6 day
vignettes	automatic fix of indentation	6 day
_Rbuildignore	ignore .editorconfig when building	3 month
.editorconfig	fix tab with spaces and add editorconfig	4 month
gitignore	ignore docs, update news	5 month
travis.yml	Fix indentation	5 month
CONDUCT.md	Merge branch 'master' into devel	3 year
DESCRIPTION	bump devel version on gh	6 day
NAMESPACE	fix notes and warnings	7 day
NEWS	bump devel version on gh	6 day
NEWS.md	bump devel version on gh	6 day

The screenshot shows the Bioconductor website for the **pRoloc** package. The Bioconductor logo is at the top, with the tagline "OPEN SOURCE SOFTWARE FOR BIOINFORMATICS". The navigation bar includes "Home", "Install", and "Help". The main content area shows the package name **pRoloc** and the version **3.8**. It includes a table of statistics:

platforms	all	fork	254 / 164	posts	1 / 2 / 2 / 0	in BiOC	6 years
build	warnings	updated	< 1 week				

The DOI is [10.18129/B3.bioc.pRoloc](https://doi.org/10.18129/B3.bioc.pRoloc). The description states: "A unifying bioinformatics framework for spatial proteomics". The author is Laurent Gatto, Oliver Crook and Lisa M. Breckels, with contributions from Thomas Burger and Samuel Wiecek. The citation is: Gatto L, Breckels LM, Wiecek S, Burger T, Lilley KS (2014). "Mass-spectrometry based spatial proteomics data analysis using pRoloc and pRolocdata." *Bioinformatics*.

Figure: Gatto et al. (2014a) Left: Public repository for the pRoloc software (<https://github.com/lgatto/pRoloc>). Right: official Bioconductor page.

# Open and reproducible research

The figure displays three screenshots illustrating the open and reproducible research workflow for the project "Assessing sub-cellular resolution in spatial proteomics experiments".

**Left Screenshot (GitHub):** Shows the repository `lgatto/QSep-manuscript`. The file tree includes:

- `data`: add marker transfer code/figs
- `figure`: Update for bioRxiv
- `pipeline`: add cover letter 2
- `travis.yml`: add travis file
- `Makefile`: addressing more reviewers comments
- `README.md`: Update README.md
- `cover.pdf`: add cover letter
- `cover.tex`: add cover letter
- `cover2.pdf`: add cover letter 2
- `chlam_rdx`: qsep assessment section with rib cluster sims
- `hdm_rdx`: qsep assessment section with rib cluster sims
- `mk.R`: Calculate qsep distribution medians
- `reknash_qsep.pdf`: incorporate Kathryn and Lisa's comments
- `reknash_qsep.pdf`: incorporate Kathryn and Lisa's comments
- `reknash_qsep.pdf`: minor updates and change marker transfer paragraph
- `qsep.R`: fix table
- `qsep.Rnw`: Update for bioRxiv
- `qsep.bib`: changes to new part in col
- `qsep.pdf`: Update for bioRxiv
- `qsep.tex`: Update for bioRxiv
- `sim.pdf`: incorporate Kathryn and Lisa's comments
- `sim.R`: incorporate Kathryn and Lisa's comments

**Middle Screenshot (bioRxiv):** Shows the preprint page for "Assessing sub-cellular resolution in spatial proteomics experiments" by Laurent Gatto, Lisa M. Breckels, and Kathryn S. Lilley. The page includes a "New Results" banner and a "View current version of this article" link.

**Right Screenshot (ScienceDirect):** Shows the published paper in "Current Opinion in Chemical Biology" (Volume 48, February 2019, Pages 123-149). The page includes the title, authors, and an abstract section.

Figure: Gatto et al. (2018) reproducible document  
(<https://github.com/lgatto/QSep-manuscript>), preprint  
(<https://doi.org/10.1101/377630>) and paper  
(<https://doi.org/10.1016/j.cbpa.2018.11.015>).

Working with open and reproducible research in mind doesn't mean releasing everything prematurely, it means

- ▶ managing research in a way one can find data and results at every stage
- ▶ one can reproduce results, re-run/compare them with new data or different methods/parameters, and
- ▶ one can release data (or parts thereof) when/if appropriate.

# Conclusions

- ▶ Protein sub-cellular localisation: *localisation is function*.
- ▶ Reliance on computational biology, statistics and dedicated software (for example MSnbase ([Gatto and Lilley, 2012](#)), pRoLoc ([Gatto et al., 2014a](#))) to interpret data and acquire biological knowledge (details not shown).
- ▶ Rigorous computational infrastructure and sound data analysis and interpretation is a **long term investment**.

# References |

- Lisa Breckels, Thomas Naake, and Laurent Gatto. *pRolocGUI: Interactive visualisation of spatial proteomics data*, 2017. URL <http://ComputationalProteomicsUnit.github.io/pRolocGUI/>. R package version 1.11.2.
- H Chen, H Sun, F You, W Sun, X Zhou, L Chen, J Yang, Y Wang, H Tang, Y Guan, W Xia, J Gu, H Ishikawa, D Gutman, G Barber, Z Qin, and Z Jiang. Activation of stat6 by sting is critical for antiviral innate immunity. *Cell*, 147(2):436–46, Oct 2011. doi: 10.1016/j.cell.2011.09.022.
- A Christoforou, C M Mulvey, L M Breckels, A Geladaki, T Hurrell, P C Hayward, T Naake, L Gatto, R Viner, A Martinez Arias, and K S Lilley. A draft map of the mouse pluripotent stem cell spatial proteome. *Nat Commun*, 7:8992, Jan 2016. doi: 10.1038/ncomms9992.
- Oliver M. Crook, Claire M. Mulvey, Paul D. W. Kirk, Kathryn S. Lilley, and Laurent Gatto. A bayesian mixture modelling approach for spatial proteomics. *PLOS Computational Biology*, 14(11):1–29, 11 2018. doi: 10.1371/journal.pcbi.1006516. URL <https://doi.org/10.1371/journal.pcbi.1006516>.
- OM Crook, LM Breckels, KS Lilley, PDW Kirk, and L Gatto. A bioconductor workflow for the bayesian analysis of spatial proteomics [version 1; peer review: 1 approved, 2 approved with reservations]. *F1000Research*, 8(446), 2019. doi: 10.12688/f1000research.18636.1.

## References II

- TPJ Dunkley, S Hester, IP Shadforth, J Runions, T Weimar, SL Hanton, JL Griffin, C Bessant, F Brandizzi, C Hawes, RB Watson, P Dupree, and KS Lilley. Mapping the Arabidopsis organelle proteome. *PNAS*, 103(17):6518–6523, Apr 2006.
- LJ Foster, CL de Hoog, Y Zhang, Y Zhang, X Xie, VK Mootha, and M Mann. A mammalian organelle map by protein correlation profiling. *Cell*, 125(1):187–199, Apr 2006.
- L Gatto and KS Lilley. MSnbase - an R/Bioconductor package for isobaric tagged mass spectrometry data visualization, processing and quantitation. *Bioinformatics*, 28(2):288–9, Jan 2012.
- L Gatto, JA Vizcaino, H Hermjakob, W Huber, and KS Lilley. Organelle proteomics experimental designs and analysis. *Proteomics*, 2010.
- L Gatto, L M Breckels, S Wiczorek, T Burger, and K S Lilley. Mass-spectrometry based spatial proteomics data analysis using pRoloc and pRolocdata. *Bioinformatics*, Jan 2014a.
- L Gatto, LM Breckels, T Burger, DJ Nightingale, AJ Groen, C Campbell, N Nikolovski, CM Mulvey, A Christoforou, M Ferro, and KS Lilley. A foundation for reliable spatial proteomics data analysis. *MCP*, 13(8):1937–52, Aug 2014b.
- Laurent Gatto, Lisa M Breckels, and Kathryn S Lilley. Assessing sub-cellular resolution in spatial proteomics experiments. *bioRxiv*, 2018. doi: 10.1101/377630.

# References III

- Aikaterini Geladaki, Nina Koccevar Britovsek, Lisa M. Breckels, Tom S. Smith, Claire M. Mulvey, Oliver M. Crook, Laurent Gatto, and Kathryn S. Lilley. LOPIT-DC: A simpler approach to high-resolution spatial proteomics. *bioRxiv*, 2018. doi: 10.1101/378364. URL <https://www.biorxiv.org/content/early/2018/07/26/378364>.
- W Huber, V J Carey, R Gentleman, S Anders, M Carlson, B S Carvalho, H C Bravo, S Davis, L Gatto, T Girke, R Gottardo, F Hahne, K D Hansen, R A Irizarry, M Lawrence, M I Love, J MacDonald, V Obenchain, A K Oleś, H Pagès, A Reyes, P Shannon, G K Smyth, D Tenenbaum, L Waldron, and M Morgan. Orchestrating high-throughput genomic analysis with Bioconductor. *Nat Methods*, 12(2):115–21, Jan 2015. doi: 10.1038/nmeth.3252.
- D N Itzhak, S Tyanova, J Cox, and G H Borner. Global, quantitative and dynamic mapping of protein subcellular localization. *Elife*, 5, Jun 2016. doi: 10.7554/eLife.16950.
- C M Mulvey, L M Breckels, A Geladaki, N K Britovek, DJH Nightingale, A Christoforou, M Elzek, M J Deery, L Gatto, and K S Lilley. Using hyperlopit to perform high-resolution mapping of the spatial proteome. *Nat Protoc*, 12(6): 1110–1135, Jun 2017. doi: 10.1038/nprot.2017.026.



## Acknowledgements

- ▶ **Mr Oliver Crook** and **Dr Lisa Breckels**, (U of Cambridge): spatial proteomics, machine learning, software.
- ▶ Funding: BBSRC (UK), Wellcome Trust (UK), FNRS (BE)

**Thank you for your attention**