

## Corotating Clumps Around Adolescent Low-Mass Stars: Four Years of Complex Quasiperiodic Variables from TESS

LUKE G. BOUMA,<sup>1,\*</sup> RAHUL JAYARAMAN,<sup>2</sup> SAUL RAPPAPORT,<sup>2</sup> LUISA M. REBULL,<sup>1</sup> ALEXANDRE DAVID-URAZ,<sup>3,4</sup>

LYNNE A. HILLENBRAND,<sup>1</sup> GÁSPÁR Á. BAKOS,<sup>5</sup>

—SUGGESTED CONTRIBUTORS, PENDING DISCUSSION, COMMENTS, ANALYSES, ETC—,<sup>6</sup>

MAXIMILIAN N. GÜNTHER,<sup>6</sup> GEORGE R. RICKER,<sup>2</sup> AND JOSHUA N. WINN<sup>5</sup>

<sup>1</sup>Department of Astronomy, MC 249-17, California Institute of Technology, Pasadena, CA 91125, USA

<sup>2</sup>MIT Kavli Institute and Department of Physics, 77 Massachusetts Avenue, Cambridge, MA 02139

<sup>3</sup>Department of Physics and Astronomy, Howard University, Washington DC, 20059

<sup>4</sup>Center for Research and Exploration in Space Science and Technology, and X-ray Astrophysics Laboratory, NASA/GSFC, Greenbelt, MD 20771, USA

<sup>5</sup>Department of Astrophysical Sciences, Princeton University, 4 Ivy Lane, Princeton, NJ 08540, USA

<sup>6</sup>European Space Agency (ESA), European Space Research and Technology Centre (ESTEC), Keplerlaan 1, 2201 AZ Noordwijk, Netherlands

(Received —; Revised —; Accepted —)

### ABSTRACT

Complex quasiperiodic variables (CQVs) are low-mass pre-main-sequence stars with nearly periodic optical modulation. The modulation is likely induced by dust or gas clumps in orbit at the Keplerian corotation radius. Here, we report new CQVs discovered in TESS short-cadence data collected between July 2018 and Sep 2022. Our search of 65,760 K and M dwarfs with  $T < 16$  and  $d < 150$  pc yielded 50 high-quality CQVs. Most of these discoveries are new, and they include the brightest ( $T \approx 9.5$ ), closest ( $d \approx 20$  pc), and oldest ( $\approx 200$  Myr) examples of this class of object currently known. A few objects are outliers: LP 12-502 for instance showed a “dip complex” with a period and duration that were fixed over 1,500 cycles, but in detail, this system exhibited between four and eight local minima per cycle, and at times also displayed multiple periods simultaneously. We demonstrate that transient corotating material is the most viable explanation for this class of object, and present evidence supporting the hypothesis that this material is sculpted by stellar magnetic fields dominated by high order multipoles. We expect that our sample will facilitate future efforts aimed at connecting CQVs to the broader contexts of star, disk, and perhaps even exoplanet evolution.

**Keywords:** Weak-line T Tauri stars (1795), Periodic variable stars (1213), Circumstellar matter (241), Star clusters (1567), Stellar magnetic fields (1610), Stellar rotation (1629)

### 1. INTRODUCTION

All pre-main-sequence stars vary in optical brightness, and the origin of such variability is, in most cases, understood. Well-explored sources of optical variability include inhomogeneities on stellar surfaces such as starspots and faculae (e.g. Basri 2021), occultations by gas-rich circumstellar disks (e.g. Bodman et al. 2017), and, in geometrically favorable circumstances, eclipses by stars and planets (e.g. Winn 2010). More exotic forms of optical variability relevant to this work include transiting exocomets (e.g.  $\beta$  Pic; Zieba et al. 2019) and disintegrating rocky bodies around both M dwarfs (e.g. KOI-2700; Rappaport et al. 2014) and white dwarfs (e.g. WD 1145; Vanderburg et al. 2015).

Corresponding author: Luke G. Bouma  
luke@astro.caltech.edu

\* 51 Pegasi b Fellow

Data from K2 and TESS have yielded a new class of variable star whose root cause is only beginning to become clear: complex quasiperiodic variables (CQVs). These objects are identified from their optical light curves, which show nearly periodic troughs that are either sharp or broad; these troughs are often superposed on smooth spot-like modulation (Stauffer et al. 2017, 2018; Zhan et al. 2019). Some CQVs show up to eight local minima (“dips”) per cycle. Most are pre-main-sequence M dwarfs without near-infrared excesses, ages of  $\approx 5\text{--}150$  million years (Myr), and rotation periods of at most two days; they are observed to comprise  $\approx 1\text{--}3\%$  of M dwarfs younger than 100 Myr (Rebull et al. 2016; Günther et al. 2022). The dips can be chromatic, with a reddening law plausibly consistent with dust (Onitsuka et al. 2017; Bouma et al. 2020; Günther et al. 2022; Koen 2023). And finally, while the dip shapes can “jump” between different depths and durations over less than one cycle, they more often evolve gradually, over tens to hundreds of cycles (e.g. Stauffer et al. 2017; Palumbo et al. 2022; Popinchalk et al. 2023).

The CQV light curves cannot be explained by pure starspots. Although all young M dwarfs are spotted, this tends to produce flux variations over timescales of the rotation period,  $P_{\text{rot}}$ , and its half-harmonic,  $0.5 P_{\text{rot}}$ . With fine-tuned viewing geometries, starspots can produce dip durations as short as  $\approx 0.2 P_{\text{rot}}$ , and the dip amplitudes in such geometries are small due to limb-darkening (see Stauffer et al. 2017, Figures 37-41). The observed dips occur over durations as short as  $0.05 P_{\text{rot}}$  with characteristic depths of a few percent; a “starspot-only” scenario can be discarded for any object with dips that are sufficiently large in depth and short in duration (Stauffer et al. 2017; Koen 2021). The dip durations and amplitudes instead require sharp geometries with material extrinsic to the stellar surface (e.g. Stauffer et al. 2017; Günther et al. 2022). Since some CQVs require such geometries, and the CQVs probably all have a single explanation, we therefore discard the “spot-only” scenario.

A few competing explanations for what causes the complex quasiperiodic variability are shown in Figure 1. The “clump” scenario invokes opaque dust clumps periodically transiting the star near the Keplerian corotation radius,  $R_{\text{cr}} = (GM/\Omega^2)^{1/3}$  (Stauffer et al. 2017; Sanderson et al. 2023). The “prominence” scenario invokes long-lived condensations of cool and dense marginally-ionized gas, embedded within the hotter corona, that would be centrifugally supported to also be near corotation (Collier Cameron & Robinson 1989; Jardine & Collier Cameron 2019; Waugh & Jardine 2022). Such structures are analogous to quiescent prominences and filaments seen in the solar corona (see e.g. Vial & Engvold 2015), though at much larger relative distances from the stellar surface. A final possibility is one of a “screen”, in which the inner wall of a quiescent circumstellar disk blocks a portion of the stellar surface to produce sudden dips whenever spots come into view (Zhan et al. 2019).

The dust clump and prominence hypotheses seem most plausible. They are qualitatively similar, except that one invokes opacity from dust, while the other invokes opacity from gas, likely bound-free transitions in hydrogen or perhaps a molecular opacity. The screen scenario seems inconsistent with the sharpness of the observed dips, the lack of infrared excess, and the observed lifetime of CQVs extending an order of magnitude longer than the  $\approx 10^7$  year timescale often quoted for primordial disk dispersal. However, unambiguous evidence in support of any one scenario has yet to be acquired. Such evidence might include a spectroscopic detection of silicate  $10 \mu\text{m}$  dust absorption during a dip, or perhaps detection of transient Balmer-line excesses as a function of cycle phase, similar to observations made in systems such as AB Dor (see Collier Cameron 1999) or PTFO 8-8695 (Johns-Krull et al. 2016).

CQVs remain mysterious because they have been both hard to discover and hard to characterize. Discoverability is tied to rarity: CQVs comprise  $\approx 1\%$  of the youngest  $\approx 1\%$  of M dwarfs (Rebull et al. 2018). Out of the millions of stars monitored by K2 and TESS, about 50 CQVs have been reported to date (Rebull et al. 2016; Stauffer et al. 2017, 2018; Zhan et al. 2019; Bouma et al. 2020; Günther et al.

2022; Popinchalk et al. 2023). The known CQVs are correspondingly faint; the initial K2 discoveries (Rebull et al. 2016; Stauffer et al. 2017) were M2-M6 dwarfs at distances  $\gtrsim 100$  pc, yielding optical brightnesses of  $V \approx 15.5$  to  $V > 20$ . This renders time-series spectroscopy at high resolution out of reach for current facilities, despite its potential utility in ruling out some of these models.

In this work, we aim to find bright and nearby CQVs, since these objects will be the most amenable to detailed photometric and spectroscopic analyses. To do this, we use 120-second cadence data acquired by TESS between 2018 July and 2022 September (Sectors 1-55; Cycles 1-4). We present our search methods in Section 2, and the properties of the resulting CQV catalog in Section 3. The evolution of many CQVs over a two-year baseline is described in Section 4, including a deep-dive into LP 12-502. We discuss a few implications in Section 5, and conclude in Section 6.

A point on nomenclature. CQVs have been called “transient flux dips”, “persistent flux dips”, “scallop shells”, “batwings”, (Stauffer et al. 2017) “complex rotators”, (Zhan et al. 2019; Günther et al. 2022; Popinchalk et al. 2023), and “complex periodic variables” (Koen 2023). The CQVs should not be conflated with “dippers”, which are classical T Tauri stars with infrared excesses, and which show large-amplitude variability linked to obscuring inner disk structures and accretion hot spots (Cody et al. 2014; Robinson et al. 2021). While a few similarities between CQVs and dippers do exist (see Sections 3.3 and 5.7), their phenomenology and stellar properties are quite different. At the risk of introducing yet another standard, we hope to use a nomenclature that reflects how, when observed over timescales of more than tens of cycles, CQVs are almost but not exactly periodic. They are quasiperiodic. While the three-type classification scheme proposed by Stauffer et al. (2017) may indeed provide some helpful visual distinctions amongst the CQVs, it seems likely that they are all explained by a single underlying phenomenon, and so we opt to refer to them by a single empirically descriptive name. We also considered names that focused on the presence of transient corotating clumps, or more generally on circumstellar plasma tori; for the time being, “CQV” is agnostic to the true underlying phenomenon.

## 2. METHODS

### 2.1. Stellar selection function

To construct our sample, we analyzed the short-cadence data acquired by TESS between 2018 July 25 and 2022 September 1 (Sectors 1-55). Specifically, we used the 120-second cadence light curves produced by the Science Processing and Operations Center at NASA Ames (Jenkins et al. 2016). While the TESS data products from these sectors also included full frame images with cadences of 600 and 1800 seconds, we limited our scope in this study for the sake of simplicity in data handling and some degree of uniformity in the properties of the data itself. In exchange, we sacrificed both completeness and homogeneity of the selection function. While TESS cumulatively observed  $\approx 90\%$  of the sky



**Figure 1. Complex quasiperiodic variables (CQVs):** *Top:* Phase-folded TESS light curves of three CQVs. Each is stacked over one month. Gray circles are raw 2-minute data; black circles bin to 300 points per cycle. Periods in hours are in the bottom right of each panel. Left-to-right, the objects are LP 12-502 (TIC 402980664; Sector 19), TIC 94088626 (Sector 10), and TIC 425933644 (Sector 28). *Bottom:* Cartoon explanations for the phenomenon. The dust clump scenario (lower left) and prominence scenario (lower center) both invoke centrifugally-supported material at the corotation radius. We disfavor the screen scenario (see Section 1).

for at least one lunar month between 2018 July and 2022 September, the 120-second cadence data were collected for only a subset of observable stars during this time due to telemetric constraints (see Fausnaugh et al. 2021).

To assess the completeness of the resulting 120-second cadence data that is the basis of this study, we cross-matched TIC8.2 (Stassun et al. 2019; Paegert et al. 2021) against the Gaia DR2 point-source catalog (Gaia Collaboration et al. 2018). We opted for Gaia DR2 rather than DR3 because the base catalog for TIC8 was Gaia DR2, which facilitated a one-to-one crossmatch using the Gaia source identifiers. This exercise showed that for  $T < 16$  M dwarfs, the TESS 2-minute data are  $\approx 50\%$  complete at  $\approx 50$  pc. At  $< 20$  pc,  $\gtrsim 80\%$  of the  $T < 16$  M dwarfs have at least one sector of short-cadence data; at  $> 100$  pc,  $\lesssim 10\%$  of such M dwarfs have at least one sector of short-cadence data. Armed with this understanding, we then used our cross-match between Gaia DR2 and TIC8 to select our stars of interest, which we defined as stars with 120-second cadence TESS light curves satisfying these four conditions:

$T < 16$	(Bright for TESS)	(1)
$G_{\text{BP}} - G_{\text{RP}} > 1.5$	(Red stars only)	(2)
$M_G > 4$	(Dwarf stars only)	(3)
$d < 150 \text{ pc}$	(Close stars only)	(4)

Here,  $M_G = G + 5\log(\varpi_{\text{as}}) + 5$  is the Gaia  $G$ -band absolute magnitude,  $\varpi_{\text{as}}$  is the parallax in units of arcseconds, and  $d$  is a geometric distance defined by inverting the parallax and ignoring any zero-point correction. The target sample

therefore includes 65,760 M dwarfs and late-K dwarfs, down to  $T < 16$  and out to  $d < 150$  pc.

## 2.2. CQV discovery

Previous methods for finding CQVs have included visually examining stars known to be in young clusters (Rebull et al. 2016; Stauffer et al. 2017), and automatically flagging rapid rotators with a large number of strong Fourier harmonics (Zhan et al. 2019). The latter approach still requires visual vetting, since “stars with many Fourier harmonics” is a designation that includes objects such as eclipsing binaries or multiple stars blended into a single photometric aperture. In this work, we implemented a new search approach based on counting the number of sharp local minima in phase-folded light curves, while also using the previously tested Fourier approach. We applied these two search techniques independently.

### 2.2.1. Counting dips

The dip counting technique aims to count sharp local minima in phase-folded light curves. CQVs will preferably have at least three such minima in order to be distinct from false positives such as synchronized and spotted binaries (“RS CVn” stars).

For our dip-counting pipeline, we began with the PDC\_SAP flux for each sector, removed non-zero quality flags, and normalized the light curve to one by dividing out its median value. We then flattened the light curve using a 5-day sliding median filter, as implemented in `wotan` (Hipke et al. 2019). On the resulting cleaned and flattened light

curve, we ran a periodogram search, opting for the [Stellingwerf \(1978\)](#) phase dispersion minimization (PDM) algorithm implemented in `astrobase` ([Bhatti et al. 2021](#)) due to its shape agnosticism. If a period  $P$  below 2 days was identified, we reran the periodogram at a finer grid to improve the accuracy of the period determination.

Once a star’s period was identified, we binned the phased light curve to 100 points per cycle. To separate “sharp” local minima from smooth but double-dipped spot variability, we then iteratively fit robust penalized splines to the wrapped phase-folded light curve, excluding points more than two standard deviations away from the local continuum ([Hippke et al. 2019](#)). The maximum number of equidistant spline knots per cycle is the parameter in this framework that controlled the meaning of “sharp”—we allowed at most 10 such knots per cycle, though for most stars fewer knots were preferred based on cross-validation using an  $\ell^2$ -norm penalty. An example fit is shown in panel (e) of Figure 2.

We then identified local minima in the resulting residual light curve using the SciPy `find_peaks` utility ([Virtanen et al. 2020](#)), which is based on comparing adjacent values in an array. For a peak to be flagged as significant, we required it to have a width of at least  $0.02P$ , and a height of at least twice the point-to-point RMS. This latter quantity is defined as the 68<sup>th</sup> percentile of the distribution of the residuals from the median value of  $\delta f_i \equiv f_i - f_{i+1}$ , where  $f$  is the flux and  $i$  is an index over time.

To identify local minima near the edges of the phased light curve, which usually would cover phases  $\phi \in [0, 1]$ , we performed the entire procedure over a phase-folded light curve spanning  $\phi \in [-1, 2]$ , by duplicating and concatenating the ordinary phase-folded light curve. The free parameters we adopted throughout this analysis procedure, for instance the maximum number of spline knots per cycle, and how large and wide of a local minimum to consider a “true dip”, were chosen during testing based on their ability to correctly re-identify a large fraction (>90%) of known CQVs, while also being able to consistently reject common false positives such as rapidly rotating spot-induced variability and typical eclipsing binaries.

Overall, for a star to proceed to manual examination, we required that it have a peak PDM period below two days, and that it exhibited at least three sharp local minima (as algorithmically reported) in at least one observed TESS sector.

### 2.2.2. Fourier analysis

We performed an independent search for strongly periodic phenomena using a Fourier-based approach, following [Zhan et al. \(2019\)](#) and [Pribulla et al. \(2023, their Section 1.3\)](#). Starting with the `PDC_SAP` light curves, we normalized each light curve, and then re-binned it into equal width 120-second bins to account for the uneven spacing in the TESS data, as well as the data gap caused by satellite downlink during each sector. We then padded the data to ensure that the light curve had a length that was a power of two, as described by [Zhan et al.](#). Taking the Fourier transform of the corrected, padded light curve using `numpy.fft.fft`, we then searched for

peaks with a significance of over  $12\sigma$  within a set of 500 frequency bins.

If a peak of such significance was found, we generated a “summary sheet” with information about the star, its full and folded light curves, Fourier transform, potential contaminating stars, and information about these contaminating stars. In the sectors for which we performed this analysis, we found that “summary” sheets were generated for  $\approx 10\%$  of the 20 000 120-second targets in a given sector. We then manually reviewed each summary sheet, and visually labelled each star based on its morphology (including e.g., eclipsing binary, CQV, RS CVn, or cataclysmic variable). CQVs that were observed in multiple sectors had their light curves stitched together and analyzed for a more accurate period determination.

### 2.2.3. Manual vetting

We assessed whether the objects found using the dip-counting (Section 2.2.1) and Fourier (Section 2.2.2) techniques were consistent with expectations for CQVs by assembling the data shown in Figure 2. We labeled a star as a “good” CQV if at least one TESS sector showed what we viewed as the unambiguous signatures of the class (short period; at least three dips or else otherwise oddly-shaped dips; relative stability over a timescale of 30 days). We independently noted stars that we thought could be CQVs, but that were more ambiguous.

Broadly speaking, the most common false positives for both the Fourier and dip-counting techniques were eclipsing binaries, spot-induced variability from rapid rotators, and variability from neighboring, off-target stars. Typical false positive rates from our dip-counting pipeline were 5:1, with 368 unique stars flagged, and about 20% being labeled either “good” or “possible” CQVs. The Fourier analysis was not amenable to calculating a similar false-positive rate, because it was implemented as a general variability search; its results are also being used for analyses on topics other than CQVs.

## 2.3. Stellar properties

**Ages**—We estimated the stellar ages by making probabilistic spatial and kinematic associations between the CQVs and known clusters in the solar neighborhood. For most stars in our sample, we did this using `BANYAN Σ` ([Gagné et al. 2018](#)).<sup>1</sup> This algorithm calculates the probability that a given star belongs to one of 27 young clusters (or “associations”) within 150 pc of the Sun, by modeling the clusters as multivariate Gaussians in 3-D position and 3-D velocity space. We used the Gaia DR2 sky positions, proper motions, and distances to calculate the membership probabilities. `BANYAN Σ` in turn analytically marginalizes over the radial velocity dimension. The probabilities returned by this procedure are qualitatively useful, but should be assessed with caution due to the non-Gaussian nature of most groups

<sup>1</sup> [https://github.com/jgagneastro/bayan\\_sigma](https://github.com/jgagneastro/bayan_sigma), git commit 394b486



**Figure 2. Validation plots used to label CQVs.** The complete figure set, with one image per sector for each of the 66 objects in Table 1 is available in the online journal. **For internal collaboration review:** <https://www.dropbox.com/sh/khtwk5a6z0zgrif/AABRG21Ire7VS4BUnaD1hOR8a?dl=0>. Panels are as follows. *a*): Phase-folded light curve; gray points are raw 2-minute data and black points are binned to 200 points per cycle. The adopted period is given in the lower-right corner. *b*): Phase-dispersion minimization (PDM) periodogram. Dotted lines show up to the 10<sup>th</sup> harmonic and subharmonic. *c*): DSS finder chart, with 1- and 2-TESS pixel radius circles displayed for scale. *d*): Cleaned light curve, binned to 20-minute cadence, in Barycentric TESS Julian Date (BTJD). *e*): Phase-folded light curve, binned to 100 points per cycle. The gray line denotes the automated spline-fit to the wrapped phase-folded light curve, and small gray triangles denote automatically identified local minima. *f*): Phase-folded light curve at twice the peak period. *g*): Phase-folded light curve at half the peak period. *h*): Phase-folded time-series within the “background” aperture defined in the SPOC light curves. *i*): Phase-folded flux-weighted centroid in the column direction. *j*): Phase-folded flux-weighted centroid in the row direction. *k*): Gaia DR2 color–absolute magnitude diagram. The gray background denotes stars within 100 pc. *l*): Information from Gaia DR2, TIC8, and the automated dip-counting search pipeline. “Neighbors”, abbreviated “nbhr”, are listed within apparent distances of 2 TESS pixels if  $\Delta T < 2.5$ . *m*): BANYAN- $\Sigma$  v1.2 association probabilities, calculated using positions, proper motions, and the parallax.

within the solar neighborhood (see e.g. Kerr et al. 2021, Figure 10).

For a few cases where BANYAN  $\Sigma$  yielded ambiguous results, we consulted the meta-catalog of young, age-dated, and age-dateable stars within from Bouma et al. (2022), and also searched the local volume around each star for co-moving companions.<sup>2</sup>

<sup>2</sup> <https://github.com/adamkraus/Comove>, git commit 278b372

*Effective temperatures, radii, and masses*—We determined the stellar effective temperature and radii by fitting the broad-band spectral energy distributions (SEDs); we then estimated the masses by interpolating against the sizes, temperatures, and ages of the PARSEC v1.2S models (Bressan et al. 2012; Chen et al. 2014).

For the SED fitting, we used `astroARIADNE` (Vines & Jenkins 2022). We adopted the BT-Settl stellar atmosphere models (Allard et al. 2012) assuming the Asplund et al. (2009) solar abundances, and the Barber et al. (2006) wa-

ter line lists. The broadband magnitudes we considered included  $GG_{BP}G_{RP}$  from Gaia DR2,  $Vgri$  from APASS,  $JHK_S$  from 2MASS, SDSS  $riz$ , and the WISE  $W1$  and  $W2$  passbands. We omitted UV flux measurements from our SED fit to avoid any possible bias induced by chromospheric UV excess; we similarly omitted WISE bands  $W3$  and  $W4$ , but due to reliability concerns. `astroARIADNE` compares the measured broadband flux measurements against pre-computed model grids, and by default fits for six parameters:  $\{T_{\text{eff}}, R_*, A_V, \log g, [\text{Fe}/\text{H}], d\}$ . The distance prior is drawn from Bailer-Jones et al. (2021). The surface gravity and metallicity are generally unconstrained. And finally, given our particular use-case, we assumed the following priors for the temperature, stellar size, and extinction:

$$T_{\text{eff}}/\text{K} \sim \mathcal{N}(3000, 1000), \quad (5)$$

$$R_*/R_\odot \sim \mathcal{T}_N(0.5, 0.3, 0.1, 1.5), \quad (6)$$

$$A_V/\text{mag} \sim \mathcal{U}(0, 0.2), \quad (7)$$

for  $\mathcal{N}$  the Gaussian and  $\mathcal{U}$  the uniform distributions, and  $\mathcal{T}(\mu, \sigma, a, b)$  a truncated normal distribution with mean  $\mu$ , standard deviation  $\sigma$ , and lower and upper bounds  $a$  and  $b$ . We validated our chosen upper bound on  $A_V$  using a 2MASS color-color diagram. Finally, using `Dynesty` (Speagle 2020), we sampled the posterior probability assuming the default Gaussian likelihood, and set a stopping threshold of  $d\log \mathcal{Z} < 0.01$ , where  $\mathcal{Z}$  denotes the evidence.

With the effective temperatures and stellar radii from the SED fit, we then estimated the stellar masses by interpolating against the PARSEC isochrones (v1.2S Chen et al. 2014). The need for models that incorporate some form of correction for young, active M dwarfs is well-documented (e.g. Stassun et al. 2012; David & Hillenbrand 2015; Feiden 2016; Somers et al. 2020). Plausible explanations for anomalous M dwarf colors and sizes relative to model predictions include starspot coverage (e.g. Gully-Santiago et al. 2017), and potentially incomplete line lists (e.g. Rajpurohit et al. 2013). In the PARSEC models, Chen et al. (2014) performed an empirical correction to the temperature–opacity relation drawn from the BT-Settl model atmospheres, in order to match observed masses and radii of young eclipsing binaries. This is sufficient for our goal of estimating stellar masses. Given our observed  $\{\tilde{T}_{\text{eff}}, \tilde{M}_*, \tilde{t}\}$ , and approximating their uncertainties as Gaussian  $\sigma_{\tilde{T}_{\text{eff}}}$ ,  $\sigma_{\tilde{M}_*}$  and  $\sigma_{\tilde{t}}$ , we evaluate a distance  $d$  between our observations and any model PARSEC grid-point  $\{T_{\text{eff}}, M_*, t\}$  as

$$d^2 = \left( \frac{\tilde{T}_{\text{eff}} - T_{\text{eff}}}{\sigma_{\tilde{T}_{\text{eff}}}} \right)^2 + \left( \frac{\tilde{M}_* - M_*}{\sigma_{\tilde{M}_*}} \right)^2 + \left( \frac{\tilde{t} - t}{\sigma_{\tilde{t}}} \right)^2, \quad (8)$$

in order to assign equal importance to each dimension. The preferred model mass is then one that minimizes this distance, and is quoted in Table 1. **Igb todo: real interpolation, rather than nearest neighbor**

### 3. RESULTS

#### 3.1. CQV catalog

Table 1 lists the 66 objects identified by our search. The “high-quality sample” includes 50 objects, which demonstrated what we viewed as unambiguous characteristics of the CQV phenomenon in at least one TESS sector. The classification of 13 CQV candidates was ambiguous, and the 3 remaining objects were false positives that we discuss below. The quality column in the table divides the three classes; additional data from TESS or other instruments could help resolve the classification of the ambiguous cases. Of the 63 CQVs and candidate CQVs, 32 were found using both the dip-counting and Fourier techniques, 23 were found using only the dip-counting technique, and 8 were found using only the Fourier technique. In the following, we will restrict our discussion to the high-quality sample, irrespective of discovery method.

The mosaic in Figure 3 shows phased light curves for the 50 CQVs. The objects are sorted first in order of the number of TESS 120-second cadence sectors in which they clearly demonstrated the CQV phenomenon, and secondarily by descending brightness. The top five objects by this metric are TIC 300651846 ( $T=13.5$ , 12 sectors); TIC 402980664 ( $T=11.1$ , 7 sectors); TIC 89463560 ( $T=13.5$ , 5 sectors); TIC 363963079 ( $T=12.9$ , 5 sectors); and TIC 294328887 ( $T=14.2$ , 4 sectors). The brightest five CQVs span  $9.3 < T < 11.1$ ; the faintest five span  $14.5 < T < 15.0$ . The fastest five have periods spanning  $3.6 \text{ hr} < P < 6.2 \text{ hr}$ , and the slowest five span  $27 \text{ hr} < P < 38 \text{ hr}$ .


In terms of the light curve shapes, Figure 3 shows a broad range of variability, with anywhere from two to eight local minima per cycle. Some stars show relatively ordinary modulation during one portion of the phased light curve, and highly structured modulation in the remainder of the cycle (e.g. TIC 206544316, TIC 224283342, TIC 402980664). Others show structured modulation over the entire span of a cycle (e.g. TIC 2234692, TIC 425933644, TIC 142173958). Others show some mix between these two modes.

A small number of objects at first glance seem reminiscent of eclipsing binaries, such as TIC 193831684, TIC 59836633, or TIC 5714469. We believe these cases are unlikely to be eclipsing binaries due to the additional coherent peaks and troughs in the light curves, which are distinct from any binary phenomena of which we are aware.

#### 3.2. Ages

Of our 63 confirmed and candidate CQVs, 61 were associated with a nearby moving group or open cluster, primarily using BANYAN  $\Sigma$ .<sup>3</sup> The relevant groups are listed in Table 1; their ages span  $\approx 5\text{--}200 \text{ Myr}$ . The most prodigious groups were Sco-Cen, Tuc-Hor, and Columba. Six CQVs were also identified in the Argus association (Zuckerman 2019), which serves as an indirect line of evidence supporting the reality and youth of that group. The yield in

<sup>3</sup> Two of the 61 memberships were made with low confidence and are flagged in Table 1. One membership, of TIC 397791443 in IC 2602, was made manually from a literature search (e.g. Cantat-Gaudin & Anders 2020).



**Figure 3. CQVs found in the TESS 2-minute data.** Phased TESS light curves over one month are shown for 50 CQVs in the high quality sample. Gray are raw 2-minute data; black bins to 300 points per cycle. Objects are ordered such that sources with the most TESS data available are on top (see Section 3.1). Each panel is labeled by the TIC identifier, the TESS sector number, the period in hours, and the three-bit binary flag from Table 1, which denotes Gaia DR3 radial\_velocity\_error outliers (bit 1), Gaia DR3 ruwe outliers (bit 2), and stars with secondary TESS periods (bit 3).

Sco-Cen is not surprising since Sco-Cen contains the majority of pre-main-sequence stars in the solar neighborhood, and many of its stars were selected for TESS 120-second cadence observations by various guest investigators. Given the  $\lesssim 10\%$  completeness of TESS beyond 100 pc, there may be many more CQVs in Sco-Cen that remain to be discovered.

For the two stars for which BANYAN  $\Sigma$  did not find any association, we were not able to confidently associate either star with any young groups. Both do seem to show the CQV signal over multiple TESS sectors, and both are photometrically elevated relative to the main sequence. Both were also noted by Kerr et al. (2021) as being in the “diffuse” population of  $< 50$  Myr stars near the Sun.

Our search confirms that the CQV phenomenon persists for at least  $\approx 150$  Myr. Table 1 includes three  $\approx 150$  Myr CQVs in AB Dor (Bell et al. 2015), a  $\approx 112$  Myr old Pleiades CQV (Dahm 2015), and a similarly-aged Psc-Eri member (Ratzenböck et al. 2020). To our knowledge, TIC 332517282 in AB Dor was the previous record-holder for the oldest-known CQV (Zhan et al. 2019; Günther et al. 2022); at least one unambiguous CQV (EPIC 211070495) and a few other candidates were also previously known in the Pleiades (Rebull et al. 2016).

The upper age limit for CQVs may even pass 200 Myr, based on the candidate membership of TIC 294328887 in the Carina Near moving group (Zuckerman et al. 2006). This group’s  $200 \pm 50$  Myr age is based on the lithium sequence of its G-dwarfs (Zuckerman et al. 2006), which shows a co-eval population of stars older than the Pleiades but younger than the 400 Myr Ursa Major moving group. The formal BANYAN- $\Sigma$  membership probability is however somewhat low, perhaps due to the missing radial velocity. This lack of information could be resolved by acquiring even a medium-resolution spectrum. An independent assessment of the group’s kinematics using Gaia data, and its rotation sequence using TESS, could lend further credence in such an analysis.

### 3.3. Infrared excesses

Most CQVs in our catalog did not show infrared excesses in the  $W1-W4$  bands, which is typical for this class of object (Stauffer et al. 2017). Visually inspecting the SEDs of our 66 star sample and the WISE images available through IRSA, we labeled two objects as having reliable infrared excesses (both  $W3$  and  $W4$  at  $> 3\sigma$  above the photospheric prediction).

The two sources exhibiting an IR excess were TIC 193136669 (TWA 34) and TIC 57830249 (TWA 33). Both are in the TW Hydreae association ( $\approx 10$  Myr), and have relatively long periods of 38 hr and 44 hr respectively. In our initial labeling, we labeled both as “ambiguous” CQVs because the dips in their light curves did not show the rigid periodicity typical of CQVs; their periods were also relatively long. However, inspection of further sectors clarified that both sources are dippers (see online plots in Figure 2). Independently, TIC 193136669 has a cold molecular disk based on observed 1.3 mm continuum emission and resolved Keplerian  $^{12}\text{CO}(2 - 1)$  emission (Rodriguez et al. 2015). It was also labeled a dipper by Capistrant et al. (2022);

we agree with their designation, and label it an “impostor” CQV in Table 1. TIC 57830249 (TWA 33) also has previously detected 1.3 mm continuum emission (Rodriguez et al. 2015), suggestive of cold dust grains being present. It is also a dipper. Section 5.7 highlights plausible evolutionary connections between CQVs and dippers in light of these “misclassifications”.

### 3.4. Binarity

The main types of binaries of interest in this work were those that were unresolved, because they could confuse our understanding of CQVs. For instance, unresolved binaries might produce multiple photometric signals and hinder our ability to correctly identify the star hosting the CQV signal. Unresolved binaries could also bias photometric magnitude and color measurements, which would affect our stellar parameter estimates. To attempt to identify binaries, we considered the following lines of information.


*Radial velocity scatter*—We examined diagrams of the Gaia DR3 “radial velocity error” as a function of stellar color for all 63 CQVs and candidate CQVs. Since this quantity represents the standard deviation of the non-published Gaia RV time series, outliers can suggest single-lined spectroscopic binarity. These plots showed two clusters of stars, at  $\lesssim 10 \text{ km s}^{-1}$  and  $20-25 \text{ km s}^{-1}$ . We there adopted a threshold of  $20 \text{ km s}^{-1}$  to flag possible single-lined spectroscopic binaries, which selected three stars: TIC 405910546, TIC 224283342, and TIC 280945693.

*RUWE*—We examined plots of Gaia DR3 RUWE as a function of color.<sup>4</sup> Elevated RUWEs imply excess noise relative to a single-source astrometric model. This can be caused by marginally resolved point sources skewing the single-transit centroid measurements, intrinsic photometric variability, or intrinsic astrometric motion. Based on this exercise, we adopted a threshold of  $\text{RUWE}_{\text{DR3}} > 2$  to flag sources with excess astrometric noise. This threshold was met for 16/50 high-quality CQVs; 0/13 of the ambiguous ones had this characteristic. There is some subjectivity in where to set the threshold, since the RUWE distribution has an extended tail (e.g. Penoyre et al. 2022). If we had instead required  $\text{RUWE}_{\text{DR3}} > 1.4$ , 21/50 high-quality CQVs and 2/13 of the ambiguous sample would have been flagged.

*Gaia DR3 non-single stars*—Gaia DR3 included a `non_single_star` column that flagged known or suspected eclipsing, astrometric, and spectroscopic binaries. None of the stars in our CQV sample were identified as possible binaries in this column.

*Multiple periodic TESS signals*—During our visual analysis of the TESS light curves and PDM periodograms, we flagged sources with beating light curves, and with PDM periodograms that showed multiple periods. We also attempted to disentangle the two signals by subtracting the

<sup>4</sup> For an explanation of the renormalized unit weight error (RUWE), see the GAIA DPAC technical note [http://www.rssd.esa.int/doc\\_fetch.php?id=3757412](http://www.rssd.esa.int/doc_fetch.php?id=3757412).



**Figure 4. Properties of CQVs identified by our search.** CQVs are mostly pre-main-sequence M dwarfs, younger than  $\approx 150$  Myr, with rotation periods faster than  $\approx 1$  day. The 50 bona fide CQVs in Table 1 are the dark blue circles; 13 ambiguous CQV candidates are light blue circles. Unresolved binaries are flagged either if the Gaia DR3 radial velocity scatter exceeded  $20 \text{ km s}^{-1}$ , or if  $\text{Gaia RUWE}_{\text{DR3}} > 2$  and multiple photometric signals were present in the TESS light curve. The top panels show the 65,760 target stars with 120-second cadence TESS data as the shaded gray background; darker regions correspond to a larger relative number of searched stars. The lower-left panel compares the rotation–color distribution of CQVs against the rotation periods of K and M dwarfs in the Pleiades from Rebull et al. (2016). The lower-middle panel plots the derived corotation radii  $R_{\text{cr}} = (GM/\Omega^2)^{1/3}$  in units of stellar radii against the measured CQV periods, in units of hours. Ages in the final panel are known from cluster membership.

mean CQV signal over each sector, and then repeating the phase-dispersion minimization analysis. The results of this effort are summarized in the secondary periods,  $P_{\text{sec}}$ , listed in Table 1. This process yielded 22/50 high-quality CQVs with secondary periods; 3/13 of the ambiguous sample met the same criterion. Of the 16 good CQVs with  $\text{RUWE}_{\text{DR3}} > 2$ , 15 also showed secondary periods in the TESS light curves. Considering the weaker threshold of  $\text{RUWE}_{\text{DR3}} > 1.4$ , 18/21 such CQVs showed secondary TESS periods.

Table 1 summarizes each of the above sources of information into a single bitwise column. We discuss possible connections between binarity and the CQV phenomenon in Section 5.2.


#### 4. CQV EVOLUTION

##### 4.1. Evolution over two year baseline


Figure 5 shows “before” and “after” views of 27 CQVs for which TESS 120-second cadence observations were available at least two years apart. Such a baseline was available for 32 of the confirmed 50 CQVs in our catalog; for plotting purposes we show the brightest 27. We have phased each sector to its own local minimum because for most of the sources we do not know the period at the precision necessary to be able to accurately propagate an ephemeris over two years. The achievable period precision,  $\sigma_P$ , can be estimated as

$$\sigma_P = \frac{\sigma_\phi P}{N_{\text{baseline}}}, \quad (9)$$

for  $N_{\text{baseline}}$  the number of cycles in the observed baseline and  $\sigma_\phi$  the phase precision with which any one feature (e.g. a dip, or the overall shape of the sinusoidal envelope) can be tracked. Assuming  $\sigma_\phi \approx 0.02$  and a 20-day baseline over a single TESS sector yields  $\sigma_P \approx 0.25^{+0.38}_{-0.14}$  minutes for the pop-



**Figure 5. CQVs keep their periods but change their shapes.** Out of the 50 CQVs in Figure 3, 32 had 120-second cadence TESS data available for a baseline of at least two years; the 27 brightest are shown here due to space constraints. Each panel shows one sector of TESS data, and is phased to its deepest minimum in flux. Each panel's title shows the TIC identifier and approximate period in hours. Text insets show the TESS sector numbers, which generally span two years, or at least 1,000 cycles. The vertical scale is fixed across sectors to clarify shape changes. Gray circles are raw 2-minute data; colored circles bin to 300 points per cycle.



**Figure 6.** LP 12-502 (TIC 402980664) light curve, where each time chunk represents one TESS orbit. Data were acquired in Sectors 18-19, 25-26, 53, and 57-58. Flares are drawn in gray. The light curve is binned to 15-minute intervals so that there are 96 points per day, and each point is connected by a line. Data gaps longer are not interpolated; if data are missing, nothing is plotted. The red vertical lines highlight apparently instantaneous state changes in the shape of the dip pattern.

ulation shown in Figure 5; propagated forward 1,000 cycles yields a typical ephemeris uncertainty range of 2–11 hours. Measuring the periods for each sector independently, there did not seem to be any significant ( $>3\sigma$ ) period changes. This implies an absolute period stability of  $\lesssim 0.1\%$  over the two-year baseline.

A few objects in Figure 5 show clear signs of the CQV phenomenon in one sector, and marginal or non-existent signs in the other. While there is subjectivity in this assessment, to our eyes cases for which at least one sector would be flagged as “ambiguous” include TIC 368129164 (Sector 23 might be labeled an EB), TIC 177309964 (Sector 38 would be simply a rotating star), TIC 404144841 (Sector 38 looks like a rotating star), TIC 201898222 (Sector 3 looks like a rotating star), TIC 144486786 (Sector 32 might be an RS CVn), and TIC 38820496 (Sector 28 might be an RS CVn). TIC 193831684, assessed on a single-sector basis, would probably be labeled an eclipsing binary—in fact, Justesen & Albrecht (2021) already gave this source such a label. However, based on the shape evolution between Sectors 13 and 39, it is a CQV. Based on the fraction of sources overall that “turned off”, the observed shape evolution implies that CQVs have a duty cycle of  $\approx 75\%$ . This type of correction is worth including in population-level estimates of how intrinsically common CQVs are in the low-mass stellar population (e.g. Günther et al. 2022).

#### 4.2. Evolution over adjacent sectors, & LP 12-502

A small fraction of our objects were in regions of sky with many sectors of adjacent TESS data. Out of this already small sample, LP 12-502 (TIC 402980664;  $d=21\text{ pc}$ ,  $J=9.4$ ,  $T=11.1$ ) stood out due to the quality and content of its data. We discuss another interesting source, TIC 300651846, in Appendix A. In this section, we first present the LP 12-502 observations and then highlight their implications.

##### 4.2.1. LP 12-502 observations

Figure 6 shows all available data for LP 12-502 from TESS Sectors 18, 19, 25, 26, 53, 58, and 59, split into successive orbits. The star was observed at 120-second cadence whenever it was observable by TESS. We binned the light curve to 15-minute intervals for visual clarity, and required at least one (120-second cadence) flux measurement per bin. Points more than  $2.5\sigma$  above the median are drawn in gray, also for visual clarity. Missing data are not drawn. Figure 7 then shows the same data, but stacked into successive TESS orbits spanning half a lunar month each. Finally, Figure 8 shows “river plots” of the same data, split into similar intervals: the Sector 18–19 data, 25–26 data, 53 data, and 58–59 data. The river plots are subject to one additional processing step: we fitted and subtracted a maximum-likelihood two-harmonic sinusoid independently from the Sector 18–19 data, 25–26 data, and 53, 58, and 59 data in order to accentuate changes in the dip timing and structure.

The average period, determined by measuring the PDM peak period over each sector independently, was  $\langle P \rangle = 18.5560\text{ hr}$ . The range between the maximum and minimum

sector-specific periods was measured to be about one minute. However, in detail, a period shift of  $\pm 1$  minute would yield major phase drifts over the baseline; that time interval corresponds to  $\approx 1/1000^{\text{th}}$  of a period, and we have observed 1500 cycles. By fine-tuning over a grid in period, we found  $P = 18.5611 \pm 0.0001\text{ hr}$ , which seems to track certain features in the LP 12-502 light curve well over the entire dataset.

What exactly is observed? For the first 64 cycles, the star shows a pattern reminiscent of an eclipsing binary, with four obvious local minima. We dub these dips {1, 2, 3, 4} at phases  $\{-0.28, -0.08, 0, 0.25\}$ , respectively. Over cycles 0–64, the depth of dips 1 and 3 remain roughly fixed. However dip 4 decreases in depth by about 2%, while dip 2 increases in depth, by about the same amount (see Figure 7). During cycles 48–64, a fifth dip may also be emerging, in the main “large” dip group.


There is then a 6-month (184 cycle) gap to Cycles 248–315, which show two highly structured dip complexes, plus a small leading dip. The single leading dip is present at the same phase as in cycles 0–64, and is therefore likely to be the same structure. Along a similar line of logic, it seems plausible that the first “dip complex” during cycles 248–264 represents an evolution of the initial complex seen during cycles 0–64, though with reduced depth. During cycles 266–310, an additional local minimum develops between the two complexes; this feature is best visualized on the river plots (Figure 8), and we describe its (shorter) period below.

The second dip complex during cycles 248–315 shows the most substructure. During e.g. cycles 283–298, this single complex shows six local minima. The deepest dip is sharp: it shows a flux excursion of 3.5% over about 22 minutes ( $0.02 P$ ), which is the steepest slope exhibited anywhere in the LP 12-502 dataset. After the sharp dip, there is a roughly exponential fall-off spanning about a quarter of a period, punctuated by coherent local minima and maxima which in detail (Figure 8) have slightly longer periods than the sharp dip. The sharp leading dip only decreases in depth following a sudden state-switch at BTJD 2030.7 (cycles 299–315), which happens during a flare (Figure 6). The trailing dips remain thereafter.

Sectors 53–58 (cycles 1233–1481) are comparatively tame; they showed only four to six dips per cycle. Some dips remain stable in depth and duration over this five month interval. Other dips grow, like the one at  $\phi = +0.06$  between cycles 1499 and 1481. Other dips, such as the one at  $\phi = +0.12$  in cycles 1233–1264, disappear entirely. The most dramatic state switch occurs during cycles 1233–1247, when a large dip “switches” from a depth of 3% and duration of 3 hours to a depth of 0.3%, and a duration of 1 hour.

##### 4.2.2. Lessons from LP 12-502

STATE-SWITCHES REVEAL DIP INDEPENDENCE—The state-switches seen in cycles 1233–1247 and 299–315 confirm that dips can disappear in less than one cycle, a point which has been previously appreciated (Stauffer et al. 2017). What is new in these particular changes is that the morphology changes show that the dips can be *independent* and *ad-*



**Figure 7. Evolution of LP 12-502 ( $P=18.6$  h)** at fixed period and epoch over three years. Each panel shows one (stacked) TESS orbit; small text denotes relative cycle number. There are 200 binned black points per cycle. The TESS pointing law dictates the large time gaps between Cycles 64-248, 315-1233, and 1264-1410; larger gaps tend to yield larger shape changes. The dips usually evolve over tens to hundreds of cycles. However cycles 1233-1264 show a dip that “switched” from a depth and duration of 3% and 3 hr to 0.3% and 1 hr over less than one cycle (cf. Figure 6).


ditive. For example, throughout cycles 1233-1264, there are three local minima between phases of 0 and 0.3. They all have identical ingress times. The shape change during the transition implies that the leading dip that “turned off” (reduced its depth and duration), while the trailing two dips remained fixed. This is visible in Figure 7. In other words, the structures producing these dips are independent to the degree that one can undergo a severe change while the others remain identical. The state switches during cycles 248-264 and 299-315 share the same characteristic: it is always the *leading* dip of a complex that “switches off”, leaving the (fixed-depth) trailing dips in its wake.

**SLOW GROWTH; RAPID DEATH**—Although there are a few instances in which we observe dips switch off over less than one cycle, dip growth seems to happen more slowly. For instance, the dip that grows between phase 0 and 0.1 between cycles 260-290 begins to become visible around BTJD 1993.2, and grows in depth by about 2% over about six cycles, to become visible by eye by BTJD 1997.7. The evolution of this particular dip is most clear in the river plots.

The evolution of the latter dip group in cycles 1410-1481 is another example of this slow mode of dip growth.

**DIP DURATIONS**—The shortest dip duration for any of the individual LP 12-502 dips seems to be  $\approx 0.06 P \approx 1.08$  hr. In comparison, using the stellar radius and mass derived in Section 2.3, the characteristic timescale  $T_{\text{dur}} \equiv R_* P / (\pi a)$  for the transit of a point-source at corotation is  $1.02 \pm 0.10$  hr. This means that while some of the LP 12-502 dips are sufficiently long to require structures that are extended in orbital azimuth, the durations of other dips are consistent with effective radii for the occulting material  $R_{\text{eff}} \ll R_*$ . This implies  $a/R_* \approx 5.8$  for this material; the analogous timescale at the stellar surface is about six times slower.

**DIP PERIODS**—Most of the LP 12-502 dips repeat with a period of  $P = 18.5611 \pm 0.0001$  hr. However the river plots (Figure 8) reveal that the light curve has specific dips with simultaneous but distinct periods. For instance, in sectors 25-26, the local minimum that develops around cycle 262 has a period faster than the mean period by  $\approx 0.1\%$ , while some of the trailing local minima in the main dip complex have periods slower than the mean period, by  $\approx 0.04\%$ . In addition,



**Figure 8.** River plots of the LP 12-502 light curve, showing (clockwise from top-left) Sectors 18-19, 25-26, 53, and 58-59. A two-harmonic sinusoid has been subtracted to highlight the sharp dips. In Sectors 25-26 (cycles 248-315), periods are visible at the fundamental period of 18.5611 hr, as well as at faster and slower relative periods (e.g.  $\phi \approx 0.3$  and  $\phi \approx 0.1$ ). Similar instances of multiple simultaneous periods are also visible in Sector 53 and 58-59. White chunks denote missing data.

tion to the fundamental period, we were able to identify at least four distinct periods shown by specific dips over the full Sectors 18-59 dataset, including periods at 18.5683, 18.5672, 18.5473, and 18.5145 hr, with a typical measurement uncertainty of  $\approx 0.0002$  hr. If each period corresponds to an individual dust or gas clump, then this implies that multiple distinct clumps can orbit the star simultaneously, at marginally different separations.

## 5. DISCUSSION

### 5.1. Typical and extreme CQVs

Figure 4 shows derived stellar properties for our CQV catalog, and compares the CQVs against both the target star sample (top panels), and against stars in the Pleiades (Rebull et al. 2016). Typical CQV masses span  $0.1\text{--}0.4 M_{\odot}$ , and typical ages span 2–150 Myr. This mass and age range in-

cludes both fully convective stars ( $M_* \lesssim 0.25 M_\odot$ ), and stars with a combination of radiative cores and convective envelopes ( $M_* \gtrsim 0.25 M_\odot$ ; Baraffe & Chabrier 2018). We found no obvious light curve morphology differences between the two classes. In terms of their rotation rates relative to the Pleiades, the CQVs are among the more rapidly rotating half of M dwarfs.

The closest CQV in our catalog is DG CVn (TIC 368129164;  $d=18$  pc), a member of AB Dor. To our knowledge, this work is the first time that it has been noted as a CQV. The three brightest CQVs are DG CVn ( $T=9.3$ ), TIC 405754448 ( $T=9.6$ ), and TIC 167664935 ( $T=10.3$ ). The shortest period belongs to TIC 201789285, at 3.64 hr. The longest period belongs to TIC 405910546, at 37.9 hr. If the latter source turns out to be an eclipsing binary, the next-longest would be TIC 193831684 (31.0 hr).

The lowest mass ( $\approx 0.12 M_\odot$ ) belongs to TIC 267953787. The catalog contains a few other stars with similar mass. Given the small number of sub-stellar mass objects in our target sample, future studies of brown dwarf photometric variability might also yield CQVs, though there could be degeneracies in interpretation with planetary surface features such as clouds and latitudinal bands (e.g. Apai et al. 2021; Vos et al. 2022).

## 5.2. CQVs and binarity

### 5.2.1. Binary statistics

In Section 3.4, we found that a significant fraction of the CQVs show indications of unresolved binarity. Excess noise above the Gaia single-source astrometric model is common (16/50 high-quality CQVs with  $\text{RUWE}_{\text{DR}3} > 2$ ), as is the presence of multiple periods in the TESS light curves (22/50). Elevated astrometric noise is almost always accompanied by multiple detectable TESS periods (15/16 high-quality CQVs). The latter point strongly corroborates the idea that most sources with  $\text{RUWE}_{\text{DR}3} > 2$  are intermediate separation binaries with projected apparent separations below 1'', and projected physical separations  $\lesssim 50$  AU.

The multiplicity rate of M dwarfs near the Sun is  $26.8 \pm 1.4\%$  (Winters et al. 2019). Based on the same study, the peak of the separation distribution decreases from  $\approx 49$  AU for  $0.30\text{--}0.60 M_\odot$  stars, to  $\approx 11$  AU for  $0.15\text{--}0.30 M_\odot$  M dwarfs. The multiplicity fraction in our CQV sample seems either consistent or perhaps marginally higher than the field fraction. This could plausibly be understood in a physical framework in which the presence of intermediate-separation binaries causes early disk dispersal, freeing the star to contract. The CQV phenomenon seems to require rapid rotation; thus if binary systems tend to produce more rapidly rotating stars, we would expect a sample of CQVs to have a larger binary fraction than the field.

### 5.2.2. Do K dwarf CQVs exist?

To date, the only stars reported to show the CQV phenomenon are M dwarfs, with typical stellar masses  $\lesssim 0.3 M_\odot$  (Stauffer et al. 2017; Günther et al. 2022). However the

two most massive CQVs in our sample, TIC 405754448 and TIC 405910546, appear to have masses of  $\approx 0.82 M_\odot$  and  $\approx 0.60 M_\odot$  respectively. The next-highest masses in our sample are  $\approx 0.40 M_\odot$ .

Based on their locations locations in the color–absolute magnitude diagram and membership in LCC, both of the K dwarfs do indeed appear to be high mass. However in detail, both are subject to ambiguities in interpretation. The TIC 405910546 light curve has a unique shape, suggestive of an eclipsing binary. Independently, TIC 405910546 was one of only three CQVs flagged with a Gaia DR3 radial velocity scatter exceeding  $20 \text{ km s}^{-1}$ . Combined, these factors suggest that TIC 405910546 might be a pre-main-sequence eclipsing binary; it should be studied further to clarify this classification. For TIC 405754448, we believe that the source *is* an unresolved binary, because  $\text{RUWE}_{\text{DR}3}=6.8$ , and because the raw light curves in Sectors 11, 37, and 38 show an additional  $\approx 6.5$  day,  $\approx 0.3\%$  amplitude sinusoidal signal suggestive of an unresolved photometric companion. If TIC 405754448 is a K+M binary, then the flux ratio between the primary and secondary would be expected to be  $\approx 10:1$ . It would be challenging for the M dwarf to produce the observed 12.9 hr signal with such a flux deficit, but not impossible. While both of these objects suggest that the CQV phenomenon may extend up in mass to pre-main-sequence K dwarfs, more data are needed to fully substantiate this claim.

### 5.2.3. An astrophysical CQV false positive: TIC 435903839

We originally classified TIC 435903839, with  $\text{RUWE}_{\text{DR}3}=17.7$ , as an “ambiguous” CQV with a 10.8 hr period, because this period minimized the dispersion in the phase-folded light curve. More careful inspection however showed that this source is an impostor: this source is a photometric blend of two ordinary rotating stars with  $P_0=3.60$  hr, and  $P_1=5.41$  hr, with a beat period  $(P_0^{-1}-P_1^{-1})^{-1}$  of 10.8 hr. This is a novel false positive scenario for CQVs: two rapidly-rotating stars with near-integer ratios of rotation periods. The beat between the two rotation signals produces the apparent CQV signal. Such false positives can be excluded through careful accounting of all peaks in a periodogram. For instance, TIC 435903839 shows a peak at 16.27 hr, which is not an integer multiple of the dispersion-minimizing 10.82 hr period.

### 5.2.4. Multiple CQVs in the same system: TIC 425937691 and TIC 142173958

TIC 142173958 and TIC 425937691 both show evidence for two separate CQV signals in their TESS light curves. For TIC 142173958, the signals are at 11.76 hr and 12.84 hr. For TIC 425937691, they are at  $\approx 4.82$  hr and  $\approx 3.22$  hr, near the 3:2 commensurability. Given that both sources have two photometric signals and elevated RUWEs, they are probably unresolved binaries consisting of two independent CQVs. Recent work has shown that the orbits of binaries closer than  $\lesssim 700$  AU tend to be aligned with their planetary systems (e.g. Christian et al. 2022). If we assume that observing CQV

variability similarly requires near edge-on viewing geometries, then we might expect a sufficiently close binary with one CQV to also have another, since the spin axes in sufficiently close binaries would tend to be aligned. In this scenario, the two periods of TIC 425937691 being within  $\lesssim 1\%$  of the 3:2 period commensurability would simply be a coincidence.

### 5.3. CQVs are quasiperiodic

A periodic signal repeats exactly; the CQVs do not (Figure 5). While their periods appear to remain constant to within measurement precision over thousands of cycles, the light curve shapes evolve over 10 to 1,000 cycles. They are therefore *quasiperiodic*. While this observation is consistent with the analyses by Günther et al. (2022) and Popinchalk et al. (2023), it marks a qualitative departure from the “persistent” vs. “transient” flux dip distinction described by Stauffer et al. (2017), since all CQV dips seem to be transient over timescales of more than 1,000 cycles (Figure 5).

With that said, one might expect a truly quasiperiodic process to be able to explore all phase angles with equal weight. LP 12-502, and perhaps other CQVs, seem to have preferred phases. For LP 12-502, all of the dips happen over phases corresponding to only two thirds of the period (Figure 7). The remaining third seems to be “out of limits” for any dipping material. This could be evidence that the stellar magnetic field is strongly asymmetric, and can hold extrinsic material at corotation, but only over two thirds of the equatorial circle. Alternatively, the source of the material (e.g. a planetesimal swarm) might be distributed over an arc of the same angular extent ( $240^\circ$ ). We favor the former explanation, for reasons discussed below.

### 5.4. Dip asymmetries and dust geometries

The asymmetry of a dip can help diagnose the optical depth of the occulting material as a function of orbital phase angle. Sharp leading edges with trailing exponential egresses for instance have been previously seen for transiting exocomets and disintegrating rocky bodies (e.g. Rappaport et al. 2012; Brogi et al. 2012; Vanderburg et al. 2015; Zieba et al. 2019).

Examining Figure 3, it is not obvious whether CQVs as a whole show any preference for sharper ingresses, or sharper egresses. In some cases (e.g. TIC 425933644), the continuum itself is not well-defined, and so the meaning of “ingress” and “egress” are not clear. In others, such as Sector 36 of TIC 89463560, there is a single clear sharp ingress with an exponential egress, which could be fitted using e.g. a model analogous to those used for exocomets (e.g. Zieba et al. 2019). The main quantities of interest in such models would be the exponential decay time- and therefore length-scale, as well as the impact parameter and the inferred transit depth (and its implications for the equivalent “radius” of the transiting cloud). We briefly explored such models, until it became clear that careful modeling of sources such as LP 12-502 merits its own in-depth study. Connections could likely also be made to the toroidal geometries that can be pro-

duced by outflowing atmospheres of transiting planets (e.g. McCann et al. 2019; MacLeod & Oklopčić 2022).

### 5.5. Dust, or gas?

We believe that the most likely scenarios to explain CQVs are either the dust clump scenario or the prominence scenario (Figure 1). Both invoke clumpy material that would be centrifugally supported at the corotation radius. The prominence idea has a longer history, based on analogy with quiescent prominences/filaments observed to exist in the solar corona for up to a few weeks (see Vial & Engvold 2015). In an extrasolar context, spectroscopic detections of transient Balmer- and resonance-line absorption seen for stars such as AB Dor and Speedy Mic (e.g. Collier Cameron & Robinson 1989; Jeffries 1993; Dunstone et al. 2006; Leitzinger et al. 2016) led to the interpretation that the data could best be explained by similar structures: cold, minimally ionized hydrogen clouds or filaments that scatter chromospheric emission from the star to produce the observed spectroscopic line variations (see Collier Cameron & Robinson 1989). The short-term mechanical stability of such gas configurations is theoretically plausible (Ferreira 2000; Waugh & Jardine 2022), and the interpretation of this class of observations seems at least somewhat secure.

A clear link between the dense gas clumps (prominences) that likely exist around rapidly rotating low-mass stars and the CQV phenomenon has yet to be made. A simple visual examination of the TESS light curves for five prominence-hosting systems studied by Jardine & Collier Cameron (2019)—AB Dor, Speedy Mic, LQ Lup, HK Aqr, and V374 Peg—revealed no obvious CQV behavior, though all show differential evolution, and Speedy Mic shows two closely-spaced periods and a strong beat. Spectroscopically observable prominences do not imply photometric CQV-like dips.

The key difference between the prominence and dust clump scenarios is in whether the occulting material of interest is neutral hydrogen, or dust. In the phrasing of the “frozen flux” condition of ideal rigid field magnetohydrodynamics, the tendency of both to become trapped at the corotation radius in the equatorial plane is tied to how of the four relevant forces (gravity, Lorentz, inertial Coriolis, and inertial centrifugal), the Lorentz and Coriolis only act perpendicular to field lines, while gravity and the centrifugal force are in balance at  $R_{\text{cr}}$  (see Townsend & Owocki 2005, Sec. 2). The magnetic field strength is only relevant in this formulation of the system in that we must have  $R_{\text{cr}} < R_{\text{Alfvén}}$  in order for closed loops to exist that can support prominences (e.g. Nakajima 1985; Jardine & Collier Cameron 2019). In detail however, the locations of such magnetic fixed points depends on the star’s magnetic field topology.

#### 5.5.1. Gas absorption microphysics

CQVs show broadband flux variations that can be 1-2× deeper in the blue than in the red (Onitsuka et al. 2017; Bouma et al. 2020; Günther et al. 2022; Koen 2023). For neutral hydrogen to produce this effect in absorption, its opacity must be larger in the blue than in the red. Gray (1992)

suggests that this might be microphysically possible: while bound-bound absorption provides opacity only at narrow resonant lines, the hydrogen opacity due to bound-free absorption is “jagged” (see Gray 1992, Figure 8.5 and Eq. 8.8), such that at temperatures of  $\approx 3,000$  K to  $\approx 10,000$  K the observed chromaticities might be reproduced. While bound-free absorption of H $^{-}$  seems like it should be important at such cool temperatures, in detail this opacity source has the *opposite sign* from what is required to produce deeper dips in the blue than in the red. Thomson scattering is similarly ruled out as a relevant opacity source, because it is gray. The final plausible alternative opacity source is dust, which has the requisite larger absorption cross-section in the blue than the red (Cardelli et al. 1989).

An instructive point of comparison is the rapidly rotating magnetic B star,  $\sigma$  Ori E, which shows dips that are deeper in the blue than in the red (Hesser et al. 1977). Photometric and spectroscopic observations of this star have been understood in terms of a warped torus of circumstellar material, analogous to the geometries we are discussing for much cooler M dwarfs (Landstreet & Borra 1978; Nakajima 1985; Townsend et al. 2005). This material is unlikely to be dust, due to the relevant sublimation timescales. The opacity is instead thought to be sourced from bound-free absorption from neutral hydrogen (Nakajima 1985), although to our best knowledge direct evidence for this conclusion has yet to be acquired. Separate and smaller-amplitude emission in  $\sigma$  Ori E may also come from electrons scattering photospheric light toward the observer when the clouds are not in transit (Berry et al. 2022).

Given these complexities, it seems important for a future theoretical study to be conducted to determine to what degree the observed chromaticities in CQVs match, or do not match, expectations from radiative transfer. This issue has a key ability to resolve the question of whether the CQVs are explained by dust or by gas, which has bearing on whether the material producing the dips is coming from the star, or whether it is a byproduct of the protoplanetary disk.

### 5.5.2. The lifetime constraint

Independent of the microphysical basis for gas absorption, the observed lifetime of the CQV phenomenon could provide another dimension to discern between the gas vs. dust clump scenarios. Based on the available statistics from e.g. Rebull et al. (2022) and references therein, it seems plausible that CQV occurrence decreases in time, bottoming to zero well before the  $\approx 700$  Myr age of Praesepe. This is odd in the context of the prominence scenario, because pre-main-sequence M dwarfs spin up over the first  $\approx 10^8$  yr; prominences might even be expected to be *more* common at the age of the Pleiades than for younger stars. This broadly assumes that any star that can support prominences will do so, and that the magnetic topology of rapidly rotating M dwarfs at 100 Myr is similar to that of rapidly rotating M dwarfs at say, 10 Myr. Based on the degree to which M dwarfs remain rapid rotators over the first few gigayears of their lives (e.g. Rebull et al. 2022; Pass et al. 2022), it seems surprising in

the gas clump (prominence) scenario that very few stars older than 150 Myr have been seen to exhibit the phenomenon. In the dust clump scenario this is not a problem, because if the dust were externally sourced, it would be expected to have a finite supply.

### 5.6. Planets or planetesimal swarms near corotation?

Close-in planets are common around M-dwarfs; studies from Kepler have shown that high-mass M dwarfs have  $\approx 0.1$  planets per star with sizes between  $1\text{--}4 R_{\oplus}$  and orbital periods within 3 days (Dressing & Charbonneau 2015). The number increases to  $\approx 0.7$  planets per star considering planets out to  $P < 10$  days. Extrapolating to all small close-in planets with say  $0.1\text{--}4 R_{\oplus}$  and within 10 days, it is reasonable to expect on average one close-in planet per M dwarf.

In the context of planet formation, the stopping location for the inner-most planet is set by the location of the protoplanetary disk’s truncation radius (Izidoro & Raymond 2018). Although this disk truncation radius is defined by equating the magnetic pressure from the stellar magnetosphere with the ram pressure from the inflowing gas, it also tends to coincide with the Keplerian corotation radius for low accretion rates (Hartmann et al. 2016; Li et al. 2022). Within models that have compact multiplanet resonant chains migrating inward due to gas drag, the inner-most planets therefore arrive at  $\approx 5\text{--}10$  stellar radii before the disk becomes depleted.

Given this context, it is tempting to try to interpret features of the CQV light curves in terms of the possible presence of either close-in exoplanets or close-in planetesimals. Rocky planets this young could have molten global magma oceans analogous to those that existed on the Earth and Moon (see Lichtenberg et al. 2022), and thus might be undergoing significant outgassing and atmospheric escape. The system would then be analogous to the Jupiter-Io plasma torus (e.g. Neubauer 1980), although with magnetic fields stronger by a factor of  $\approx 1,000$ . However we emphasize that while this type of configuration seems *a priori* plausible, no direct evidence currently supports it.

In systems like LP 12-502, the observed variability would perhaps drive one toward a picture of a disintegrating planetesimal swarm, rather than just a single planetesimal. The need for more than just one launching body is because we see dips appear and disappear at many different phases, but with roughly the same period. The observed changing dip depths in response to flares would be understood through the planetesimal outflow rates being stalled by changes in the local magnetic field or coronal plasma density. The sizes of these purported planetesimals would need to be  $\lesssim 1 R_{\oplus}$ , based on the non-detections of their transits, analogous to e.g. K2-22 (Sanchis-Ojeda et al. 2015) or KOI-2700 (Rappaport et al. 2014). Typical sizes of bodies in the asteroid belt for instance span hundreds of meters to a few kilometers.

There are at least three problems with this disintegrating planetesimal swarm concept. First, the model would predict that certain orbital phases would produce recurrent dips if observed over sufficiently long baselines, because the launching planetesimal would be massive enough to remain in orbit,

while stochastically ejecting material. An analogous system would be K2-22 (Sanchis-Ojeda et al. 2015). For most CQVs (Figure 5), the data are in tension with this expectation. A second issue is that many of the dips show asymmetries in the wrong direction relative to the naive expectation of a trailing comet tail. Invoking non-exponentially decaying dust distributions as a function of azimuth might be one way out, but such an idea lacks firm theoretical footing. Finally, dynamical stability is a concern: in a model in which any new dip is “explained” by invoking a new planetesimal, one might eventually pack the corotation radius to a degree where instability over short timescales would be guaranteed.

It is certainly possible that exoplanets or exoplanetessimals could end up being connected to the CQV phenomenon, for instance as a possible sources for the occulting dust or gas. However for the time being, none of the CQVs that we have studied in this work show clear evidence for the nearby planets that we statistically expect should be common.

### 5.7. From dippers to debris disks

In identifying the two candidate CQVs with outlying SEDs (TICs 193136669 and TIC 57830249; Section 3.3), we were prompted to reconsider our light curve-based labeling, and ultimately concluded that these sources are dippers. This episode suggests that there could be overlap between CQVs and dippers. It is also worth emphasizing that our labeling of e.g. TIC 57830249 was based on a single sector of TESS data (Sector 36) when its behavior was relatively periodic and it showed dip depths of a few percent. However, in other TESS sectors (e.g. Sector 10), this source looks completely different, varying in apparent flux by a factor of two, with no discernible periodicity at all.

Assessing these results against the backdrop of our increasing understanding of dippers (e.g. Cody et al. 2014; Ansdell et al. 2016; Robinson et al. 2021; Capistrant et al. 2022; Gaidos et al. 2022), it is clear that the loss of an infrared excess is associated with strong changes in a star’s optical variability. It is reasonable to imagine connections between CQVs and dippers: both classes of object can show transient flux dips that are relatively narrow in duration. The dips in both are probably associated with clumps of dust or gas. However the CQV dips are both more periodic and less deep than those of dippers; they also display far less transience over timescales of a few to tens of cycles. This is presumably because CQVs have demonstrably less dust than (most) dipper stars. At a population level, the CQV stars are also older. A common mystery between the CQVs and dippers, in our own estimation, is how exactly the *narrowness* of the dust clumps is produced. It is not unreasonable to imagine a similar mechanism operating for both types of object, tied perhaps to a shared magnetic topology, or perhaps to a preference for dust to inspiral to the star in clumped structures.

### 5.8. Are half-cycles important?

The interval of half a cycle period could be significant in the context of CQVs for two reasons. The first is that for material on a circular orbit viewed edge-on, a half-cycle corre-

sponds to the interval between transit and secondary eclipse. The second is that the half-cycle also corresponds to the interval over which half of the star’s surface is visible. Of the CQVs in Figures 3 and 5, a fraction that seems greater than random might exhibit a preference for showing dips or peaks that correspond to the half-cycle interval.

One set of CQVs shows “CQV behavior” (some form of dip complex), but which only lasts for half of any given cycle. TIC 206544316 (Sector 2) is a canonical example; TIC 405910546 (Sector 38), TIC 167664935 (Sector 38), TIC 146539195 (Sector 5), TIC 118449916 (Sector 44), and TIC 312410638 (Sector 38) seem to show the same tendency.


An independent set of CQVs exhibits small dips roughly half a cycle after large dips; it is tempting to label the small dips secondary eclipses. Such sources include TIC 402980664 (Sector 25; relative to the sharpest, deepest minimum), TIC 89463560 (Sector 36), TIC 224283342 (Sector 2), and TIC 442571495 (Sector 12). For the particular case of TIC 224283342, this observation, coupled with Gaia DR3 radial velocities suggesting a radial velocity scatter exceeding  $20 \text{ km s}^{-1}$ , suggests that the system could be a low-mass eclipsing binary transiting a giant polar spot, analogous to TOI-3884 (Almenara et al. 2022).

In general however, it is challenging to interpret the significance of the potential “half-period regularities”; the CQVs exhibit such a wide range of variability that for seemingly any regularity one might posit, it is easy to come up with counter-example objects which do not follow the trend. In other words, these half-cycle CQVs could simply be a product of the human tendency to pattern match.

### 5.9. Mass flux estimate

We can estimate the mass of a transiting cloud by first converting the transit depth to an effective cloud radius,  $R_{\text{cloud}}$ . For most CQVs in Figure 3, this yields  $\approx 2\text{-}20 R_{\oplus}$ . A minimum constraint on the number density follows by requiring the cloud to be optically thick. For cases like LP 12-502, this is reasonable because the transit duration of the shortest dips implies  $R_{\text{cloud}} \ll R_{\star}$ . Carrying out the relevant calculation assuming the occulting material is dust grains  $1 \mu\text{m}$  in size, Sanderson et al. (2023) reported minimum cloud masses of order  $10^{12} \text{ kg}$  (their Eq. 23), which scale linearly with both the optical depth and dust grain radius. This is comparable to a small asteroid; the asteroid belt itself has a mass of order  $\approx 10^{21} \text{ kg}$  (Park et al. 2019). If the material were cool gas prominences rather than dust, the mass would need to be of order  $10^{14} \text{ kg}$  (Collier Cameron et al. 1990), about  $100\times$  larger than the requisite dust mass.

Our observations provide a direct measurement of how often dips appear and disappear, both due to sudden state-switches, and due to more gradual, secular evolution. For instance, LP 12-502 showed three “state-switch” events over the six months of available TESS observations, during cycles 248–264, 299–315, and 1233–1247. In each case, a dip “turned off”. It is plausible to imagine that these events correspond to either mass being ejected from the corotating clumps, or else perhaps being accreted onto the star. In either



**Figure 9.** The magnetic B star connection.  $\sigma$  Ori E and HD 345439 (top row) are magnetic B stars with predominantly dipolar magnetic fields known to host circumstellar plasma tori. HD 37776 and HD 64740 (middle row) are analogous magnetic B stars with field topologies dominated by high order multipoles. The bottom row compares the latter systems against the “best-matching” CQV light curves, selected by eye from Figure 3. CQVs have light curves that are visually similar to the topologically complex magnetic B stars. Stellar masses rounded to one significant figure are given in the lower right of each panel; the star, TESS sector, and period are listed in each subtitle.

case, the corresponding  $\dot{M} \equiv M \cdot dN/dt$  time-averaged over six months is  $\approx 3 \times 10^{-18} M_{\odot} \text{ yr}^{-1} \approx 1 \times 10^{-12} M_{\oplus} \text{ yr}^{-1}$ . Considered cumulatively over the  $\approx 10^8$  years for which the CQV phenomenon is observed, this yields a cumulative moved dust mass of  $10^{-4} M_{\oplus}$ , of order the Solar System’s asteroid belt. If the occulting material is gas, the masses involved would be of order 100 times larger. For cases in which we observe the growth of dips, such as the Sector 29 data for TIC 224283342, or Sector 5 of TIC 294328885, the dip depths typically increase by of order a few percent over ten to twenty days. This growth rate yields a mass flux one order of magnitude larger than the earlier estimate.

#### 5.10. Strengthening the magnetic B star connection

Stauffer et al. (2017) previously noted a possible connection between the CQVs and rapidly rotating magnetic B stars such as  $\sigma$  Ori E, which can have circumstellar gas clouds trapped in corotation (Townsend et al. 2005). The  $\sigma$  Ori class is distinct from Be-star decretion disks, which do not have any obvious connection to the stellar magnetic field (Rivinius et al. 2013).

An argument against the connection between CQVs and the  $\sigma$  Ori E analogs is that the light curve of  $\sigma$  Ori E is much simpler than those in Figure 3, with only two broad local minima, and one “hump” (Figure 9; see also Jayaraman et al. 2022). Within the model proposed by Townsend et al., this is explained by the simplicity of the star’s magnetic field, which is well-approximated by a dipole, similar to most other magnetic B stars (Aurière et al. 2007; Donati & Landstreet 2009). The magnetic axis needs to be tilted relative to the stellar spin axis in order to match the qualitative behavior of both the broadband light curves, and the line-profile variations seen in hydrogen, helium, and carbon (Oksala et al. 2012).

Two exceptions to the rule of “simple field topologies for magnetic B stars” are HD 37776 and HD 64740, which are B stars that were previously known from spectropolarimetry to have field geometries dominated by high order multipoles (Kochukhov et al. 2011). Reviewing the literature, we learned that recent TESS light curves of these two B stars appear surprisingly similar to the CQV light curves (Mikulášek et al. 2020). The middle row of Figure 9 shows the phased TESS light curves for these two outlying magnetic B stars, with by-eye “best matching” CQVs shown for comparison. The number of dips per cycle, the shapes of the dips, and the dip depths relative to the sinusoidal envelope all appear to be similar. The implication of this surprising connection is that the highly-structured M dwarf light curves may imply multipolar magnetic fields – since it is the complex fields which are the defining characteristic of these two B stars, relative to other known magnetic B stars like  $\sigma$  Ori E. This is consistent with the non-axisymmetric topologies that have been reported in a small number of M dwarfs for which Zeeman Doppler Imaging is technically feasible (see the review by Kochukhov 2021).

The physical similarity between the B stars and the M dwarfs presumably has its origin in the existence of a “centrifugal magnetosphere” (see Petit et al. 2013). In other

words, both classes of objects probably satisfy the condition  $R_A > R_{\text{cr}}$ , for  $R_A$  the Alfvén radius. When this condition holds, stable points exist wherever the outward centrifugal force is balanced by the combined effect of gravity and magnetic tension along closed field loops (Nakajima 1985). In the converse “dynamical” case, when  $R_A < R_{\text{cr}}$ , any material launched by the star returns to its surface over the free-fall timescale. A simple order of magnitude estimate assuming a dipole field with  $B_0 \approx 1 \text{ kG}$  at the star’s surface, a local plasma number density  $n \approx 10^9 \text{ cm}^{-3}$ , and a plasma temperature  $10^6 \text{ K}$  gives Alfvén radii of order a few times the corotation radii,  $R_{\text{cr}}$ . This suggests that the existence of a centrifugal magnetosphere is plausible in young, rapidly rotating M dwarfs.

## 6. CONCLUSIONS

In this work, we searched 120-second cadence TESS data collected from 2018 July to 2022 September for complex quasiperiodic variables (CQVs). The target stars were 65,760 K and M dwarfs within 150 pc. The selection function was  $>80\%$  complete within 30 pc, and  $<10\%$  complete at distances exceeding 100 pc.

In our target sample, we found 50 objects that showed complex quasiperiodic behavior over at least one TESS sector. These 50 bona fide CQVs are listed in Table 1. This table also includes an additional 13 ambiguous CQVs, whose designation is less certain, and 3 impostors. We inferred ages for all but two of the 66 objects based on association memberships; we also derived temperatures and radii using SED fitting, and then inferred stellar masses by interpolating against stellar evolutionary models. We caution that our selection function was not volume-limited: the TESS 120-second stellar sample had a heterogeneous selection function which may have been biased in favor of young stars over field stars. Previous work however has shown that  $\approx 1\text{-}3\%$  of M dwarfs younger than  $\approx 100 \text{ Myr}$  show the CQV phenomenon (Rebull et al. 2016; Günther et al. 2022).

Analyzing the TESS light curves and stellar properties of our CQVs, we draw the following conclusions.

1. CQVs are quasiperiodic. The mean periods remain fixed over the  $>1,000$ -cycle baseline of available observations; but the light curve shapes always evolve (Figure 5).
2. The same CQV can show multiple periods simultaneously. LP 12-502, for instance, showed dips with four distinct periods within  $\pm 0.3\%$  of its fundamental period, sometimes simultaneously, and each lasting for up to 50 cycles (Figure 8).
3. CQVs evolve over timescales that are both secular ( $>100$  cycles) and impulsive ( $<1$  cycle). Dip growth seems to happen over durations of at least ten cycles, and slow dip decay can also occur. “State-switches” correspond to dips to collapsing instantaneously, and are almost always linked with observed optical flares. This suggests that the occulting material is sensitive to sudden changes in the magnetic field.

4. The rate of dip evolution can be used to place a model-dependent constraint on how much material is either being accreted or ejected during the state changes (Section 5.9). Order of magnitude estimates require at least an asteroid belt’s worth of dust ( $10^{-4} M_{\oplus}$ ) over  $10^8$  years, or  $\approx 10^{-2} M_{\oplus}$  if the occulting material is gas.
5. The CQV phenomenon persists for  $\gtrsim 150 \text{ Myr}$ , based on the existence of multiple CQVs in AB Dor, the Pleiades, and Psc-Eri (Section 2.3). It may even extend to 200 Myr, based on the one CQV we found in the Carina Near moving group (TIC 294328887;  $\approx 200 \text{ Myr}$ ). However the lack of detected CQVs in the Hyades and Praesepe suggests that the lifetime of the phenomenon is limited to the first few hundred million years.
6. The duty cycle for CQVs seems to be  $\approx 75\%$ , based on the fraction of bona fide CQVs that either turned on or turned off during TESS re-observations, two years after the initial observation.
7. Most CQVs are M dwarfs with masses  $0.1\text{-}0.5 M_{\odot}$ . Two sources, TIC 405754448 and TIC 405910546, have masses that appear to exceed the M dwarf limit. However both are potentially binaries, and this may confuse our ability to accurately identify the source of the CQV signal (Section 5.2.2). We encourage additional scrutiny of these objects in future work.
8. Surprising analogs to the CQVs exist in magnetic B stars, but only those with multipolar field topologies. Since most magnetic B stars have dipolar magnetic fields, this suggests that the CQV dips and warps are similarly being sculpted by the stellar magnetic fields, and that the magnetic fields themselves are probably also multipolar.
9. The closest CQVs to the Sun are at distances of 15–20 pc, and the brightest have  $V \approx 12$  ( $J \approx 7.5$ ). We have found most of them in this work, since our CQV sample was  $\gtrsim 80\%$  complete within 30 pc. The lack of CQVs in the volume-complete  $<15 \text{ pc}$  sample of  $0.1\text{-}0.3 M_{\odot}$  stars analyzed by Winters et al. (2021) is consistent with this estimate. Expanding our analysis of the TESS data to the full frame images would yield a truly volume-limited selection function, and would expand the CQV census by about a factor of two within 50 pc, and by a factor of ten within 100 pc.

While many questions remain, two in particular will be important for clarifying what these objects might teach us in a broader astrophysical context: 1) Is the eclipsing material responsible for the phenomenon gas or dust? 2) What sets the characteristic clumping size for the circumstellar material?

The distinction between gas or dust is important because it will clarify whether the CQV phenomenon is intrinsic, so that material comes from the star, or extrinsic, so that it is sourced through some generic evolutionary phase of debris disks. This knowledge would in turn propagate to our understanding of whether the phenomenon is primarily teaching

us about dust production and processing in gas-poor disks, or whether it is teaching us about the ability of cold gas to remain stable in hot stellar coronae for long durations. Observationally, acquisition of medium- or high-resolution time-series spectra holds a good chance at resolving the gas vs. dust question. Given our observed  $\approx 75\%$  duty cycles, such data must be acquired simultaneously with photometric time-series observations (e.g. during TESS re-observation) in order for detections and non-detections to be interpretable.

In both the gas and dust scenarios, CQVs are preferentially viewed edge-on. This implies that after correcting for the line-of-sight inclination, roughly one third of low mass stars (those that rotate rapidly enough [Günther et al. 2022](#)) could trap circumstellar material in the same way. It also suggests that CQVs may preferentially show transiting planets at larger distances than the corotating material, though this may be dependent on whether the magnetic and stellar spin axes tend to be aligned. Given these points, observational follow-up work should include searching for outer transiting planets, and measuring equatorial velocities in order to test whether the stellar inclination angles are indeed preferentially edge-on. Any source of empirical information on the stellar magnetic field, whether from the Zeeman effect (e.g. [Kochukhov 2021](#)) or perhaps radio emission (e.g. [Hallinan et al. 2015](#)), could also help clarify the strength of the magnetospheres for these objects.

On the theoretical front, building a physical understanding what sets the characteristic size scale of the clumping material would help clarify why the light curves have the bizarre shapes that are observed. The relevant puzzles in plasma physics and radiative transfer could perhaps be connected to our understanding of the close-in rocky planets that are expected to be present around most of these stars. The challenges intrinsic to both the observational and theoretical work seem worth the effort.

## ACKNOWLEDGMENTS

LGB gratefully acknowledges support from the Heising-Simons 51 Pegasi b Fellowship. A few thank-yous in particular (from LGB) are owed to G. Laughlin, A. Mann, and J. Spake for clarifications and ideas, and to E. Schaller and the 51 Peg science team for organizing the annual summits. In addition, this paper relied on data collected by the TESS mission, which are publicly available from the Mikulski Archive for Space Telescopes (MAST). Funding for the TESS mission is provided by NASA's Science Mission Directorate. TIC 402980664 in particular was observed at 120-second cadence thanks to the TESS Guest Investigator programs G022252 (PI: J. Schlieder; Sectors 18, 19, 25, 26) and G04168 (PI: R. Jayaraman; Sector 53).

LGB conceived the project and performed the dip-counting search, light curve classification, cluster membership, SED, variability, and secondary-period analyses. RJ and SR performed the Fourier-based analysis and contributed to light curve classification. LR cross-examined the light curve clas-

sification, and contributed an independent SED analysis. AD identified the magnetic B star connection. LAH contributed to project design and to the interpretation of the light curves. GAB acquired and maintained the servers used to run the dip-finding pipeline. GRR is an architect of the TESS mission. **JNW? MNG?** All authors assisted in manuscript writing and revision.

**Software:** astrobase ([Bhatti et al. 2021](#)), astropy ([Astropy Collaboration et al. 2013, 2018, 2022](#)), lightkurve ([Lightkurve Collaboration et al. 2018](#)), numpy ([Harris et al. 2020](#)), pyGAM ([Servén & Brummitt 2018](#)), scipy ([Virtanen et al. 2020](#)), TESS-point ([Burke et al. 2020](#)), wotan ([Hippke et al. 2019](#)).

**Facilities:** Astrometry: Gaia ([Gaia Collaboration et al. 2018, 2022](#)). Imaging: Second Generation Digitized Sky Survey. Spectroscopy: Keck:I (HIRES; [Vogt et al. 1994](#)). Photometry: TESS ([Ricker et al. 2015](#)), Broadband photometry: 2MASS ([Skrutskie et al. 2006](#)), APASS ([Henden et al. 2016](#)), Gaia ([Gaia Collaboration et al. 2018, 2022](#)), SDSS ([York et al. 2000](#)), WISE ([Wright et al. 2010](#)).

## REFERENCES

- Allard, F., Homeier, D., & Freytag, B. 2012, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 370, 2765
- Almenara, J. M., Bonfils, X., Forveille, T., et al. 2022, *A&A*, 667, L11
- Ansdell, M., Gaidos, E., Rappaport, S. A., et al. 2016, *ApJ*, 816, 69
- Apai, D., Nardiello, D., & Bedin, L. R. 2021, *ApJ*, 906, 64
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, *ARA&A*, 47, 481
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, 156, 123
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, *ApJ*, 935, 167
- Aurière, M., Wade, G. A., Silvester, J., et al. 2007, *A&A*, 475, 1053
- Bailer-Jones, C. A. L., Rybizki, J., Fouesneau, M., Demleitner, M., & Andrae, R. 2021, *AJ*, 161, 147
- Baraffe, I., & Chabrier, G. 2018, *A&A*, 619, A177
- Barber, R. J., Tennyson, J., Harris, G. J., & Tolchenov, R. N. 2006, *MNRAS*, 368, 1087
- Basri, G. 2021, An Introduction to Stellar Magnetic Activity
- Bell, C. P. M., Mamajek, E. E., & Naylor, T. 2015, *MNRAS*, 454, 593
- Berry, I. D., Owocki, S. P., Shultz, M. E., & ud-Doula, A. 2022, *MNRAS*, 511, 4815
- Bhatti, W., Bouma, L., Joshua, et al. 2021, waqasbhatti/astrobase: astrobase v0.5.3, Zenodo
- Bodman, E. H. L., Quillen, A. C., Ansdell, M., et al. 2017, *MNRAS*, 470, 202
- Bouma, L. G., Winn, J. N., Ricker, G. R., et al. 2020, *AJ*, 160, 86
- Bouma, L. G., Curtis, J. L., Masuda, K., et al. 2022, *AJ*, 163, 121
- Bressan, A., Marigo, P., Girardi, L., et al. 2012, *MNRAS*, 427, 127
- Brogi, M., Keller, C. U., de Juan Ovelar, M., et al. 2012, *A&A*, 545, L5
- Burke, C. J., Levine, A., Fausnaugh, M., et al. 2020, TESS-Point: High precision TESS pointing tool, Astrophysics Source Code Library, record ascl:2003.001
- Cantat-Gaudin, T., & Anders, F. 2020, *A&A*, 633, A99
- Capistrant, B. K., Soares-Furtado, M., Vanderburg, A., et al. 2022, *ApJS*, 263, 14
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Chen, Y., Girardi, L., Bressan, A., et al. 2014, *MNRAS*, 444, 2525
- Christian, S., Vanderburg, A., Becker, J., et al. 2022, *AJ*, 163, 207
- Cody, A. M., Stauffer, J., Baglin, A., et al. 2014, *AJ*, 147, 82
- Collier Cameron, A. 1999, in Astronomical Society of the Pacific Conference Series, Vol. 158, Solar and Stellar Activity: Similarities and Differences, ed. C. J. Butler & J. G. Doyle, 146
- Collier Cameron, A., Duncan, D. K., Ehrenfreund, P., et al. 1990, *MNRAS*, 247, 415
- Collier Cameron, A., & Robinson, R. D. 1989, *MNRAS*, 238, 657
- Curtis, J. L., Agüeros, M. A., Mamajek, E. E., Wright, J. T., & Cummings, J. D. 2019, *AJ*, 158, 77
- Dahn, S. E. 2015, *ApJ*, 813, 108
- David, T. J., & Hillenbrand, L. A. 2015, *ApJ*, 804, 146
- Donati, J. F., & Landstreet, J. D. 2009, *ARA&A*, 47, 333
- Dressing, C. D., & Charbonneau, D. 2015, *ApJ*, 807, 45
- Dunstone, N. J., Barnes, J. R., Collier Cameron, A., & Jardine, M. 2006, *MNRAS*, 365, 530
- Fausnaugh, M., Morgan, E., Vanderspek, R., et al. 2021, *PASP*, 133, 095002
- Feiden, G. A. 2016, *A&A*, 593, A99
- Ferreira, J. M. 2000, *MNRAS*, 316, 647
- Gagné, J., Mamajek, E. E., Malo, L., et al. 2018, *ApJ*, 856, 23
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *A&A*, 616, A1
- Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2022, *arXiv e-prints*, arXiv:2208.00211
- Gaidos, E., Mann, A. W., Rojas-Ayala, B., et al. 2022, *MNRAS*, 514, 1386
- Gray, D. F. 1992, The observation and analysis of stellar photospheres., Vol. 20
- Gully-Santiago, M. A., Herczeg, G. J., Czekala, I., et al. 2017, *ApJ*, 836, 200
- Günther, M. N., Berardo, D. A., Ducrot, E., et al. 2022, *AJ*, 163, 144
- Hallinan, G., Littlefair, S. P., Cotter, G., et al. 2015, *Nature*, 523, 568
- Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, *Nature*, 585, 357
- Hartmann, L., Herczeg, G., & Calvet, N. 2016, *ARA&A*, 54, 135
- Henden, A. A., Templeton, M., Terrell, D., et al. 2016, *VizieR Online Data Catalog*, II/336
- Hesser, J. E., Ugarte, P. P., & Moreno, H. 1977, *ApJL*, 216, L31
- Hippke, M., David, T. J., Mulders, G. D., & Heller, R. 2019, *arXiv:1906.00966 [astro-ph]*, arXiv: 1906.00966
- Hippke, M., David, T. J., Mulders, G. D., & Heller, R. 2019, *AJ*, 158, 143
- Izidoro, A., & Raymond, S. N. 2018, in *Handbook of Exoplanets*, ed. H. J. Deeg & J. A. Belmonte, 142
- Jardine, M., & Collier Cameron, A. 2019, *MNRAS*, 482, 2853
- Jayaraman, R., Hubrig, S., Holdsworth, D. L., et al. 2022, *ApJL*, 924, L10
- Jeffries, R. D. 1993, *MNRAS*, 262, 369

- Jenkins, J. M., Twicken, J. D., McCauliff, S., et al. 2016, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 9913, Software and Cyberinfrastructure for Astronomy IV, ed. G. Chiozzi & J. C. Guzman, 99133E
- Johns-Krull, C. M., Prato, L., McLane, J. N., et al. 2016, *ApJ*, 830, 15
- Justesen, A. B., & Albrecht, S. 2021, *ApJ*, 912, 123
- Kenyon, S. J., & Hartmann, L. 1995, *ApJS*, 101, 117
- Kerr, R. M. P., Rizzuto, A. C., Kraus, A. L., & Offner, S. S. R. 2021, *ApJ*, 917, 23
- Kochukhov, O. 2021, *A&A Rv*, 29, 1
- Kochukhov, O., Lundin, A., Romanyuk, I., & Kudryavtsev, D. 2011, *ApJ*, 726, 24
- Koen, C. 2021, *MNRAS*, 500, 1366
- . 2023, *MNRAS*, 518, 2921
- Landstreet, J. D., & Borra, E. F. 1978, *ApJL*, 224, L5
- Leitzinger, M., Odert, P., Zaqrashvili, T. V., et al. 2016, *MNRAS*, 463, 965
- Li, R., Chen, Y.-X., & Lin, D. N. C. 2022, *MNRAS*, 510, 5246
- Lichtenberg, T., Schaefer, L. K., Nakajima, M., & Fischer, R. A. 2022, arXiv e-prints, arXiv:2203.10023
- Lightkurve Collaboration, Cardoso, J. V. d. M., Hedges, C., et al. 2018, Lightkurve: Kepler and TESS time series analysis in Python, Astrophysics Source Code Library, record ascl:1812.013
- MacLeod, M., & Oklopčić, A. 2022, *ApJ*, 926, 226
- McCann, J., Murray-Clay, R. A., Kratter, K., & Krumholz, M. R. 2019, *ApJ*, 873, 89
- Mikulášek, Z., Krtička, J., Shultz, M. E., et al. 2020, in Stellar Magnetism: A Workshop in Honour of the Career and Contributions of John D. Landstreet, ed. G. Wade, E. Alecian, D. Bohlender, & A. Sigut, Vol. 11, 46
- Murphy, S. J., Lawson, W. A., & Bessell, M. S. 2013, *MNRAS*, 435, 1325
- Nakajima, R. 1985, *Ap&SS*, 116, 285
- Neubauer, F. M. 1980, *J. Geophys. Res.*, 85, 1171
- Oksala, M. E., Wade, G. A., Townsend, R. H. D., et al. 2012, *MNRAS*, 419, 959
- Onitsuka, M., Fukui, A., Narita, N., et al. 2017, *PASJ*, 69, L2
- Paegert, M., Stassun, K. G., Collins, K. A., et al. 2021, arXiv e-prints, arXiv:2108.04778
- Palumbo, E. K., Montet, B. T., Feinstein, A. D., et al. 2022, *ApJ*, 925, 75
- Park, R. S., Vaughan, A. T., Konopliv, A. S., et al. 2019, *Icarus*, 319, 812
- Pass, E. K., Charbonneau, D., Irwin, J. M., & Winters, J. G. 2022, *ApJ*, 936, 109
- Pecaut, M. J., & Mamajek, E. E. 2016, *MNRAS*, 461, 794
- Penoyre, Z., Belokurov, V., & Evans, N. W. 2022, *MNRAS*, 513, 5270
- Petit, V., Owocki, S. P., Wade, G. A., et al. 2013, *MNRAS*, 429, 398
- Popinchalk, M., Faherty, J. K., Curtis, J. L., et al. 2023, *ApJ*, 945, 114
- Pribulla, T., Borkovits, T., Jayaraman, R., et al. 2023, *MNRAS*, 524, 4220
- Rajpurohit, A. S., Reylé, C., Allard, F., et al. 2013, *A&A*, 556, A15
- Rappaport, S., Barclay, T., DeVore, J., et al. 2014, *ApJ*, 784, 40
- Rappaport, S., Levine, A., Chiang, E., et al. 2012, *ApJ*, 752, 1
- Ratzenböck, S., Meingast, S., Alves, J., Möller, T., & Bomze, I. 2020, *A&A*, 639, A64
- Rebull, L. M., Stauffer, J. R., Cody, A. M., et al. 2018, *AJ*, 155, 196
- Rebull, L. M., Stauffer, J. R., Hillenbrand, L. A., et al. 2022, *AJ*, 164, 80
- Rebull, L. M., Stauffer, J. R., Bouvier, J., et al. 2016, *AJ*, 152, 114
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, *Journal of Astronomical Telescopes, Instruments, and Systems*, 1, 014003
- Rivinius, T., Carciofi, A. C., & Martayan, C. 2013, *A&A Rv*, 21, 69
- Robinson, C. E., Espaillat, C. C., & Owen, J. E. 2021, *ApJ*, 908, 16
- Rodriguez, D. R., van der Plas, G., Kastner, J. H., et al. 2015, *A&A*, 582, L5
- Sanchis-Ojeda, R., Rappaport, S., Pallè, E., et al. 2015, *ApJ*, 812, 112
- Sanderson, H., Jardine, M., Collier Cameron, A., Morin, J., & Donati, J. F. 2023, *MNRAS*, 518, 4734
- Servén, D., & Brummitt, C. 2018, pyGAM: Generalized Additive Models in Python
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Somers, G., Cao, L., & Pinsonneault, M. H. 2020, *ApJ*, 891, 29
- Speagle, J. S. 2020, *MNRAS*, 493, 3132
- Stassun, K. G., Kratter, K. M., Scholz, A., & Dupuy, T. J. 2012, *ApJ*, 756, 47
- Stassun, K. G., Oelkers, R. J., Paegert, M., et al. 2019, *AJ*, 158, 138
- Stauffer, J., Collier Cameron, A., Jardine, M., et al. 2017, *AJ*, 153, 152
- Stauffer, J., Rebull, L., David, T. J., et al. 2018, *AJ*, 155, 63
- Stellingwerf, R. F. 1978, *ApJ*, 224, 953
- Townsend, R. H. D., & Owocki, S. P. 2005, *MNRAS*, 357, 251
- Townsend, R. H. D., Owocki, S. P., & Groote, D. 2005, *ApJL*, 630, L81
- Vanderburg, A., Johnson, J. A., Rappaport, S., et al. 2015, *Nature*, 526, 546
- Vial, J.-C., & Engvold, O. 2015, *Astrophysics and Space Science Library*, Vol. 415, Solar Prominences
- Vines, J. I., & Jenkins, J. S. 2022, *MNRAS*, 513, 2719
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *Nature Methods*, 17, 261

- Vogt, S. S., Allen, S. L., Bigelow, B. C., et al. 1994, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 2198, *Instrumentation in Astronomy VIII*, ed. D. L. Crawford & E. R. Craine, 362
- Vos, J. M., Faherty, J. K., Gagné, J., et al. 2022, *ApJ*, 924, 68
- Waugh, R. F. P., & Jardine, M. M. 2022, *MNRAS*, 514, 5465
- Winn, J. N. 2010, in *Exoplanets*, ed. S. Seager, 55
- Winters, J. G., Charbonneau, D., Henry, T. J., et al. 2021, *AJ*, 161, 63
- Winters, J. G., Henry, T. J., Jao, W.-C., et al. 2019, *AJ*, 157, 216
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, 140, 1868
- York, D. G., Adelman, J., Anderson, John E., J., et al. 2000, *AJ*, 120, 1579
- Zhan, Z., Günther, M. N., Rappaport, S., et al. 2019, *ApJ*, 876, 127
- Zieba, S., Zwintz, K., Kenworthy, M. A., & Kennedy, G. M. 2019, *A&A*, 625, L13
- Zuckerman, B. 2019, *ApJ*, 870, 27
- Zuckerman, B., Bessell, M. S., Song, I., & Kim, S. 2006, *ApJL*, 649, L115

**Table 1.** Bonafide, candidate, and debunked complex quasiperiodic variables from the TESS 2-minute data. **For internal review, full versions are available here** <https://www.dropbox.com/scl/fo/twn4s9ckbevf75jqhtoy1/h?rlkey=t5cn8cx2uoc2ptdm9e570kp1b&dl=0>

TIC –	<i>T</i> mag	<i>d</i> pc	$G_{\text{BP}} - G_{\text{RP}}$ mag	RUWE –	<i>P</i> hr	Assoc	Age Myr	$T_{\text{eff}}$ K	$R_*$ $R_{\odot}$	$M_*$ $M_{\odot}$	$R_{\text{cr}}$ $R_*$	$P_{\text{sec}}$ hr	Quality –	Bin –	$N_{\text{sector}}$ –
368129164	9.29	18.3	2.89	6.95	6.44	ABDMG	149	3140	0.71	0.4	1.81	2.60	1	011	3
405754448	9.63	92.6	1.75	6.81	12.92	LCC	15	4273	1.5	0.82	1.74	134.4	1	011	5
167664935	10.31	62.5	2.52	5.05	14.05	UCL	16	3325	1.42	0.38	1.5	10.71	1	011	3
311092148	11.03	26.8	3.03	1.5	7.86	COL	42	3035	0.5	0.27	2.6	–	1	000	1
402980664	11.11	21.3	3.04	1.48	18.56	COL	42	3080	0.37	0.22	5.76	–	1	000	10
50745567	11.28	38.0	3.22	3.95	6.34	BPMG	24	3014	0.67	0.28	1.68	28.55	1	011	2
59836633	11.38	61.9	2.71	1.21	14.96	BPMG	24	3282	0.82	0.48	2.91	–	1	000	3
425933644	11.4	43.2	2.82	10.29	11.67	THA	45	3151	0.63	0.4	3.06	–	1	010	6
142173958	11.61	70.6	3.09	2.9	11.76	TWA	10	3028	1.15	0.26	1.45	12.84	1	011	3
146539195	11.62	48.2	3.37	2.86	6.73	BPMG	24	2898	0.8	0.24	1.38	7.29	1	011	2
206544316	11.63	42.8	2.89	1.26	7.73	THA	45	3114	0.57	0.35	2.44	–	1	000	6
335598085	11.9	105.5	2.85	2.79	15.85	LCC	15	3119	1.34	0.28	1.56	17.94	1	011	3
405910546	12.11	111.9	2.36	1.09	37.99	LCC	15	3455	0.92	0.6	5.26	–	1	100	4
272248916	12.15	80.5	2.83	5.5	8.9	UCL	16	3193	0.81	0.4	1.97	50.7	1	011	3
178155030	12.17	46.8	2.91	1.29	11.67	THA	45	3097	0.49	0.3	3.53	–	1	000	4
224283342	12.29	38.0	3.04	1.27	21.3	COL	42	3050	0.39	0.22	6.08	–	1	100	3
89026133	12.31	131.6	2.82	4.0	11.2	UCL	16	3188	1.33	0.31	1.29	27.83	1	011	3
234295610	12.51	48.1	3.04	1.13	18.29	THA	45	3074	0.44	0.27	5.09	–	1	000	3
118449916	12.54	97.1	3.09	25.18	12.31	TAU	2	3025	1.04	0.28	1.7	6.71	1	011	4
67897871	12.55	148.2	3.01	2.55	6.23	USCO	10	3082	1.5	0.18	0.65	6.72	1	011	2
353730181	12.65	106.6	2.75	1.23	13.51	TAU	2	3253	0.8	0.41	2.65	–	1	000	4
201898222	12.68	42.2	3.21	1.29	10.7	THA	45	2996	0.39	0.2	3.69	13.62	1	001	5
264767454	12.73	123.3	2.93	12.3	10.01	COL(?)	42	3150	1.0	0.42	1.76	20.62	1	011	13
442571495	12.75	80.8	3.03	1.64	9.59	UCL	16	3099	0.65	0.3	2.35	13.82	1	001	3
2234692	12.8	53.7	3.0	1.2	6.52	COL	42	3098	0.44	0.26	2.56	59.8	1	001	7
94088626	12.88	57.6	3.07	1.12	6.6	ARG	45	3090	0.46	0.27	2.53	–	1	000	2
264599508	12.88	79.7	3.01	1.89	7.9	COL	42	3098	0.62	0.4	2.4	8.99	1	001	7
363963079	12.92	83.1	3.09	8.0	7.82	ARG	45	3040	0.67	0.4	2.21	7.41	1	011	7
193831684	13.03	51.6	3.23	1.16	31.02	BPMG	24	2971	0.42	0.2	6.87	–	1	000	3
177309964	13.1	91.0	2.94	1.15	10.88	CAR	45	3125	0.62	0.4	2.95	–	1	000	34
425937691	13.18	43.1	3.77	2.86	4.82	THA	45	2782	0.41	0.16	1.91	3.22	1	011	5
141146667	13.28	57.6	3.28	1.23	3.93	FIELD	NaN	2968	0.42	NaN	NaN	–	1	000	6
332517282	13.29	39.0	3.27	1.05	9.67	ABDMG	149	2975	0.28	0.2	4.87	–	1	000	3
144486786	13.3	77.4	3.05	15.05	6.82	COL	42	3074	0.51	0.3	2.38	11.49	1	011	4
38820496	13.3	44.1	3.37	1.08	15.73	THA	45	2903	0.34	0.16	5.13	–	1	000	5
289840926	13.31	40.2	3.75	1.16	4.8	BPMG	24	2807	0.36	0.14	2.05	15.64	1	001	3
404144841	13.33	77.1	3.19	1.11	10.74	TWA	10	3008	0.52	0.22	2.86	–	1	000	4
89463560	13.45	123.9	2.97	1.31	9.43	ARG	45	3055	0.75	0.37	2.15	7.76	1	001	10
300651846	13.49	109.2	2.86	1.16	8.26	CAR	45	3136	0.62	0.4	2.44	–	1	000	31
267953787	13.49	130.5	3.59	1.2	17.46	TAU	2	2826	1.06	0.12	1.59	–	1	000	4
68812630	13.6	123.8	3.22	1.63	9.04	TAU	2	2996	0.76	0.27	1.86	5.28	1	001	3
141306513	13.65	50.2	3.4	1.08	13.36	THA	45	2964	0.32	0.16	4.85	–	1	000	2
201789285	14.03	45.4	3.82	1.19	3.64	THA	45	2757	0.3	0.12	2.02	–	1	000	5
294328887	14.23	97.1	3.22	1.05	8.51	CARN	200	2994	0.45	0.35	3.33	–	1	000	35

**Table 1** *continued*

**Table 1** (*continued*)

312410638	14.3	136.9	3.12	1.09	28.06	UCL	16	3030	0.58	0.25	5.11	-	1	000	3
38539720	14.52	129.4	3.37	1.2	9.16	PERI	120	2924	0.57	0.25	2.43	-	1	000	1
359892714	14.53	95.5	4.04	1.07	11.33	EPSC	3	2675	0.55	0.13	2.36	-	1	000	6
118769116	14.58	119.0	3.6	1.13	8.56	TAU	2	2852	0.56	0.2	2.18	-	1	000	4
440725886	14.69	135.1	2.96	1.06	3.92	PLE	112	3109	0.45	0.35	1.99	-	1	000	5
397791443	15.01	151.1	3.1	1.06	6.95	IC2602	46	3031	0.48	0.26	2.46	-	1	000	6
160329609	9.65	8.7	3.4	1.18	24.31	ARG	45	2912	0.35	0.16	6.6	-	0	000	3
148646689	12.14	140.4	2.44	1.7	10.63	UCL	16	3466	1.25	0.55	1.61	13.37	0	001	3
280945693	12.27	98.2	2.97	1.16	15.27	LCC	15	3103	1.09	0.31	1.94	-	0	100	5
165184400	12.37	43.2	3.02	1.24	15.91	THA	45	3076	0.42	0.25	4.74	-	0	000	4
245834739	12.55	115.4	2.85	1.49	10.47	TAU	2	3112	1.02	0.36	1.68	9.86	0	001	6
125843782	13.01	127.7	2.86	1.21	44.17	TAU	2	3135	0.9	0.35	4.97	-	0	000	4
244161191	13.17	44.7	3.54	1.28	7.17	COL	42	2860	0.38	0.18	2.8	8.39	0	001	3
231058925	13.17	51.2	3.25	1.35	8.87	THA	45	2978	0.38	0.2	3.33	-	0	000	5
301676454	13.4	70.7	3.07	1.24	9.18	ARG	45	3009	0.47	0.25	2.96	-	0	000	1
58084670	13.58	140.2	2.82	1.06	11.16	FIELD	NaN	3138	0.77	NaN	NaN	-	0	000	6
67745212	13.63	27.8	3.8	1.11	5.12	COL	42	2781	0.21	0.09	3.17	-	0	000	2
5714469	13.73	78.3	3.65	1.12	10.35	UCL	16	2828	0.54	0.18	2.53	-	0	000	3
259586708	13.82	95.6	2.93	1.17	22.52	COL	42	3133	0.46	0.29	5.84	-	0	000	7
435903839	11.95	80.7	2.49	17.7	10.82	ABDMG(?)	149	3458	0.76	0.54	2.66	-	-1	010	6
57830249	11.96	48.8	3.2	1.34	43.82	TWA	10	2948	0.7	0.25	5.63	-	-1	000	3
193136669	13.06	61.1	3.49	1.22	37.64	TWA	10	2855	0.58	0.19	5.62	-	-1	000	4

NOTE—This table includes 50 CQVs (Quality flag 1), 13 ambiguous CQVs (Quality flag 0), and 3 impostors (Quality flag -1). The three-bit binarity flag “Bin” is for Gaia DR3 `radial_velocity_error` outliers (bit 1), Gaia DR3 `ruwe` outliers (bit 2), and stars with multiple TESS periods (bit 3). The machine-readable version, available online, includes additional columns for the Gaia DR2 and DR3 source identifiers, as well as the stellar parameter uncertainties. The age uncertainties are typically  $\approx \pm 10\%$ , but can be asymmetric. **The median temperature, radius, and mass uncertainties are  $\pm XXX\text{ K}$ ,  $\pm XX\%$ , and  $\pm YY\%$  respectively.**  $N_{\text{sector}}$  denotes the number of TESS sectors for which *any* data are expected to be acquired between July 2018 and Oct 2024. This number is generally greater than the number of sectors for which 120-second cadence data exist. Association names and provenance follow conventions adopted by Gagné et al. (2018): ABDMG: AB Doradus moving group (Bell et al. 2015). ARG: Argus (Zuckerman 2019). BPMG:  $\beta$  Pic moving group (Bell et al. 2015). CARN: Carina Near moving group (Zuckerman et al. 2006). COL: Columba (Bell et al. 2015). EPSC:  $\epsilon$  Chamæleonis (Murphy et al. 2013). LCC: Lower Centaurus Crux (Pecaut & Mamajek 2016). PERI: Pisces-Eridani (Curtis et al. 2019). PLE: Pleiades (Dahm 2015). TAU: Taurus (Kenyon & Hartmann 1995). THA: Tucana-Horologium association (Bell et al. 2015). TWA: TW Hydriæ association (Bell et al. 2015). UCL: Upper Centaurus Lupus (Pecaut & Mamajek 2016). USCO: Upper Scorpius (Pecaut & Mamajek 2016). The “(?)” string denotes low-confidence membership.

## APPENDIX


### A. TIC 300651846

Figures 10 and 11 show 120-second cadence data for TIC 300651846, a CQV in the TESS continuous viewing zone. If it were not for the existence of TIC 402980664, this source would probably have received greater attention. With the exception of a few sectors, TESS data will exist for TIC 300651846 for at least Sectors 1-12, 27-39, and 61-69. While most of the available data will exist in the full frame images, in Figures 10 and 11 focus only on the currently available 120-second cadence data.

During Sectors 32-39, the source shows between one and four local minima per cycle. During the early portions of Sectors 61-65, it is somewhat more complex, with at least five clear local minima per cycle. As the source evolves, this complexity decreases, while the sharpness of one of the minima appears to increase. We did not find evidence for any obvious “state-switches” analogous to those that we observed in LP 12-502; gradual evolution over timescales of  $\approx 50$ -100 cycles seem to be the norm for TIC 300651846. Unlike the TIC 402980664 river plots (Figure 8), we did not subtract any “continuum sinusoid” for this source, because the continuum is not as obviously defined.


### B. NO SIGNIFICANT POWER AT 20 SECOND CADENCE

TESS was the first instrument to show that CQV light curves contain power at timescales of a few minutes (Zieba et al. 2019; Günther et al. 2022). This advance was enabled by the fifteen-fold faster cadence in the TESS 2-minute data, relative to K2. A logical follow-up is to ask whether the periodic components of the CQV light curves contain power at timescales below one



**Figure 10.** Light curve evolution of TIC 300651846. All available 120-second cadence data as of 2023 Aug 11 are shown. Cycles 0 to 622 span TESS Sectors 32-39 (Nov 2020–June 2021); cycles 2296-2676 span Sectors 61-65 (Jan–June 2023). We assumed a 8.254 hr period and a fixed reference epoch (BTJD 2174.127) for both panels. Light curve segments are split based on the presence of gaps longer than three hours. Cycle numbers are listed in the lower-right of each light curve segment.

minute. Between 2020 and 2021, we observed 10 CQVs at 20-second cadence with TESS in order to explore this question (TESS DDT029). The stars were TICs 142173958, 146539195, 24518895, 276453848, 264599508, 363963079, 144486786,



**Figure 11. River plots of TIC 300651846.** This is an alternative visualization of the data in Figure 10. All available 120-second cadence data as of 2023 Aug 11 are shown. Cycles 0 to 622 span TESS Sectors 32-39 (Nov 2020–June 2021); cycles 2296-2676 span Sectors 61-65 (Jan–June 2023). We assumed  $P=8.254$  hr and  $t_0=2174.127$  [BTJD]. Note that the two panels have slightly different color scales.

408188366, 300651846, 262400835. These sources were selected from CQVs known at the time to have short periods and sharp features when observed at 2-minute cadence. Comparing the 20-second to 120-second data for these stars (data available on MAST), we concluded that these CQVs did not contain appreciable power at timescales shorter than a few minutes.