

## Rotation and Lithium Confirmation of a 500 Parsec Halo for the Open Cluster NGC 2516\*

2 L. G. BOUMA,<sup>1</sup> J. L. CURTIS,<sup>2,3</sup> J. D. HARTMAN,<sup>1</sup> J. N. WINN,<sup>1</sup> AND G. Á. BAKOS<sup>1,4</sup>

3 <sup>1</sup>*Department of Astrophysical Sciences, Princeton University, 4 Ivy Lane, Princeton, NJ 08540, USA*

4 <sup>2</sup>*Department of Astronomy, Columbia University, 550 West 120th Street, New York, NY 10027, USA*

5 <sup>3</sup>*Department of Astrophysics, American Museum of Natural History, Central Park West, New York, NY 10024, USA*

6 <sup>4</sup>*Institute for Advanced Study, 1 Einstein Drive, Princeton, NJ 08540, USA*

7 (Received April 29, 2021; Revised —; Accepted —)

8 Submitted to AAS journals.

### ABSTRACT

Recent analyses of the Gaia data have identified diffuse stellar populations surrounding nearby open clusters. It is important to verify that these “halos”, “tails”, and “strings” are of similar ages and compositions as stars in the denser part of the cluster. We present an analysis of NGC 2516 ( $\approx 150$  Myr), which has a classical tidal radius of 10 pc and an apparent halo of stars spanning 500 pc ( $20^\circ$  on-sky). Combining photometry from Gaia, rotation periods from TESS, and lithium measurements from Gaia-ESO and GALAH, we find that the halo of NGC 2516 is the same age as the cluster’s core. Two thirds of kinematically selected halo members out to 250 pc from the cluster center have rotation periods consistent with a gyrochronological age of 150 Myr. A comparison sample of field stars shows no such trend. The lithium abundances of stars in the halo are higher than in the field, and are correlated with the stellar rotation rate and binarity fraction, as has been noted in other young open clusters. Broadly speaking, this work supports a new paradigm wherein the halos of open clusters are often more populous than their cores. We highlight implications for spectroscopic survey targeting, open cluster dispersal, and planet searches around young stars.

Keywords: stellar associations (1582), open star clusters (1160), stellar rotation (1629), stellar ages (1581), stellar kinematics (1608)

### 1. INTRODUCTION

Star clusters form when dense filaments in hierarchically structured molecular clouds collide and collapse (Shu et al. 1987). Over the first 10 Myr, feedback effects including protostellar outflows, photoionization, radiation pressure, and supernova shocks disperse the gas (Krumholz et al. 2019). Since the gas represents about 99% of the mass of the original cloud, gas dispersal enables the majority ( $\sim 90\%$ ) of stars in the cluster to escape the cluster’s gravitational well (Lada & Lada 2003).

From 10 to 1000 Myr, the cluster remnants that survive gas dispersal suffer an onslaught of supernovae, AGB winds, and close stellar encounters that almost always leads to dissolution (Lamers et al. 2010). Extrinsic to the cluster, collisions with giant molecular clouds (Spitzer 1958), and perturbations from the galactic tide in both the radial and vertical dimen-

sions further promote stellar escape (e.g., Fukushige & Heggie 2000; Bergond et al. 2001).

Finding the stars that have dispersed from their clusters into the galactic field is a pressing topic for a few reasons. One is to understand the spatial extent and duration of star formation events (e.g., Wright & Mamajek 2018). Another is to understand the Sun’s birth environment (Adams 2010). The plausibility of finding the stars that formed with the Sun, whether through chemical tagging (Freeman & Bland-Hawthorn 2002; Hogg et al. 2016; Ness et al. 2018), kinematic analyses, or both, can be tested by first using the remnants of open clusters that have only recently dissipated into the field.

A separate project that benefits from the new discoveries of dispersed stellar populations is that of finding young transiting exoplanets (e.g., Mann et al. 2016; Ciardi et al. 2018; David et al. 2019; Livingston et al. 2019; Bouma et al. 2020; Rizzuto et al. 2020; Plavchan et al. 2020; Newton et al. 2021; Tofflemire et al. 2021; Zhou et al. 2021). Young transiting planets are hard to find because young stars are rare, and mostly reside in the crowded galactic plane (Kharchenko et al. 2013; Piskunov et al. 2018). If the dispersed halos of nearby star clusters could be reliably identified, this could

Corresponding author: L. G. Bouma  
luke@astro.princeton.edu

\* This is paper 3 of the Cluster Difference Imaging Photometric Survey.

expand the census of nearby young stars by up to a factor of 10, based on the expected fraction of stars thought to be lost during gas dispersal.

Although it has been possible to detect dispersed stellar associations for a long time, Gaia is now enabling major advances (e.g., de Zeeuw et al. 1999; Bergond et al. 2001; Zuckerman & Song 2004; Oh et al. 2017; Cantat-Gaudin et al. 2018; Gagné et al. 2018; Gagné & Faherty 2018; Kounkel et al. 2018; Zari et al. 2018; Kounkel & Covey 2019; Fürnkranz et al. 2019). The populations found by the most recent studies can be summarized as follows. On one end are groups with no discernable cores (e.g., Psc-Eri and  $\mu$ -Tau, Meingast et al. 2019; Curtis et al. 2019; Gagné et al. 2020). On the other are hierarchically structured associations with many over and under-densities (e.g. the Sco-Cen and Vela associations, Pecaut & Mamajek 2016; Cantat-Gaudin et al. 2019). Here, we focus on an intermediate regime: low-density halos associated with a single overdensity, typically an open cluster that was known before Gaia (see Kounkel & Covey 2019; Kounkel et al. 2020; Meingast et al. 2021). In some cases, these halos may actually be tidal tails, as have been reported in the Hyades (Meingast & Alves 2019; Röser et al. 2019), the Ursa Major moving group (Gagné et al. 2020), and Coma Berenices (Tang et al. 2019; Fürnkranz et al. 2019).

One point of difficulty however is that different algorithms for identifying clusters yield differing levels of sensitivity and precision (Hunt & Reffert 2020). For example, using a Gaussian Mixture Model tautologically yields open clusters that are elliptical (e.g., Wallace 2018). Some unsupervised methods yield dispersed and asymmetric structures with number densities down to 100 times lower than the field around the same regions (e.g., Kounkel & Covey 2019 and Meingast et al. 2021). Since different techniques yield quantitatively and qualitatively different outcomes, the purity of kinematically selected samples is important to verify through independent lines of evidence, including rotation periods and spectroscopy.

As part of a Cluster Difference Imaging Photometric Survey (CDIPS, Bouma et al. 2019), we have been making TESS light curves of stars reported to be members of coeval populations and searching them for planets (Bouma et al. 2020). Our analysis of TESS Sectors 1-13 yielded light curves of 483,407 candidate cluster members in the Southern Ecliptic hemisphere, which are available on MAST.<sup>1</sup>

As part of this project, we focus here on a single southern open cluster, and ask: is the cluster halo truly coeval with the core? We chose NGC 2516 for this analysis because it is young (100-200 Myr) and close enough ( $d \approx 400$  pc) to facilitate photometric rotation measurements using TESS. Its halo was reported by Kounkel & Covey (2019) to span  $\approx 20 \times 10 \times 350$  pc. We want to know: is this halo real or is it just a collection of interlopers, foreground and background stars? To what extent can we use Gaia alone to reliably iden-

tify age-dated needles in the haystack of field stars? What are the implications, observationally and theoretically, if we can identify the dispersed halos of open clusters?

A brief note on terminology. Low-density stellar associations connected to a dense population (a “core”) have been described as being in “halos”, “strings”, “coronae”, “snakes”, “outskirts”, and “tidal tails” (e.g., Chumak & Rastorguev 2006a; Davenport & Sandquist 2010; Kounkel & Covey 2019; Röser et al. 2019; Tian 2020; Meingast et al. 2021). The latter term implies a particular model for the formation of the dispersed group. “Halo” is model agnostic, but is not an ideal term because it can connote spherical symmetry, which is rarely the case. The halo of NGC 2516 is most accurately described as consisting of a leading and trailing tail. We call it a halo for simplicity.

Section 2 summarizes the astrometric analyses of the Gaia data that led to our interest in NGC 2516. Section 3 measures the age of the cluster’s halo and core using Gaia photometry (Section 3.1), TESS gyrochronology (Section 3.2), and lithium equivalent widths (Section 3.3). In Section 4 we discuss the implications of our analysis for open cluster dispersal, stellar spin-down, and the lithium-rotation correlation. Section 5 gives our conclusions.

## 2. A DISPERSED HALO AND A CORE?


We selected candidate NGC 2516 members based on those reported in the literature. While some pre-Gaia analyses were available (Jeffries et al. 2001; Kharchenko et al. 2013), the purity and accuracy of the Gaia-derived results are the current state of the art. We therefore adopted what we viewed as the most important Gaia-based samples to compare: those of Cantat-Gaudin et al. (2018), Kounkel & Covey (2019) and Meingast et al. (2021). We refer to these studies in the following as CG18, KC19, and M21 respectively. A useful visualization of these samples is available online.<sup>2</sup>

The Gaia clustering studies each used different selection functions, which yielded different results. CG18 considered stars brighter than  $G = 18$  mag within a few degrees of the center of NGC 2516, and reported 1106 candidate cluster members. KC19 and M21 considered stars up to  $\approx 1$  mag fainter, and reported 3003 and 1860 members respectively. The unsupervised clustering techniques that each of these studies applied to the second Gaia data release are discussed in Appendix A, as is the overlap between their resulting membership catalogs.

Figure 1 shows the cluster members reported by each study in the space of observed positions, proper motions, and radial velocity when available. In Figure 1, and for the remainder of the study, we describe the CG18 members as the “core” of the cluster, and any non-overlapping KC19 and M21 members as the “halo”. This definition implies that there are 1,106 core members, and 2,192 halo members. The distinction is tautological, in the sense that CG18 did not extend their

<sup>1</sup> <https://archive.stsci.edu/hlsp/cdips>

<sup>2</sup> <https://homepage.univie.ac.at/stefan.meingast/coronae.html>, made by Meingast et al. (2021), last accessed 2021/04/05.



**Figure 1. The dense and diffuse components of NGC 2516 across position and velocity space.** The core (black) was analyzed by Cantat-Gaudin et al. (2018) using Gaia DR2, and is coincident with the visual overdensity of stars canonically accepted as “the cluster”. The halo (blue) is a concatenation of studies by Kounkel & Covey (2019) and Meingast et al. (2021), which used less restrictive membership assignment algorithms described in Appendix A. The field sample is randomly drawn from an  $(\alpha, \delta, \pi)$  cube centered on the cluster. The low volume density of the halo stars makes it difficult to visually distinguish the field and the halo samples.

search for members out to tens of degrees from the cluster center. Nonetheless, the CG18 membership catalog is consistent with that of many earlier investigators (e.g., Jeffries et al. 2001; Kharchenko et al. 2013), and is consistent with the general visual overdensity one sees when viewing NGC 2516 in the sky: it appears to span  $\approx 2^\circ$ , not  $\approx 20^\circ$ .

Outside of the core and halo, we also define a set of nearby field stars in the “neighborhood” of NGC 2516. Based on the observed distribution of halo members, we drew these


stars randomly from the following intervals of right ascension, declination, and parallax:

$$\alpha [\text{deg}] \in [108, 132], \quad (1)$$

$$\delta [\text{deg}] \in [-76, -45], \quad (2)$$

$$\pi [\text{mas}] \in [1.5, 4.0]. \quad (3)$$

<sup>169</sup> We imposed a magnitude limit of  $G = 19$  mag, and ran the  
<sup>170</sup> queries using the `astroquery.gaia` module (Ginsburg



**Figure 2. Position and orbit of NGC 2516 in the Galaxy.** Points are halo members with  $\pi/\sigma_\pi > 20$ ; the cluster is most elongated viewed top-down (left). The Sun is at  $\{X, Y, Z\} = \{-8122, 0, +20.8\}$  pc. Gray lines trace the past (solid) and future (dashed) cluster orbit, and blue arrows denote the mean cluster velocity after subtracting the local standard of rest:  $\{v_X, v_Y, v_Z\} = \{-22.2, -25.3, -4.6\}$  km s<sup>-1</sup>.

et al. 2018). We allowed the number of stars in the comparison sample to exceed that in the cluster sample by a factor of  $\approx 5$ , to ensure broad sampling of stellar masses and evolutionary states. We also required the comparison sample to not overlap with the cluster sample.

The style of visualization given in Figure 1 does not make it clear, in our eyes, why the clustering algorithms have decided to associate certain stars with the halo. Canonically, open clusters are spherical, and span  $\approx 10$  pc (e.g., Kharchenko et al. 2013). Inverting the parallaxes in Figure 1 shows that members have been reported from line-of-sight distances ranging from  $\approx 300$  to  $\approx 600$  pc. Is this structure real? What explains its existence?

An initial step in visualizing the structure and kinematics of the candidate cluster members is to consider their Cartesian coordinates (Figure 2). We computed these positions by transforming from  $(\alpha, \delta, \pi)$  to galactic  $(X, Y, Z)$  assuming the astropy v4.0 coordinate standard (Astropy Collaboration et al. 2018). The Galaxy rotates in the  $+Y$  direction. The cluster orbit (gray line) was evaluated by taking the median parameters for core members for which CG18 reported membership probabilities exceeding 70%. We then integrated the orbit using gala and the MWPotential2014 potential (Bovy 2015; Price-Whelan 2017).

The general shape of the cluster itself seems to include a central overdensity and two tails (the halo). One tail is leading the cluster's orbit, and is angled toward the center of the galaxy when viewed top-down. The other tail is trailing the cluster's orbit, and is pointed away from the center of the galaxy. The elongation of the cluster in both the  $(X, Y)$  and  $(Z, Y)$  planes is correlated with the direction of the LSR-corrected median cluster velocity. Possible explanations for this overall morphology are discussed in Section 4.1.

### 3. AGE-DATING THE HALO OF NGC 2516

#### 3.1. HR Diagram from Gaia


The first check on whether the candidate cluster members share the same age is to analyze the Gaia Hertzsprung-Russell (HR) diagrams. Comparable analyses have previously been performed by CG18, KC19, and M21.

Figure 3 shows the HR diagram in the space of absolute Gaia  $G$  magnitude as a function of observed  $G_{\text{BP}} - G_{\text{RP}}$  color. Magnitudes are from Gaia EDR3; we performed the Gaia DR2 to EDR3 cross-match using the pre-computed gaiaedr3.dr2\_neighbourhood table available at the Gaia archive, and required the closest (proper motion and epoch-corrected) source to be the single match. Spectral types are interpolated from the Pecaut & Mamajek (2013) table, after accounting for reddening as described below.<sup>3</sup>

Stars in the core appear consistent with having a fixed age and metallicity, and varying mass. The halo stars show a similar sequence, though with greater scatter. One possible explanation for the additional scatter is that the halo is more contaminated by field stars. For instance,  $\approx 5$  red giants in the halo must be field interlopers, because their isochronally implied ages would be inconsistent with that of the cluster.

Another possible explanation for scatter in the halo's HR diagram is differential reddening across different sightlines. The reported halo spans  $20^\circ$  on-sky, and varies in position from about  $b = -12^\circ$  to  $b = -20^\circ$ , with the stars closest to the galactic plane also being further from the Sun by up to 300 pc (Figure 1). An HR diagram binned by galactic latitude did show some minimal evidence for differential reddening, and so we expect that both field star contamination and differential reddening could play a role in the larger scatter of the halo relative to the core. A third possibility that we have not explored is whether the binary fraction could also differ between the different regions of the cluster.

<sup>3</sup> [http://www.pas.rochester.edu/~emamajek/EEM\\_dwarf\\_UBVIJHK\\_colors\\_Teff.txt](http://www.pas.rochester.edu/~emamajek/EEM_dwarf_UBVIJHK_colors_Teff.txt), version 2021/03/02.



**Figure 3. HR diagrams of NGC 2516.** *Left:* Gaia EDR3 photometry for the core (black) shows a main-sequence and turnoff consistent with stars of a fixed age and metallicity. The halo (blue) is similar, though the binary sequence is less defined. The faintest M dwarfs in the core and halo are brighter than field stars (gray) with the same color, consistent with the cluster M dwarfs not having yet reached the main-sequence. *Right:* PARSEC isochrones ( $[{\rm Fe/H}] = +0.06$ ) are overplotted at intervals of  $\log_{10}(t/{\rm yr}) = \{8, 8.25, 8.5\}$ . The arrow represents the average reddening correction applied to the models. The opacity of the M-dwarf model atmospheres is less well-calibrated for  $M_* \lesssim 0.45M_\odot$  ( $\approx M2V$ ); these model values are shown with dashed lines.

In the right panel of Figure 3, we compare the observed Gaia EDR3 photometry with PARSEC isochrones (Bressan et al. 2012; Chen et al. 2014, 2015; Marigo et al. 2017). We used the web interface<sup>4</sup> to interpolate these isochrones at  $\log_{10}(t/{\rm yr}) = \{8, 8.25, 8.5\}$ .

To determine the reddening correction across NGC 2516, we queried the 2MASS DUST service<sup>5</sup> and retrieved the total line-of-sight extinction parameters at the positions of each NGC 2516 member. The mean and standard deviation of the  $E(B-V)$  values for the Schlegel et al. (1998) and Schlafly & Finkbeiner (2011) maps were consistent, so we adopted the average from Schlegel et al. (1998):  $E(B-V)_s = 0.206 \pm 0.039$ . Bonifacio et al. (2000) found however that

the Schlegel et al. (1998) maps overestimate the reddening values when the color excess exceeds about 0.10 mag. We therefore applied the correction proposed by Bonifacio et al. (2000), and additionally corrected for the mean cluster distance by a factor  $\exp(-|d \sin(b)|/h)$  where  $d$  is the average cluster distance,  $b$  is the average galactic latitude, and  $h$  is the scale height of the galactic disk, assumed to be 125 pc. This yielded our adopted  $E(B-V) = 0.102 \pm 0.019$ , where the uncertainty reflects the standard deviation of the individual Schlegel et al. (1998) values. Finally, to convert to Gaia colors we used the calibration of Stassun et al. (2019), namely  $E(G_{BP} - G_{RP}) = 1.31E(B-V)$  and  $A_G = 2.72E(B-V)$ . This yielded  $E(G_{BP} - G_{RP}) = 0.134 \pm 0.025$ , which is used in the plots. To redden the isochrones, we assumed  $R_V = 3.1$ , and applied the O'Donnell (1994) SED-dependent extinction law star-by-star through the PARSEC interface.

For the metallicity, we considered a range of super and sub-solar metallicities to fit as much of the locus from  $0.5 < G_{BP} - G_{RP} < 1.5$  as possible, and settled by eye on the

<sup>4</sup> <http://stev.oapd.inaf.it/cgi-bin/cmd>, 2021/02/26 CMD3.4, YBC bolometric corrections as in Chen et al. (2019).

<sup>5</sup> <http://irsa.ipac.caltech.edu/applications/DUST>; query performed using the astrobase.services.dust module (Bhatti et al. 2018).

slightly super-solar  $[M/H] = 0.06$  (Cummings 2011). Sub-solar metallicities led to model predictions too blue along the main sequence by  $\approx 0.1$  mag. Our adopted metallicity agrees with the spectroscopic metallicity from Cummings (2011, Sec 4.4.4), and with the iron abundance determined by the Gaia-ESO team (Baratella et al. 2020). The latter result represented an up-revision from an earlier sub-solar metallicity (Randich et al. 2018). We caution though that other sub-solar metallicities have been reported (Bailey et al. 2018), and note that the cluster metallicity and reddening are degenerate; if the reddening is lowered, the implied metallicity will increase. While a detailed redetermination of these parameters from the Gaia photometry is beyond our scope, our adopted values for both the reddening and metallicity are within the range of previously reported values in the literature.

Overall, the data and models agree for masses above  $\approx 0.45 M_{\odot}$ . Below this mass, the data and models diverge at colors redder than  $G_{\text{BP}} - G_{\text{RP}} \approx 2.2$ , in the sense that the model isochrones have lower luminosities and bluer colors than the observations.<sup>6</sup> The MIST isochrones showed a comparable disagreement (Choi et al. 2016). Proposed explanations for the discrepancy between the models and observations include starspots and incomplete molecular line lists (e.g., Stauffer et al. 2003; Feiden & Chaboyer 2013; Rajpurohit et al. 2013; Mann et al. 2013; Choi et al. 2016). Ultimately, we adopted the PARSEC isochrones because Chen et al. (2014) implemented an empirical temperature-opacity calibration, which leads the PARSEC isochrones to diverge at slightly lower mass than the MIST models. Regardless, for purposes of age-dating the cluster we focus on the main-sequence turn-off (MSTO), since this is where the models have maximum sensitivity to age.

Cummings & Kalirai (2018) presented techniques for mitigating some of the complexities of MSTO age-dating (e.g., sparse turnoffs, stellar rotation, high binarity fractions, and the presence of blue stragglers). Combining a *UBV* color-color analysis with Gaia DR2 cluster memberships, they found MSTO ages for NGC 2516 ranging from 165 to 195 Myr, depending on the choice of model isochrone ( $\chi^2$ , PARSEC, MIST, or SYCLIST).

Our goal is to determine whether the ages of the core and halo are consistent. The left panel of Figure 3 suggests that isochronally, they are consistent: stars past the turnoff in both the halo and core are on the same locus. The right panel of Figure 3 demonstrates the precision with which the claim can be made. The MSTO stars are consistent with the 178 and 316 Myr models, but are bluer than the 100 Myr model. The red-giant-branch (RGB) stars are consistent only with the 178 Myr ( $\log_{10} t/\text{yr} = 8.25$ ) model. Based on the assumption that the width of the MSTO can be attributed to binary stars, we are most interested in the blue edge (blue stragglers are, however, a concern). The blue edge appears most

compatible with the 178 Myr model. These results are consistent with the MSTO age range of 165 to 195 Myr found by Cummings & Kalirai (2018), and we refer to their work for the more precise model-dependent comparison. Based on Table 3 of Cummings & Kalirai (2018), we therefore adopt a MSTO age for NGC 2516 of  $175 \pm 35$  Myr, where our quoted uncertainty is a quadrature sum of the statistical and systematic components from Cummings & Kalirai (2018).

### 3.2. Rotation from TESS

#### 3.2.1. Methodology


We began our rotation analysis by considering all 3,298 candidate members of NGC 2516. For each source, we retrieved all available CDIPS light curves, on a per-sector basis. This yielded 2,205 stars with at least one sector from a CDIPS light curve. Each of these stars is brighter than the CDIPS magnitude limit of  $G_{\text{RP}} = 16$  mag. 2,270 of the stars from the NGC 2516 source list have  $G_{\text{RP}} < 16$ . The difference (65 stars) is due to 35 stars from Meingast et al. (2021) which were not available at the time of the CDIPS reductions, and 30 stars falling on the TESS chip gaps. At the distance of NGC 2516, the  $G_{\text{RP}} = 16$  mag cutoff imposed during the CDIPS processing corresponds roughly to  $(G_{\text{BP}} - G_{\text{RP}})_0$  of 2.2, or a spectral type of  $\approx$ M2V.

We removed systematic trends common to the light curves on each CCD in each individual sector, and stitched together the resulting light curves before searching for rotation-induced periodicity. Details regarding our detrending approach are presented in Appendix B. After detrending, we proceeded with a few cleaning steps: we masked 0.7 days at the beginning and end of each spacecraft orbit, and ran a sliding standard-deviation rejection window over the light curve, which removed any outlying points within  $\pm 3 \times \text{MAD}$  of the median in each window.

We then measured the rotation period from the resulting light curve using the periodogram implementations in astrobase, namely the traditional Lomb-Scargle, and the Stellingwerf phase-dispersion minimization algorithm (Lomb 1976; Stellingwerf 1978; Scargle 1982; Stellingwerf 2011; Bhatti et al. 2018). We used the CDIPS aperture radius that, based on theoretical expectations, was expected to give the optimal balance between light from the target star and the sky background (Sullivan et al. 2015). This typically resulted in a circular aperture radius of either 1 or 1.5 pixels. We recorded the top five periodogram peaks from each method, and their corresponding powers. Finally, as a check on crowding, we recorded the number of stars within the aperture with greater brightness than the target star, and with brightness within 1.25 and 2.5 TESS magnitudes of the target star.

The resulting rotation periods and periodogram powers are reported, regardless of detection significance, in Table 1. To clean these measurements, we designed automated cleaning criteria, which yielded sets we denote  $\mathcal{A}$  and  $\mathcal{B}$ . For both sets, we considered light curves for which the peak Lomb-Scargle periodogram period was below 15 days, the normal-

<sup>6</sup> The 100 Myr isochrone does overlap well with the lower main sequence, but it fails to fit the upper main sequence and red giants.



**Figure 4.** TESS rotations periods and dereddened Gaia colors across the core and halo of NGC 2516. Sets  $\mathcal{A}$  (top) and  $\mathcal{B}$  (bottom) undergo successive stages of automated cleaning (see Section 3.2.1). The Pleiades (125 Myr; Rebull et al. 2016) and Praesepe (650 Myr; Douglas et al. 2017) are shown for reference. Stars in the core and the halo of NGC 2516 overlap with the Pleiades at  $(G_{\text{BP}} - G_{\text{RP}})_0 < 1.2$ . From  $1.2 < (G_{\text{BP}} - G_{\text{RP}})_0 < 1.7$ , some NGC 2516 members have longer rotation periods than Pleiades members. Rotation periods for a comparison sample of field stars are shown in Appendix C.

ized Lomb-Scargle power exceeded 0.08, and for which no companions with brightness exceeding one-tenth of the target star were in the aperture (*i.e.*, excluding visual binaries with  $\Delta T < 1.25$ , according to the Gaia DR2 source catalog). These limits were chosen after inspecting the light curves visually, to eliminate low-quality rotation periods while retaining high completeness. For Set  $\mathcal{A}$ , this yielded 987 light curves, out of 1,641 stars for which rotation periods might plausibly have been detected. For Set  $\mathcal{B}$ , we additionally required that the Lomb-Scargle and Stellingwerf phase-dispersion methods yielded identical periods to within 10%. This yielded 507 light curves. Boolean flags for each set are included in Table 1. A set of field stars was also analyzed for comparison. The results are discussed in Appendix C.

### 3.2.2. Rotation-Color Diagrams

Figure 4 shows the resulting rotation periods, with Set  $\mathcal{A}$  shown on top and Set  $\mathcal{B}$  on the bottom. While Set  $\mathcal{A}$  is more complete than Set  $\mathcal{B}$ , this completeness comes at the expense of greater contamination. For instance, stars above the slow sequence (perhaps field contaminants) mostly disappear in Set  $\mathcal{B}$ . The “void” or “gap” underneath the slow sequence is also more pronounced in Set  $\mathcal{B}$ . To facilitate a comparison against the Pleiades and Praesepe, we used the rotation periods and dereddened colors from Table 5 of Curtis et al. (2020), which drew on data from Rebull et al. (2016) and Douglas et al. (2019) respectively. The first order conclusion is that the Pleiades and NGC 2516 appear gyrochronologically coeval for colors spanning  $0.5 < (G_{\text{BP}} - G_{\text{RP}})_0 < 1.2$ . The implications of this comparison for the age of NGC 2516 are discussed in Section 4.4.

Figure 5 shows the same data, with the points colored by indicators of binarity. To assess astrometric binarity, we used the renormalized unit weight error (RUWE) from Gaia EDR3. For plotting purposes, we labeled anything with RUWE exceeding 1.2 as an astrometric binary (11% of the overall cluster sample; see *e.g.*, Belokurov et al. 2020). To assess photometric binarity, we fitted a spline to Figure 3, fixing the nodes by hand. We then labeled any points brighter than 0.3 mag above the cluster sequence as photometric binaries. This included 20% of the overall cluster sample. Many of the resulting candidate binaries, though not all of them, appear below the slow rotation sequence. In Set  $\mathcal{B}$ , over the color range where the slow and fast sequence are present ( $0.5 < (G_{\text{BP}} - G_{\text{RP}})_0 < 1.3$ ), there are 281 stars, of which 103 (37%) show signs of binarity. Dividing these 281 stars into “fast” and “slow” sequences by eye, the fraction of stars showing signs of binarity in the slow and fast subsamples are 30% (63/210) and 56% (40/71), respectively. We speculate on possible explanations in Section 4.2.

### 3.2.3. Comparing Kinematics and Rotation

The TESS rotation periods provide an independent check on the Gaia-derived kinematic cluster memberships. We explored this by cross-matching the stars with detected TESS rotation periods against our original target list of 3,298 Gaia members.

To visualize the results, we again opted for Cartesian coordinates, and supplemented them by calculating the tangential stellar velocities relative to the cluster center. Appendix D discusses the projection effect correction required for calculating the tangential velocities: for a star at position  $(\alpha, \delta)$ , we compare the observed proper motion  $(\mu_\alpha, \mu_\delta)$  with what the proper motion at the star’s position would have been if the star were comoving with the core of NGC 2516. This yields a quantity we denote  $\Delta v^*$ , per M21. We convert these proper motion differences to physical units by multiplying by the measured parallax.

The results are shown in the top two rows of Figure 6. In these figures, to ensure a fair comparison, we only show stars with  $0.5 < (G_{\text{BP}} - G_{\text{RP}})_0 < 1.2$  for which our TESS pipeline succeeded in making light curves. In other words, the stars in the base sample are those for which we could have plausibly detected a rotation period. This color range is preferred because the slow sequence becomes less defined for spectral types later than  $\approx$ K4V. The bins in the histograms (top-right) were chosen to ensure that the same number of stars were in each bin. Rotation periods are detected for 66% of stars at least 25 pc from the cluster core. The average detection rate inside of 25 pc, 73%, is slightly higher. Within  $\approx$ 3.5 pc, the period detection rate of 61% is relatively low, due to crowding and saturation in the TESS images.


The lower panels of Figure 6 give a second view of how field star contamination affects the outskirts of the cluster. Of the outermost 100 stars in position space with detected rotation periods,  $\approx$ 8 appear as outliers above the slow sequence, compared to just a few for the innermost 100 stars. For the outermost 100 stars in velocity space,  $\approx$ 13 appear as outliers. After accounting for the stars that were expected to show rotation periods and did not, this suggests field contaminant rates of  $\approx$ 50% for the outermost cluster members in our target list.

Even with this level of contamination, our rotation period measurements show that the halo extends to separations of  $\approx$ 250 pc in physical space from the cluster core. The most widely separated F2V-K2V stars with rotation periods (blue points in Figure 6) are separated by  $\approx$ 430 pc. The total length of the structure is therefore 400-500 pc, depending on which members of the halo are chosen as the “tips” on either end. This agrees with the overall structure of the halo reported by KC19, with some minor caveats (orange circles) visible in the top-left panels of Figure 6.

In projection-corrected tangential velocity space, the fraction of stars with rotation period detections remains high out to roughly  $5 \text{ km s}^{-1}$ . Meingast et al. (2021) by comparison required a physically motivated cut in tangential velocity space of  $1.5 \text{ km s}^{-1}$ . Our results show that at the expense of higher field star contamination rates, bona fide members can be identified even out at higher velocity separations.

### 3.3. Lithium from Gaia-ESO and GALAH

The final approach we took for assessing the youth of the halo population of NGC 2516 was an analysis of the Li I 6708 Å doublet. This type of age measurement has been re-



**Figure 5. Binarity indicators for rotators in NGC 2516.** Sets  $\mathcal{A}$  (top) and  $\mathcal{B}$  (bottom) are as in Figure 4. Astrometric binaries ( $\text{RUWE} > 1.2$ ; 11% of stars) appear in red; photometric binaries ( $> 0.3$  mag above an empirical isochrone; 20% of stars) are overplotted in orange. Over  $0.5 < (G_{BP} - G_{RP})_0 < 1.3$ , Set  $\mathcal{B}$  has 281 stars. Dividing these stars into “slow” and “fast” sequences by eye, the fraction of stars showing signs of binarity in the slow and fast subsamples are 30% (63/210) and 56% (40/71) respectively.

495 viewed by Soderblom (2010), and its connection to stellar ro-  
 496 tation has been discussed by Bouvier (2020). For NGC 2516,  
 497 two spectroscopic datasets seemed important: Gaia-ESO  
 498 (Gilmore et al. 2012) and GALAH (Silva et al. 2015). At the  
 499 time of analysis, Gaia-ESO DR4 and GALAH DR3 were the  
 500 most relevant (Randich et al. 2018; Buder et al. 2020). The  
 501 target selection and results from each survey were as follows.  
 502 The Gaia-ESO collaboration chose candidate NGC 2516  
 503 members to observe with GIRAFFE and UVES based on pre-  
 504 viously reported literature members and publicly available  
 505 photometry. Since the existence of the NGC 2516 halo was  
 506 not known at the time of target selection, very few halo stars  
 507 are in the sample. Based on the data, Randich et al. (2018)  
 508 determined stellar parameters (including lithium equivalent  
 509 widths and metallicities) for 796 stars that they considered  
 510 possible NGC 2516 members. Cross-matching these against  
 511 our kinematic list of 3,298 candidate members by position  
 512 and imposing a  $0.^{\circ}5$  maximum separation limit yielded 459  
 513 kinematic members with available spectra, 417 in the core  
 514 and 42 in the halo of NGC 2516. The lop-sided ratio is due  
 515 to the Gaia-ESO selection function. An inspection of the ef-  
 516 fective temperature vs. lithium equivalent width based on Ta-  
 517 ble 2 of Randich et al. (2018) showed the surprising feature  
 518 that stars with  $T_{\text{eff}} \lesssim 4000\text{K}$  often had non-zero equivalent  
 519 widths, despite the expectation that their lithium should be  
 520 fully depleted. Whether these measurements were significant  
 521 was not clear, and so we opted to download the spectra from  
 522 the ESO archive to re-measure equivalent widths.

523 The GALAH DR3 target selection is discussed by Buder  
 524 et al. (2020). The relevant aspects for our analysis are that  
 525 the survey targeted  $12 < V < 14$  stars at  $\delta < +10^{\circ}$  in select  
 526 fields, provided the stars were at least ten degrees from the  
 527 galactic plane. We identified the candidate NGC 2516 mem-  
 528 bers for which spectra had been obtained by searching the  
 529 GALAH\_DR3\_main\_allstar\_v1 catalog, after excluding  
 530 stars with the stellar parameter bit flags 1, 2, 3. This  
 531 excludes spectra with unreliable broadening, low S/N, and  
 532 unreliable wavelength solutions (see Table 6 of Buder et al.  
 533 2020). Of our 3,298 candidate NGC 2516 members, 107 had  
 534 spectra in GALAH DR3. 51 were in the core, and 56 were in  
 535 the halo. We downloaded<sup>7</sup> the GALAH DR3 spectra for all  
 536 107 entries.

537 We measured the lithium equivalent widths using the  
 538 specutils package (Earl et al. 2020). We did this by  
 539 first shifting the spectra to the stellar rest frame, and then  
 540 focused in on a  $15\text{\AA}$  window centered on the  $6707.835\text{\AA}$   
 541 lithium doublet. We continuum normalized the spectra with  
 542 a third-degree Chebyshev series, excluding any regions that  
 543 showed absorption lines (e.g., Fe I is present at  $6703.58\text{\AA}$   
 544 and  $6705.10\text{\AA}$ ). We proceeded by fitting a Gaussian to the  
 545 continuum normalized spectra, considering only a  $\pm 1\text{\AA}$  win-  
 546 dow centered on the Li doublet. The equivalent widths were  
 547 then evaluated by numerically integrating the fitted model

548 over the same window. Our approach therefore includes the  
 549 Fe  $6707.44\text{\AA}$  blend in the reported Li equivalent widths—  
 550 which leads to systematic overestimates of 10 to  $15\text{ m\AA}$  (e.g.,  
 551 Bouvier et al. 2018). Finally, to derive uncertainties on the  
 552 EWs, we repeated the procedure twenty times, but added  
 553 noise to the spectra, drawn from a normal distribution with  
 554 a scale set by the standard deviation of the continuum. The  
 555 reported uncertainties are then drawn from the 84<sup>th</sup> and 16<sup>th</sup>  
 556 percentiles of the resulting EW distribution, with minima im-  
 557 posed at  $5\text{ m\AA}$  for FGK stars, and  $20\text{ m\AA}$  for M dwarfs. They  
 558 are included in Table 1. We verified the overall scale of  
 559 our results by comparing our measurements with those of  
 560 Randich et al. (2018). Most stars with  $T_{\text{eff}} > 4500\text{K}$  fol-  
 561 lowed a 1-1 relation. For cooler stars, our EW measure-  
 562 ments are systematically lower, as expected given the deple-  
 563 tion timescales for stars at these ages (e.g., Soderblom et al.  
 564 2014). We visually verified from the spectra that, for the most  
 565 part, we prefer our measurements.


566 Figure 7 shows the concatenation of the Gaia-ESO and  
 567 GALAH results, with the points colored according to  
 568 whether the stars are in the core or the halo of the cluster. The  
 569 GALAH spectra span a color range of  $0 < (G_{\text{BP}} - G_{\text{RP}})_0 <$   
 570 1.5, due to the  $V < 14$  brightness cutoff of the survey. The  
 571 overall increase of Li EW from  $0 < (G_{\text{BP}} - G_{\text{RP}})_0 < 1.0$  is  
 572 driven by the temperatures in the stellar chromospheres: hot-  
 573 ter stars fully ionize Li out of its ground state (e.g., Figure 4  
 574 of Soderblom et al. 1993b). The depletion of stars redder than  
 575  $(G_{\text{BP}} - G_{\text{RP}})_0 \gtrsim 1.4$ , i.e., later than K4.5V ( $0.71 M_{\odot}$ ), is vis-  
 576 ible relative to the earlier K dwarfs: these stars have burned  
 577 their lithium. For comparison, we have also plotted the Li  
 578 EWs measured by Berger et al. (2018) for planet-hosting  
 579 stars in the Kepler field, and Pleiades members from Bouvier  
 580 et al. (2018) (we interpolated from their effective tempera-  
 581 tures using the Pecaut & Mamajek 2013 table). For clarity,  
 582 we have omitted the few upper limits reported by Bouvier  
 583 et al. (2018). The field star distribution peaks at late F-type  
 584 stars. The majority of kinematically selected stars in the core  
 585 and the halo show lithium equivalent widths substantially in  
 586 excess of these field stars.

587 We can go one step further, and match our sample of  
 588 kinematically selected stars with spectra against the TESS  
 589 rotators. The result is shown in Figure 8. At fixed stel-  
 590 lar mass, the rapid rotators have lithium equivalent widths  
 591 an order of magnitude larger than the slow rotators. This  
 592 effect is mostly apparent in the K dwarfs. Similar trends  
 593 were noted in the Pleiades over three decades ago (Butler  
 594 et al. 1987), and are thought to be caused by differences in  
 595 lithium abundance (Soderblom et al. 1993b). Alternative ex-  
 596 planations, such as differences in line-formation conditions  
 597 (e.g., chromospheric temperatures, microturbulent velocities,  
 598 or the presence of starspots) are incompatible with the avail-  
 599 able data. The lithium-rotation correlation is discussed fur-  
 600 ther below (Section 4.3).


## 4. DISCUSSION

### 4.1. How did the halo form?

<sup>7</sup> Via [datacentral.org.au/services/download](http://datacentral.org.au/services/download), using the `sobject_id` identifiers.



**Figure 6.** NGC 2516 members exist hundreds of parsecs from the core, and up to  $\approx 5 \text{ km s}^{-1}$  from the core in tangential velocity. *Top left:* Cartesian positions (as in Figure 2) and 2D-tangential velocities for halo members. Stars with detected rotation periods in Set  $\mathcal{A}$  are shown in blue; halo members for which rotation periods should have been detectable, but were not detected, are shown in orange. Stars in the core are shown as black points. In the *top right*, the ratio of detected to expected stars with rotation periods is shown versus 3D separation from the core and 2D tangential velocity difference. Bin widths are chosen to enforce the same number of stars in the denominator ( $N = 54$ ) per bin. The *bottom panels* show the innermost and outermost 100 stars with detected rotation periods in position and velocity space.



**Figure 7. Lithium in the core and halo of NGC 2516.** Equivalent width of the 6708 Å doublet is shown versus dereddened color for all candidate NGC 2516 members with Gaia-ESO ( $N = 459$ ) or GALAH ( $N = 107$ ) spectra available. The GALAH spectra comprise slightly over half of the halo stars, due to the non-targeted selection function of that survey. Gray points are field stars with Kepler planets (Berger et al. 2018). Pleiades stars (125 Myr) with Li detections are from Bouvier et al. (2018); Praesepe detections (650 Myr) are shown in yellow (Soderblom et al. 1993a). Points with  $\text{EW} \approx 0 \text{ m}\text{\AA}$  are non-detections; all stars later than  $\approx \text{M}0$  do not have significant lithium. The Li measurements for both core and halo stars in NGC 2516 are consistent with a near-Pleiades age.


Any theory for the structure of NGC 2516, the Psc-Eri stream, and the strings and halos found by KC19 and M21 needs to explain a few observations. First is their aspect ratios. The halo of NGC 2516, for instance, extends over at least 500 pc, despite being only  $\approx 25$  pc wide. Second is their axial tilts in the plane of the Milky Way – why their leading and trailing arms tend to conform to the Galaxy’s differential rotation (e.g., Figure 2 or Figure 13 of M21). Third is the correlation between the mean LSR-subtracted cluster velocity and elongation axes (blue arrows in Figure 2). Such a correlation has also been noted in Coma Ber by Tang et al. (2019); the other halos may show similar trends.

The most likely explanation seems to be that the observed halo structures are tidal tails (e.g., Chumak & Rastorguev 2006a; Krumholz et al. 2019). The idea is that stars near the cluster’s tidal radius escape slowly due to the galactic tide, and subsequently form leading and lagging arms due to differential rotation in the Galaxy. The contraction rate of the cluster can affect this process. Whether the exact contraction rate of NGC 2516 (Healy & McCullough 2020), and the corresponding evaporation rate of the core are sufficient to explain the formation of the halo has yet to be assessed.

A second possibility is that the clusters form in larger and more dispersed star formation complexes: the stars in the halo need not have formed in the same “clump” as those in the cluster core. In other words, the formation environ-

ment might be a giant molecular filament, rather than a giant molecular cloud. A sample of such filaments has been collected and analyzed by Zucker et al. (2018): if the current  $\approx 20:1$  aspect ratio of NGC 2516 were primordial, its structure would match the aspect ratios of what they term either “elongated dense core complexes”, or “bone candidates”. A more immediate example is the Orion A cloud. Großschedl et al. (2018) showed using young stellar objects as tracers that Orion A is 90 pc long, and that it has a dense “head” and a lower density “tail”.

Any differences in the stellar mass function between the core and halo could be informative, since the tidal tail explanation predicts that the mean tail star should be less massive than the mean core star (e.g., Chumak & Rastorguev 2006a). Figure 9 shows the histogram of  $A_G$ -corrected absolute magnitudes, with the corresponding Pecaut & Mamajek (2013) mass estimates. The stars closest to the cluster center do seem to have a greater proportion of  $\gtrsim 0.8 M_\odot$  stars, while the outermost stars have a greater proportion of  $\lesssim 0.4 M_\odot$  stars. The mean star in the inner separation bin has  $(M_G)_0 = 7.08$  ( $\approx 0.69 M_\odot$ ), while in the outer bin the mean star has  $(M_G)_0 = 7.61$  ( $\approx 0.63 M_\odot$ ). While the difference does appear significant given the number of stars involved, the theoretical implications are not entirely clear, since the “molecular filament” explanation might be able to accommodate mass differences between the core and tail through density gradi-




**Figure 8. Lithium and rotation in NGC 2516.** Points show all candidate NGC 2516 members with TESS rotation periods and Gaia-ESO or GALAH spectra, and are colored by equivalent width. Points with reported equivalent widths below  $20\text{ m}\text{\AA}$  are shown with gray crosses. The lithium-rotation correlation for the K-dwarfs has been observed in other young clusters, and is discussed in Section 4.3.

ents across the initial filament. Observational incompleteness at the low-mass end seems unlikely to alter the result, since the innermost and outermost regions are subject to similar selection effects.

One possible approach for distinguishing the two explanations could be to observationally establish an age-size correlation. If the  $\approx 100$  Myr halos are produced primarily through evaporation and expansion, then the younger clusters should be significantly smaller (Chumak & Rastorguev 2006a,b). A theoretical approach along complementary lines would be to perform a dynamical analysis of whether filaments like Orion A, or those in the Zucker et al. (2018) sample are at all expected to evolve into structures resembling NGC 2516.


#### 4.2. Stellar Rotation at 100-200 Myr

From their rotation period analysis of NGC 2516, Fritzewski et al. (2020) demonstrated that between 100 and 200 Myr, young clusters (NGC 2516, Blanco 1, Psc-Eri, M 35, M 50, and the Pleiades) all show similar rotation period distributions. The FGK stars show a “slow sequence”, a population of “rapid rotators”, and in some cases stars in the “void” between the two populations (e.g., Psc-Eri and NGC 2516 show more stars in the void than the Pleiades). The late M dwarfs with detected rotation periods tend to rotate rapidly ( $< 3$  days), though there may also be late M dwarfs with slow ( $> 20$  days) rotation periods (Stauffer et al. 2016). Although these late M dwarfs are beyond the magnitude limit of our analysis, they have been explored in depth by Irwin et al. (2007) and Fritzewski et al. (2020). A compar-



**Figure 9. Masses and absolute magnitudes for the innermost and outermost stars in NGC 2516.** The 3,298 candidate NGC 2516 members were split into three groups according to their distance from the cluster center. The inner and outermost groups, each containing 1,099 stars, are counted in 1 mag bins. The mean star in the innermost sample has a 10% lower mass than the mean star in the outermost sample, consistent with expectations for tidal tails.

ison of our results is shown in Figure 10. The bottom panel



**Figure 10. Comparison against previous analyses of stellar rotation in NGC 2516.** The bottom panel is a concatenation of Set  $\mathcal{B}$  with the results of Irwin et al. (2007) (I07) and Fritzewski et al. (2020) (F20). I07 and F20 analyzed stars in the core of NGC 2516.

of this plot includes, in order of precedence: rotation periods from our analysis, from “Class 1 and 2” Fritzewski et al. (2020) rotators, and from the rotators in Irwin et al. (2007). The main contribution of our study beyond these previous analyses is our sensitivity across both the core and halo of NGC 2516, and a corresponding expansion in the number of stars that can be analyzed. In the following section, we therefore focus on the origins of the features in the rotation-color diagram, and whether our expanded sample can provide any new insights.

The existence of a slow sequence is the main basis of gyrochronology (*e.g.*, Barnes 2003). While the spindown relation ( $P_{\text{rot}} \sim t^{1/2}$ ) proposed by Skumanich (1972) is not an accurate enough description, the general idea is correct, and the first-order effect can be understood through magnetic braking (Weber & Davis 1967).

The other features of the rotation-color diagram are indications that magnetic braking is not always the only important effect. The question of why the rapid rotators exist, for instance, requires additional physics. Models that vary core-envelope decoupling timescales and pre-MS disk lifetimes can reproduce each type of population (Irwin et al. 2007; Gallet & Bouvier 2013, 2015); the question then shifts to why these timescales should differ for stars that are otherwise identical. Another possibility is that proposed by Matt et al. (2015): the saturation of the magnetic dynamo yields  $dP/dt \sim P$  when  $P > P_{\text{sat}}$ ,  $dP/dt \sim P^{-1}$  for  $P < P_{\text{sat}}$ , and a maximal spin-down rate in the intermediate regime. This could naturally produce the period bimodality observed at ages of  $\approx 100$  Myr (though it fails to explain the widening of the distribution observed at later times Godoy-Rivera et al. 2021). Other significant processes could be external to the

star entirely: Qureshi et al. (2018) for instance have reported that mergers between giant planets and the star could explain a non-negligible fraction of the rapid rotators.

A separate hypothesis that we can test using the existing data is the idea that rapid rotation can be explained through binarity. Correlations between rapid rotation and binarity in both young clusters and the field have previously been noted (Meibom et al. 2007; Stauffer et al. 2016; Simonian et al. 2019; Gillen et al. 2020). Related effects could include tidal synchronization for the shortest period binaries, or alternatively pre-MS magnetic locking between the disk and star (e.g., Koenigl 1991; Long et al. 2005).

The lower panels of Figure 4 show our attempt at identifying unresolved binaries in the NGC 2516 rotation sample. Binaries resolved by Gaia were already excluded in our initial definitions of the samples. In NGC 2516 we see that the fraction of stars showing signs of binarity in the slow and fast rotation subsamples are 30% (63/210) and 56% (40/71); the fast rotators show a preference for binarity. This correlation is in line with the earlier findings noted above. While many rapid rotators in NGC 2516 (34%) have no evidence for being binaries, the cuts we used to select binaries ( $\text{RUWE} > 1.2$ ; photometric excess  $> 0.3$  mag) leave open a significant fraction of parameter space. A 0.3 mag flux excess, for instance, corresponds to mass ratios  $M_2/M_1 \gtrsim 0.700$ , assuming the luminosity  $L$  scales as  $M^{3.5}$ .

Comparing the explanations of disk-locking against tidal synchronization, the population statistics seem to rule out tidal synchronization. A quarter of the NGC 2516 members are rapidly rotating. In comparison, in the field half of Sun-like stars are binaries, and only  $\approx 9\%$  of these binaries have periods below 100 days (Raghavan et al. 2010). If we assume that all binaries with sub-100 day periods become tidally synchronized, then we can only explain a rapid rotator occurrence rate of  $\approx 5\%$ . Elevated binarity fractions in pre-MS stars likely do not change this picture (see Section 4.4 of Duchêne & Kraus 2013).

A separate effect is that on the slow rotation sequence, Figure 4 shows that the binary stars appear to be either preferentially redder, or to have faster rotation periods than the single stars. One likely explanation for this could be that the unresolved binaries have a component contributing additional red light to the system, skewing the color measurement of the primary. Whether any physical effects could be at play remains a question for future work.

#### 4.3. The lithium-rotation correlation

The lithium-rotation correlation (Figure 8) carries over beyond just equivalent widths, and to the actual abundance of lithium in the photosphere (Soderblom et al. 1993b). Non-LTE corrections do not change the overall relationships between Li abundance and stellar temperature or rotation (Carlsson et al. 1994; Lind et al. 2009). The correlation has been seen in the Pleiades, the Psc-Eri stream, and M 35, among other clusters (Bouvier et al. 2018; Arancibia et al. 2020; Jeffries et al. 2020; Hawkins et al. 2020). In NGC 2516, Jeffries et al. (1998) tentatively observed it

as well, when analyzing 24 stars in the core of NGC 2516. Our concatenation of the Gaia-ESO and GALAH-DR3 spectra represents a significant expansion in volume and color range from this earlier analysis.

What causes the correlation between lithium abundance and rotation at fixed stellar mass? Hints likely lie in shared correlations between lithium abundance, rotation rate, radius inflation, internal mixing, and magnetic field strength (Chabrier et al. 2007; Somers & Stassun 2017; Jeffries et al. 2020). For instance, one interpretation of the lithium-rotation correlation based on pure hydrodynamics is that the rapid rotation leads to less efficient convective penetration (“overshoot”), which lowers the mixing efficiency at the convective-radiative boundary (Baraffe et al. 2017). The magnetic field itself may even be sufficient to inhibit the convection (Ventura et al. 1998). Alternatively, the star’s temperature and density profiles could be at fault: rapid rotators have the strongest magnetic fields and the most spotted surfaces. These spots lower the photospheric flux, and drive the stellar radius to expand while lowering the core temperatures, which in turn slows the rate of lithium-destroying proton capture reactions (Feiden & Chaboyer 2013; Somers & Pinsonneault 2015).

None of the internal processes noted above answer the initial question of why some of the G and K dwarfs rotate rapidly to begin with. One clue could be that the rapid rotators tend to be in photometric or astrometric binaries. The majority of such binaries have wide ( $a \gtrsim 200$  au) or intermediate (0.5 - 100 au) separations (Raghavan et al. 2010).

It is expected that the circumstellar disks in these binaries have different properties than those of single stars, due to e.g., gap formation in the circumbinary disk, disk truncation, and faster disk clearing (Artymowicz & Lubow 1994; Moe & Kratter 2020). It would therefore be reasonable to assume that disk lifetimes in the binary systems are shorter than in single star systems. Eggenberger et al. (2012) showed that longer disk lifetimes should lead to an extended phase of disk-locking, which leads to a greater amount of differential rotation generated in the radiative zone, more efficient shear mixing, and consequently a lower photospheric lithium abundance. This provides a plausible explanation for the overall trend that the rapid rotators are often in unresolved binaries, and that they are also lithium rich. A careful observational analysis including an assessment of the completeness to different separations and mass ratios of binaries would be useful to clarify whether or not all of the rapid rotators are in binaries. If not, this scenario would be falsified.

#### 4.4. The age of NGC 2516

*Age from Gyrochronology*—Comparing the slow sequence of the Pleiades and NGC 2516 more closely, the top row of Figure 4 shows that the two sequences overlap from  $0.5 < (G_{\text{BP}} - G_{\text{RP}})_0 < 1.2$ . At redder colors,  $1.2 < (G_{\text{BP}} - G_{\text{RP}})_0 < 1.7$ , the dispersion in rotation periods increases. The maximum rotation periods seen in NGC 2516 across both Sets  $\mathcal{A}$  and  $\mathcal{B}$  extend up to  $\approx 11$  days, rather than the  $\approx 8.5$  day upper limit

seen in the Pleiades. This is consistent with NGC 2516 being slightly older than the Pleiades.

Fitting a model to substantiate the claim that NGC 2516 is gyrochronologically older than the Pleiades (*e.g.*, Majek & Hillenbrand 2008; Angus et al. 2019; Spada & Lanzaflame 2020) is, unfortunately, an exercise in tautology. Gyrochronological models are empirically calibrated against the Pleiades and Praesepe. At  $(G_{\text{BP}} - G_{\text{RP}})_0 = 1.35$ , we observe rotation periods in NGC 2516 ranging from 7 to 10 days. The 7 day rotation periods are consistent with the Pleiades; the 10 day rotation periods are not. Fitting formulae from the previously cited studies give ages for a star with  $\{(G_{\text{BP}} - G_{\text{RP}})_0 = 1.35, (B - V)_0 = 1.10, P = 7 \text{ day}\}$  of 204 Myr, 316 Myr, and 107 Myr respectively. Given a Pleiades age of  $125 \pm 20 \text{ Myr}$  (an average of MSTO and rotation-corrected lithium-depletion results; see *e.g.* Stauffer et al. 1998, Soderblom et al. 2014, and Cummings & Kalirai 2018), only the Spada & Lanzaflame (2020) model has an absolute scale that seems to be well-calibrated at these ages and colors. For an 8 day rotation period of a star with the same color, this model quotes a 193 Myr age; at 9 days, 388 Myr. Given this degree of model uncertainty, we prefer the relative statement that NGC 2516 appears slightly gyrochronologically older than the Pleiades, and much younger than Praesepe. A gyrochronological age estimate for NGC 2516 between 150 and 200 Myr would therefore appear reasonable. Much older, and the issue of why the F and G dwarfs do not spin down becomes pressing. We caution however that the spindown rate of FGK stars between 100 and 300 Myr has not yet been empirically calibrated, and if “stalling” were to occur, it could affect our age assessment (see Curtis et al. 2020).

*Age from Lithium Depletion*—The lithium depletion boundary for NGC 2516 has to our knowledge not been measured. Given the distance to the cluster, this is not surprising: at  $\sim 150 \text{ Myr}$ , the LDB is expected to occur at a stellar mass of  $\approx 0.088 M_{\odot}$  (Soderblom et al. 2014). This would correspond to  $(G_{\text{BP}} - G_{\text{RP}})_0 \approx 4.7$ , well beyond the limits of available cluster membership lists and spectroscopy.

An easier *relative* measurement to make of lithium depletion is from the EW-color diagram (Figure 7). The NGC 2516 and Pleiades sequences in this diagram appear indistinguishable. A relative lithium-based age estimate for NGC 2516 would therefore be “Pleiades-age”. The uncertainties on this estimate are large because lithium depletion timescales are long for Sun-like stars past 100 Myr. For example, NGC 2516 FGK stars are on average more depleted than stars in IC 2602 ( $\approx 40 \text{ Myr}$ ), but only marginally so (Soderblom et al. 2014). On the older end though, Figure 7 does show that relative to Praesepe ( $\approx 650 \text{ Myr}$ ), the lithium EWs of NGC 2516 are significantly enhanced. Given these comparisons, a plausible range of ages based on lithium for NGC 2516 would therefore be between  $\approx 100$  and  $200 \text{ Myr}$ .

*Adopted Age*—As noted in Section 3.1, a color-color and color-magnitude analysis of the main sequence turnoff (MSTO) by Cummings & Kalirai (2018) found that

NGC 2516 is 40-60 Myr older than the Pleiades. Given a Pleiades age of  $125 \pm 20 \text{ Myr}$ , this yields a MSTO age for NGC 2516 of  $175 \pm 35 \text{ Myr}$ . The gyrochronological age we have argued is likely older than the Pleiades, *i.e.*, in the range of 150 to 200 Myr. The lithium age based on the depletion of the G and K dwarfs is more uncertain, but is consistent with the Pleiades and could be a bit older ( $\approx 100$  to 200 Myr). Averaging these three different indicators gives an age for NGC 2516 of  $167 \pm 20 \text{ Myr}$ , if we set the uncertainties to match the absolute Pleiades age uncertainty of  $\pm 20 \text{ Myr}$  (Soderblom et al. 2014). Whether this average age is at all meaningful, we leave the reader to judge.

## 5. CONCLUSION

The combination of astrometry from Gaia, photometry from TESS, and spectroscopy from GALAH and Gaia-ESO has clarified a few things about the structure of NGC 2516.

- *Over-densities observed in the Gaia 3D positions and 2D velocities revealed a halo of stars spanning  $\approx 500 \text{ pc}$  in the plane of the galaxy.* The earliest reference to this halo that we can find is by Kounkel & Covey (2019). The halo is more precisely described as a leading and trailing tail (Figure 2), with the leading edge angled toward the galactic center, relative to the cluster’s orbit in the Galaxy. This is consistent with the direction and amplitude of the Milky Way’s differential rotation ( $\approx -0.2 \text{ deg Myr}^{-1} \text{ kpc}^{-1}$ ), and similar halos/tails have been observed around  $\approx 10$  other nearby open clusters (Meingast et al. 2021).
- *Isochronal, rotational, and lithium dating show that the halo of NGC 2516 is coeval with its core.* In short, all the data are consistent with the halo being real. The Gaia EDR3 data show a main-sequence turnoff that suggest an isochronal age of 150 Myr for both the core and halo (Figure 3). The faintest M dwarfs in the core and halo are also brighter than field stars of the same color. Gyrochronally, stars in NGC 2516 spanning  $0.5 < (G_{\text{BP}} - G_{\text{RP}})_0 < 1.2$  (spectral types F2V-K3V) overlap with the slow sequence of the Pleiades (Figure 4). At redder colors, from  $1.2 < (G_{\text{BP}} - G_{\text{RP}})_0 < 1.7$  (spectral types K3V-K6V), there is a larger dispersion in rotation period. The upper envelope of the rotation period distribution at these redder colors extends to longer rotation periods than in the Pleiades (11 vs. 8 day rotation periods). This is consistent with NGC 2516 being slightly older than the Pleiades (*i.e.*,  $\approx 150 \text{ Myr}$ ). The lithium equivalent widths of the kinematic cluster members (Figure 7) are consistent with this assessment.
- *Bonafide NGC 2516 members exist out to  $\approx 250 \text{ pc}$  in separation and out to  $\approx 5 \text{ km s}^{-1}$  in tangential velocity separation from the cluster core.* Figure 6 shows the overlap between kinematic and rotational cluster members used to make this assessment.

- The field star contamination rate increases at larger physical and velocity separations from the cluster core.

The right and lower panels of Figure 6 quantify the statement:  $\approx 70\%$  of the innermost candidate members have detected rotation periods consistent with a gyrochronological age near that of the Pleiades. At physical separations of  $\approx 200$  pc, this fraction decreases to  $\approx 50\%$ . Given the number density ratios of cluster to field stars at the widest separations ( $\approx 10^{-3}$ , Meingast et al. 2021), these contamination fractions are not as bad as one might fear.

- The average star in the outskirts has lower mass (by  $\approx 10\%$ ) than the average star in the cluster center. This is shown in Figure 9, and discussed in Section 4.1.

While this is consistent with expectations for a “tidal tail” origin of the halo, the possibility of a primordial halo also merits exploration.

- The rapidly rotating K dwarfs show elevated lithium abundances, and elevated binarity fractions.

Each trend has been noted in comparable stellar populations (e.g., Soderblom et al. 1993b; Meibom et al. 2007; Jeffries et al. 2020; Gillen et al. 2020). The rough orders of magnitude of the effects are  $\approx 1$  dex in lithium abundance for a factor of 10 in rotation period, and a factor of  $\gtrsim 2$  enhancement in binarity fraction for fast vs. slow rotators. Section 4.3 discusses one plausible connection between the two observations: the disk-locking phase may be truncated in binary systems, which could free the stars to spin up as they contract on the pre-MS. This spin up could result in less efficient internal mixing and a longer-lasting presence of photospheric lithium (Eggenberger et al. 2012).

The existence of the halo itself is not particularly surprising. A star with a  $1 \text{ km s}^{-1}$  velocity difference from the cluster average will move away from the center at a rate of  $1 \text{ pc Myr}^{-1}$ . For a  $\sim 100$  Myr cluster, a halo with characteristic size  $\sim 100$  pc is therefore expected, given that the typical velocity dispersions of open clusters are of order kilometers per second.

On the other hand, our ability to reliably identify members of these halos is rather surprising. These stars have remained hidden in the background of the Galaxy throughout centuries of modern astronomy. Their discovery represents an important step toward understanding the dispersal of open clusters. Combined with wide-field photometric surveys, the stars themselves enable advances in studies of stellar angular momentum evolution. Combined with wide-field spectroscopic surveys (e.g., Kollmeier et al. 2017), the halos could also improve our understanding of the chemical evolution of low-mass stars. Finally, the halo stars provide an expanded sample of age-dated stars to search for planets (e.g., Newton et al. 2021), which may help in benchmarking our understanding of planetary evolution.

## ACKNOWLEDGMENTS

L.G.B. is grateful to G. Zhou, B. Tofflemire, A. McWilliam, E. Newton, M. Kounkel, A. Kraus, L. Hillenbrand, and K. Hawkins for the discussions on young stars, rotation, and lithium that encouraged this analysis. L.G.B. and J.H. acknowledge support by the TESS GI Program, programs G011103 and G022117, through NASA grants 80NSSC19K0386 and 80NSSC19K1728. L.G.B. was also supported by a Charlotte Elizabeth Procter Fellowship from Princeton University. This study was based in part on observations at Cerro Tololo Inter-American Observatory at NSF’s NOIRLab (NOIRLab Prop. ID 2020A-0146; 2020B-0029 PI: Bouma), which is managed by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation. This paper also includes data collected by the TESS mission, which are publicly available from the Mikulski Archive for Space Telescopes (MAST). Funding for the TESS mission is provided by NASA’s Science Mission directorate. We thank the TESS Architects (G. Ricker, R. Vanderspek, D. Latham, S. Seager, J. Jenkins) and the many TESS team members for their efforts to make the mission a continued success.

**Software:** astrobase (Bhatti et al. 2018), astropy (Astropy Collaboration et al. 2018), astroquery (Ginsburg et al. 2018), cdips-pipeline (Bhatti et al. 2019), corner (Foreman-Mackey 2016), gala (Price-Whelan 2017; Price-Whelan et al. 2017), IPython (Pérez & Granger 2007), matplotlib (Hunter 2007), numpy (Walt et al. 2011), pandas (McKinney 2010), scikit-learn (Pedregosa et al. 2011), scipy (Jones et al. 2001), wotan (Hipke et al. 2019).

**Facilities:** Astrometry: Gaia (Gaia Collaboration et al. 2018, 2020). Imaging: Second Generation Digitized Sky Survey. Spectroscopy: CTIO1.5m (CHIRON; Tokovinin et al. 2013), AAT (HERMES; Lewis et al. 2002; Sheinis et al. 2015), VLT:Kueyen (FLAMES; Pasquini et al. 2002). Photometry: TESS (Ricker et al. 2015).

**Table 1.** Rotation periods and lithium equivalent widths for 3,298 candidate NGC 2516 members.

Parameter	Example Value	Description
source_id	5290665204845694336	Gaia DR2 source identifier.
source_id_edr3	5290665204845694336	Gaia EDR3 source identifier.
in_SetA	True	In Set $\mathcal{A}$ ( $LSP > 0.08$ , $P < 15d$ , $n_{\text{equal}} == 0$ , $n_{\text{close}} >= 1$ ).
in_SetB	True	In Set $\mathcal{B}$ (Set $\mathcal{A}$ and periods match).
n_cdips_sector	7	Number of TESS sectors with CDIPS light curves.
period	0.228981	Lomb-Scargle best period [days].
lspval	0.657826	Lomb-Scargle periodogram value for best period.
spdmpaperiod	0.228981	Stellingwerf PDM best period [days].
spdmpval	0.372779	Stellingwerf PDM periodogram value for best period.
n_equal	0	Number of stars brighter than the target in TESS aperture.
n_close	1	Number of stars with $\Delta T > 1.25$ in TESS aperture.
n_faint	1	Number of stars with $\Delta T > 2.5$ in TESS aperture.
ra	119.234778	Gaia DR2 right ascension [deg].
dec	-61.015929	Gaia DR2 declination [deg].
ref_epoch	2015.5	Reference epoch for right ascension and declination.
parallax	2.458023	Gaia DR2 parallax [mas].
parallax_error	0.028395	Gaia DR2 parallax uncertainty [mas].
pmra	-4.225826	Gaia DR2 proper motion $\mu_\alpha \cos \delta$ [mas yr $^{-1}$ ].
pmdec	10.899938	Gaia DR2 proper motion $\mu_\delta$ [mas yr $^{-1}$ ].
phot_g_mean_mag	15.778953	Gaia DR2 $G$ magnitude.
phot_bp_mean_mag	16.629477	Gaia DR2 $G_{\text{BP}}$ magnitude.
phot_rp_mean_mag	14.85896	Gaia DR2 $G_{\text{RP}}$ magnitude.
radial_velocity	NaN	Gaia DR2 heliocentric radial velocity [km s $^{-1}$ ].
radial_velocity_error	NaN	Gaia DR2 radial velocity uncertainty [km s $^{-1}$ ].
subcluster	core	Is star in core (CG18) or halo (KC19+M21)?
in(CG18)	True	Star in <a href="#">Cantat-Gaudin et al. (2018)</a> .
in(KC19)	True	Star in <a href="#">Kounkel &amp; Covey (2019)</a> .
in(M21)	False	Star in <a href="#">Meingast et al. (2021)</a> .
(Bp-Rp)_0	1.636198	Gaia $G_{\text{BP}} - G_{\text{RP}}$ color, minus $E(G_{\text{BP}} - G_{\text{RP}}) = 0.1343$
is_phot_bin	False	True if $> 0.3$ mag above cluster isochrone.
is_astrm_bin	False	True if Gaia EDR3 RUWE $> 1.2$ .
ruwe	0.986349	Gaia EDR3 RUWE.
Li_EW_mA_GaiaESO	-1.495	Gaia-ESO Li doublet equivalent width, including the Fe blend [mÅ].
Li_EW_mA_perr_GaiaESO	12.407	Gaia-ESO Li doublet EW upper uncertainty [mÅ].
Li_EW_mA_merr_GaiaESO	5.0	Gaia-ESO Li doublet EW lower uncertainty [mÅ].
Li_EW_mA_GALAH	NaN	GALAH Li doublet equivalent width, including the Fe blend [mÅ].
Li_EW_mA_perr_GALAH	NaN	GALAH Li doublet EW upper uncertainty [mÅ].
Li_EW_mA_merr_GALAH	NaN	GALAH Li doublet EW lower uncertainty [mÅ].

NOTE— Table 1 is published in its entirety in a machine-readable format. One entry is shown for guidance regarding form and content. This table is a concatenation of all candidate NGC 2516 members reported by CG18, KC19, and M21 based on the Gaia DR2 data. Different levels of purity and completeness can be achieved using different cuts on photometric periods, periodogram powers, and lithium equivalent widths. Sets  $\mathcal{A}$  and  $\mathcal{B}$  provide two possible levels of cleaning (see Section 3.2.1). When the target star is the only star present in the TESS aperture,  $n_{\text{equal}} = 0$ ,  $n_{\text{close}} = 1$ , and  $n_{\text{faint}} = 1$ . The light curves are available at <https://archive.stsci.edu/hlsp/cdips>. Supplementary plots enabling the analysis individual stars are available at <https://lgbouma.com/notes>.

## REFERENCES

- 1025 Adams, F. C. 2010, *ARA&A*, **48**, 47
- 1026 Angus, R., Morton, T. D., Foreman-Mackey, D., et al. 2019, *AJ*, **158**, 173
- 1027 Arancibia, J., Bouvier, J., Bayo, A., et al. 2020, Boletin de la Asociacion Argentina de Astronomia La Plata Argentina, **61C**, 81
- 1028 Artymowicz, P., & Lubow, S. H. 1994, *ApJ*, **421**, 651
- 1029 Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, **156**, 123
- 1030 Bailey, J. I., Mateo, M., White, R. J., Shectman, S. A., & Crane, J. D. 2018, *MNRAS*, **475**, 1609
- 1031 Baraffe, I., Pratt, J., Goffrey, T., et al. 2017, *The Astrophysical Journal Letters*, **845**, L6
- 1032 Baratella, M., D’Orazi, V., Carraro, G., et al. 2020, *A&A*, **634**, A34
- 1033 Barnes, S. A. 2003, *ApJ*, **586**, 464
- 1034 Belokurov, V., Penoyre, Z., Oh, S., et al. 2020, *MNRAS*, **496**, 1922
- 1035 Berger, T. A., Howard, A. W., & Boesgaard, A. M. 2018, *ApJ*, **855**, 115
- 1036 Bergond, G., Leon, S., & Guibert, J. 2001, *A&A*, **377**, 462
- 1037 Bhatti, W., Bouma, L., & Yee, S. 2019, cdips-pipeline v0.1.0, <https://doi.org/10.5281/zenodo.3370324>
- 1038 Bhatti, W., Bouma, L. G., & Wallace, J. 2018, astrobase, <https://doi.org/10.5281/zenodo.1469822>
- 1039 Bonifacio, P., Monai, S., & Beers, T. C. 2000, *AJ*, **120**, 2065
- 1040 Bouma, L. G., Hartman, J. D., Bhatti, W., Winn, J. N., & Bakos, G. Á. 2019, *ApJS*, **245**, 13
- 1041 Bouma, L. G., Hartman, J. D., Brahm, R., et al. 2020, *AJ*, **160**, 239
- 1042 Bouvier, J. 2020, *Mem. Soc. Astron. Italiana*, **91**, 39
- 1043 Bouvier, J., Barrado, D., Moraux, E., et al. 2018, *A&A*, **613**, A63
- 1044 Bovy, J. 2015, *ApJS*, **216**, 29
- 1045 Bressan, A., Marigo, P., Girardi, L., et al. 2012, *MNRAS*, **427**, 127
- 1046 Buder, S., Sharma, S., Kos, J., et al. 2020, arXiv e-prints, 2011, arXiv:2011.02505
- 1047 Butler, R. P., Cohen, R. D., Duncan, D. K., & Marcy, G. W. 1987, *ApJL*, **319**, L19
- 1048 Campello, R. J. G. B., Moulavi, D., Zimek, A., & Sander, J. 2015, *ACM Transactions on Knowledge Discovery from Data*, **10**, 5:1
- 1049 Cantat-Gaudin, T., Jordi, C., Vallenari, A., et al. 2018, *A&A*, **618**, A93
- 1050 Cantat-Gaudin, T., Jordi, C., Wright, N. J., et al. 2019, *A&A*, **626**, A17
- 1051 Carlsson, M., Rutten, R. J., Bruls, J. H. M. J., & Shchukina, N. G. 1994, *A&A*, **288**, 860
- 1052 Chabrier, G., Gallardo, J., & Baraffe, I. 2007, *A&A*, **472**, L17
- 1053 Chen, Y., Bressan, A., Girardi, L., et al. 2015, *MNRAS*, **452**, 1068
- 1054 Chen, Y., Girardi, L., Bressan, A., et al. 2014, *MNRAS*, **444**, 2525
- 1055 Chen, Y., Girardi, L., Fu, X., et al. 2019, *A&A*, **632**, A105
- 1056 Choi, J., Dotter, A., Conroy, C., et al. 2016, *ApJ*, **823**, 102
- 1057 Chumak, Y. O., & Rastorguev, A. S. 2006a, *Astronomy Letters*, **32**, 157
- 1058 —. 2006b, *Astronomy Letters*, **32**, 446
- 1059 Ciardi, D. R., Crossfield, I. J. M., Feinstein, A. D., et al. 2018, *AJ*, **155**, 10
- 1060 Cummings, J. 2011, PhD thesis, Indiana University
- 1061 Cummings, J. D., & Kalirai, J. S. 2018, *AJ*, **156**, 165
- 1062 Curtis, J. L., Agüeros, M. A., Mamajek, E. E., Wright, J. T., & Cummings, J. D. 2019, *AJ*, **158**, 77
- 1063 Curtis, J. L., Agüeros, M. A., Matt, S. P., et al. 2020, *ApJ*, **904**, 140
- 1064 Davenport, J. R. A., & Sandquist, E. L. 2010, *ApJ*, **711**, 559
- 1065 David, T. J., Petigura, E. A., Luger, R., et al. 2019, *ApJL*, **885**, L12
- 1066 de Zeeuw, P. T., Hoogerwerf, R., de Bruijne, J. H. J., Brown, A. G. A., & Blaauw, A. 1999, *AJ*, **117**, 354
- 1067 Douglas, S. T., Agüeros, M. A., Covey, K. R., & Kraus, A. 2017, *ApJ*, **842**, 83
- 1068 Douglas, S. T., Curtis, J. L., Agüeros, M. A., et al. 2019, *ApJ*, **879**, 100
- 1069 Duchêne, G., & Kraus, A. 2013, *ARA&A*, **51**, 269
- 1070 Earl, N., Tollerud, E., Jones, C., et al. 2020, *astropy/specutils*
- 1071 Eggenberger, P., Haemmerlé, L., Meynet, G., & Maeder, A. 2012, *A&A*, **539**, A70
- 1072 Ester, M., Kriegel, H.-P., Sander, J., & Xu, X. 1996, in Proceedings of the Second International Conference on Knowledge Discovery and Data Mining, KDD’96 (Portland, Oregon: AAAI Press), 226
- 1073 Feiden, G. A., & Chaboyer, B. 2013, *ApJ*, **779**, 183
- 1074 Foreman-Mackey, D. 2016, *Journal of Open Source Software*, **1**, 24
- 1075 Freeman, K., & Bland-Hawthorn, J. 2002, *ARA&A*, **40**, 487
- 1076 Fritzewski, D. J., Barnes, S. A., James, D. J., & Strassmeier, K. G. 2020, *A&A*, **641**, A51
- 1077 Fukushige, T., & Heggie, D. C. 2000, *MNRAS*, **318**, 753
- 1078 Fürnkranz, V., Meingast, S., & Alves, J. 2019, *A&A*, **624**, L11
- 1079 Gagné, J., & Faherty, J. K. 2018, *ApJ*, **862**, 138
- 1080 Gagné, J., Faherty, J. K., & Popinchalk, M. 2020, *Res. Notes AAS*, **4**, 92
- 1081 Gagné, J., Roy-Loubier, O., Faherty, J. K., Doyon, R., & Malo, L. 2018, *ApJ*, **860**, 43
- 1082 Gagné, J., David, T. J., Mamajek, E. E., et al. 2020, *ApJ*, **903**, 96
- 1083 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2020, arXiv e-prints, arXiv:2012.01533
- 1084 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *A&A*, **616**, A1
- 1085 Gallet, F., & Bouvier, J. 2013, *A&A*, **556**, A36
- 1086 —. 2015, *A&A*, **577**, A98
- 1087 Gillen, E., Briegal, J. T., Hodgkin, S. T., et al. 2020, *MNRAS*, **492**, 1008

- 1121 Gilmore, G., Randich, S., Asplund, M., et al. 2012, *The Messenger*,  
1122 147
- 1123 Ginsburg, A., Sipocz, B., Madhura Parikh, et al. 2018,  
1124 *Astropy/Astroquery: V0.3.7 Release*
- 1125 Godoy-Rivera, D., Pinsonneault, M. H., & Rebull, L. M. 2021,  
1126 arXiv:2101.01183 [astro-ph], arXiv: 2101.01183
- 1127 Großschedl, J. E., Alves, J., Meingast, S., et al. 2018, *A&A*, 619,  
1128 A106
- 1129 Hawkins, K., Lucey, M., & Curtis, J. 2020, *MNRAS*, 496, 2422
- 1130 Healy, B. F., & McCullough, P. R. 2020, *ApJ*, 903, 99
- 1131 Hippke, M., David, T. J., Mulders, G. D., & Heller, R. 2019, *AJ*,  
1132 158, 143
- 1133 Hogg, D. W., Casey, A. R., Ness, M., et al. 2016, *ApJ*, 833, 262
- 1134 Hunt, E. L., & Reffert, S. 2020, *A&A*, arXiv: 2012.04267
- 1135 Hunter, J. D. 2007, *Computing in Science & Engineering*, 9, 90
- 1136 Irwin, J., Hodgkin, S., Aigrain, S., et al. 2007, *MNRAS*, 377, 741
- 1137 Jeffries, R. D., Jackson, R. J., Sun, Q., & Deliyannis, C. P. 2020,  
1138 *MNRAS*, 500, 1158
- 1139 Jeffries, R. D., James, D. J., & Thurston, M. R. 1998, *MNRAS*,  
1140 300, 550
- 1141 Jeffries, R. D., Thurston, M. R., & Hambly, N. C. 2001, *A&A*, 375,  
1142 863, number: 3 Publisher: EDP Sciences
- 1143 Jones, E., Oliphant, T., Peterson, P., et al. 2001, Open source  
1144 scientific tools for Python
- 1145 Kharchenko, N. V., Piskunov, A. E., Schilbach, E., Röser, S., &  
1146 Scholz, R.-D. 2013, *A&A*, 558, A53
- 1147 Koenigl, A. 1991, *ApJL*, 370, L39
- 1148 Kollmeier, J. A., Zasowski, G., Rix, H.-W., et al. 2017, arXiv  
1149 e-prints, arXiv:1711.03234
- 1150 Kounkel, M., & Covey, K. 2019, *AJ*, 158, 122
- 1151 Kounkel, M., Covey, K., & Stassun, K. G. 2020, *AJ*, 160, 279
- 1152 Kounkel, M., Covey, K., Suárez, G., et al. 2018, *AJ*, 156, 84
- 1153 Krone-Martins, A., & Moitinho, A. 2014, *A&A*, 561, A57
- 1154 Krumholz, M. R., McKee, C. F., & Bland-Hawthorn, J. 2019,  
1155 *ARA&A*, 57, 227
- 1156 Lada, C. J., & Lada, E. A. 2003, *ARA&A*, 41, 57
- 1157 Lamers, H. J. G. L. M., Baumgardt, H., & Gieles, M. 2010,  
1158 *MNRAS*, 409, 305
- 1159 Lewis, I. J., Cannon, R. D., Taylor, K., et al. 2002, *MNRAS*, 333,  
1160 279
- 1161 Lind, K., Asplund, M., & Barklem, P. S. 2009, *A&A*, 503, 541
- 1162 Livingston, J. H., Dai, F., Hirano, T., et al. 2019, *MNRAS*, 484, 8
- 1163 Lomb, N. R. 1976, *Astrophysics and Space Science*, 39, 447
- 1164 Long, M., Romanova, M. M., & Lovelace, R. V. E. 2005, *ApJ*, 634,  
1165 1214
- 1166 Mamajek, E. E., & Hillenbrand, L. A. 2008, *ApJ*, 687, 1264
- 1167 Mann, A. W., Gaidos, E., & Ansdell, M. 2013, *ApJ*, 779, 188
- 1168 Mann, A. W., Newton, E. R., Rizzuto, A. C., et al. 2016, *AJ*, 152,  
1169 61
- 1170 Marigo, P., Girardi, L., Bressan, A., et al. 2017, *ApJ*, 835, 77
- 1171 Matt, S. P., Brun, A. S., Baraffe, I., Bouvier, J., & Chabrier, G.  
1172 2015, *ApJL*, 799, L23
- 1173 McInnes, L., Healy, J., & Astels, S. 2017, *The Journal of Open  
1174 Source Software*, 2, 205
- 1175 McKinney, W. 2010, in *Proceedings of the 9th Python in Science  
1176 Conference*, ed. S. van der Walt & J. Millman, 51
- 1177 Meibom, S., Mathieu, R. D., & Stassun, K. G. 2007, *ApJL*, 665,  
1178 L155
- 1179 Meingast, S., & Alves, J. 2019, *A&A*, 621, L3
- 1180 Meingast, S., Alves, J., & Fürnkranz, V. 2019, *A&A*, 622, L13
- 1181 Meingast, S., Alves, J., & Rottensteiner, A. 2021, *A&A*, 645, A84
- 1182 Moe, M., & Kratter, K. M. 2020, arXiv:1912.01699
- 1183 Ness, M., Rix, H.-W., Hogg, D. W., et al. 2018, *ApJ*, 853, 198
- 1184 Newton, E. R., Mann, A. W., Kraus, A. L., et al. 2021, *AJ*, 161, 65
- 1185 O'Donnell, J. E. 1994, *ApJ*, 422, 158
- 1186 Oh, S., Price-Whelan, A. M., Hogg, D. W., Morton, T. D., &  
1187 Spergel, D. N. 2017, *AJ*, 153, 257
- 1188 Pasquini, L., Avila, G., Blecha, A., et al. 2002, *The Messenger*,  
1189 110, 1
- 1190 Pecaut, M. J., & Mamajek, E. E. 2013, *ApJS*, 208, 9
- 1191 Pecaut, M. J., & Mamajek, E. E. 2016, *MNRAS*, 461, 794
- 1192 Pedregosa, F., Varoquaux, G., Gramfort, A., et al. 2011, *Journal of  
1193 Machine Learning Research*, 12, 2825
- 1194 Pérez, F., & Granger, B. E. 2007, *Computing in Science and  
1195 Engineering*, 9, 21
- 1196 Piskunov, A. E., Just, A., Kharchenko, N. V., et al. 2018,  
1197 *Astronomy and Astrophysics*, 614, A22
- 1198 Plavchan, P., Barclay, T., Gagné, J., et al. 2020, *Nature*, 582, 497
- 1199 Price-Whelan, A., Sipocz, B., Major, S., & Oh, S. 2017, *adrn/gala:  
1200 v0.2.1*
- 1201 Price-Whelan, A. M. 2017, *The Journal of Open Source Software*,  
1202 2
- 1203 Qureshi, A., Naoz, S., & Shkolnik, E. L. 2018, *ApJ*, 864, 65
- 1204 Raghavan, D., McAlister, H. A., Henry, T. J., et al. 2010, *ApJS*,  
1205 190, 1
- 1206 Rajpurohit, A. S., Reylé, C., Allard, F., et al. 2013, *A&A*, 556, A15
- 1207 Randich, S., Tognelli, E., Jackson, R., et al. 2018, *A&A*, 612, A99
- 1208 Rebull, L. M., Stauffer, J. R., Bouvier, J., et al. 2016, *AJ*, 152, 113
- 1209 Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, *Journal of  
1210 Astronomical Telescopes, Instruments, and Systems*, 1, 014003
- 1211 Rizzuto, A. C., Newton, E. R., Mann, A. W., et al. 2020,  
1212 arXiv:2005.00013 [astro-ph]
- 1213 Röser, S., Schilbach, E., & Goldman, B. 2019, *A&A*, 621, L2
- 1214 Scargle, J. D. 1982, *ApJ*, 263, 835
- 1215 Schlafly, E. F., & Finkbeiner, D. P. 2011, *ApJ*, 737, 103
- 1216 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- 1217 Sheinis, A., Anguiano, B., Asplund, M., et al. 2015, *Journal of  
1218 Astronomical Telescopes, Instruments, and Systems*, 1, 035002
- 1219 Shu, F. H., Adams, F. C., & Lizano, S. 1987, *ARA&A*, 25, 23

- 1220 Silva, G. M. D., Freeman, K. C., Bland-Hawthorn, J., et al. 2015,  
 1221 [MNRAS, 449, 2604](#)
- 1222 Simonian, G. V. A., Pinsonneault, M. H., & Terndrup, D. M. 2019,  
 1223 [ApJ, 871, 174](#)
- 1224 Skumanich, A. 1972, [ApJ, 171, 565](#)
- 1225 Soderblom, D. R. 2010, [ARA&A, 48, 581](#)
- 1226 Soderblom, D. R., Fedele, S. B., Jones, B. F., Stauffer, J. R., &  
 1227 Prosser, C. F. 1993a, [AJ, 106, 1080](#)
- 1228 Soderblom, D. R., Hillenbrand, L. A., Jeffries, R. D., Mamajek,  
 1229 E. E., & Naylor, T. 2014, [Protostars and Planets VI, 219](#)
- 1230 Soderblom, D. R., Jones, B. F., Balachandran, S., et al. 1993b, [AJ,](#)  
 1231 [106, 1059](#)
- 1232 Somers, G., & Pinsonneault, M. H. 2015, [MNRAS, 449, 4131](#)
- 1233 Somers, G., & Stassun, K. G. 2017, [AJ, 153, 101](#)
- 1234 Spada, F., & Lanzafame, A. C. 2020, [A&A, 636, A76](#)
- 1235 Spitzer, Jr., L. 1958, [ApJ, 127, 17](#)
- 1236 Stassun, K. G., Oelkers, R. J., Paegert, M., et al. 2019, [AJ, 158, 138](#)
- 1237 Stauffer, J., Rebull, L., Bouvier, J., et al. 2016, [AJ, 152, 115](#)
- 1238 Stauffer, J. R., Jones, B. F., Backman, D., et al. 2003, [AJ, 126, 833](#)
- 1239 Stauffer, J. R., Schultz, G., & Kirkpatrick, J. D. 1998, [ApJL, 499,](#)  
 1240 [L199](#)
- 1241 Stellingwerf, R. F. 1978, [ApJ, 224, 953](#)
- 1242 —. 2011, Period Determination of RR Lyrae Stars, in conference:  
 1243 RR Lyrae Stars, Metal-Poor Stars, and the Galaxy Place.  
 1244 arXiv:1108.4984
- 1245 Sullivan, P. W., Winn, J. N., Berta-Thompson, Z. K., et al. 2015,  
 1246 [ApJ, 809, 77](#)
- 1247 Tang, S.-Y., Pang, X., Yuan, Z., et al. 2019, [ApJ, 877, 12](#)
- 1248 Tian, H.-J. 2020, [ApJ, 904, 196](#)
- 1249 Tofflemire, B. M., Rizzuto, A. C., Newton, E. R., et al. 2021, [AJ,](#)  
 1250 [161, 171](#)
- 1251 Tokovinin, A., Fischer, D. A., Bonati, M., et al. 2013, [PASP, 125,](#)  
 1252 [1336](#)
- 1253 Ventura, P., Zeppieri, A., Mazzitelli, I., & D'Antona, F. 1998,  
 1254 [A&A, 331, 1011](#)
- 1255 Wallace, J. J. 2018, [Res. Notes AAS, 2, 213](#)
- 1256 Walt, S. v. d., Colbert, S. C., & Varoquaux, G. 2011, Computing in  
 1257 Science & Engineering, 13, 22
- 1258 Weber, E. J., & Davis, Jr., L. 1967, [ApJ, 148, 217](#)
- 1259 Wright, N. J., & Mamajek, E. E. 2018, [MNRAS, 476, 381](#)
- 1260 Zari, E., Hashemi, H., Brown, A. G. A., Jardine, K., & de Zeeuw,  
 1261 P. T. 2018, [A&A, 620, A172](#)
- 1262 Zhou, G., Quinn, S. N., Irwin, J., et al. 2021, [AJ, 161, 2](#)
- 1263 Zucker, C., Battersby, C., & Goodman, A. 2018, [ApJ, 864, 153](#)
- 1264 Zuckerman, B., & Song, I. 2004, [ARA&A, 42, 685](#)


## A. CLUSTERING METHODS AND OUTCOMES

1267 Figure 11 is a Venn diagram of the three membership catalogs concatenated in this study. 99% of the **CG18** sample overlaps  
 1268 with **KC19**, **M21**, or both. By comparison, 36% of the **KC19** sample, and 15% of the **M21** sample do not overlap with any of  
 1269 the other catalogs. The data—Gaia DR2—used by all the studies was the same. What are the differences in methodology that  
 1270 produce these different membership lists?

1271 **CG18** applied a procedure that yielded what we colloquially call “the core”. Their membership assignment algorithm was to  
 1272 first query a Gaia DR2 cone around the previously reported  $\{\alpha, \delta\}$  of the cluster center, and within  $\pm 0.5$  mas of its previously  
 1273 reported parallax. The outer radius of their cone was the previously reported angular radius of the cluster ( $0.71^\circ$ ; **Kharchenko**  
 1274 et al. 2013). No proper motion cut was applied. **CG18** then applied an unsupervised classification scheme to  $G < 18$  mag stars  
 1275 within this cone (UPMASK; Krone-Martins & Moitinho 2014). The UPMASK algorithm first performs a  $k$ -means clustering  
 1276 on the astrometric parameters,  $\{\mu_{\alpha'}, \mu_{\delta}, \pi\}$ . A “veto” step is then applied to assess whether the groups of stars output from the  
 1277  $k$ -means clustering are more concentrated than a uniform distribution, by comparing the total branch lengths of their respective  
 1278 minimum spanning trees. This binary flag is converted to a membership probability by redrawing values for  $\{\mu_{\alpha'}, \mu_{\delta}, \pi\}$  using  
 1279 the reported uncertainties and covariances. The final membership probability for any given star is then the fraction of draws for  
 1280 which the star passes the veto step. In the case of NGC 2516, the resulting radius within which half of the cluster members were  
 1281 found was  $0.50^\circ$ . We included all **CG18** NGC 2516 members with reported membership probability exceeding 10% in our list of  
 1282 candidate cluster members.

1283 **KC19** applied a different unsupervised clustering method using the 5D Gaia DR2 positions and proper motions. **KC19** began  
 1284 with a sample of  $\approx 2 \times 10^7$  stars, mostly within  $\approx 1$  kpc and with  $G \lesssim 18$  mag. They then applied the hierarchical density-based  
 1285 spatial clustering of applications with noise algorithm (HDBSCAN; Campello et al. 2015, McInnes et al. 2017) on the entire  
 1286 sample, setting the minimum number of stars per cluster to be 40. **KC19** then iterated over a few different cutoff parallaxes to  
 1287 improve sensitivity to structures with distances from the Sun ranging between  $\approx 100$  pc and  $\approx 1000$  pc. They then fitted ages  
 1288 to the resulting clusters using two different methods: isochronal, and a convolutional neural network. Regarding membership  
 1289 probabilities, **KC19** reported the binary “member” or “not” from HDBSCAN. We included all **KC19** NGC 2516 members in our  
 1290 list of candidate cluster members.

1291 Finally, **M21** considered ten known open clusters within the nearest kiloparsec. One of these clusters was NGC 2516. Their  
 1292 initial list of  $\approx 3 \times 10^7$  stars was similar to that of **KC19**, mainly requiring well-behaved astrometry and photometry from Gaia.  
 1293 Given the mean positions and velocities of the clusters determined by **CG18**, **M21** then selected all stars with proper-motion  
 1294 correct tangential velocities within  $1.5 \text{ km s}^{-1}$  of the cluster mean. They then applied DBSCAN (Ester et al. 1996) on both  
 1295 the observed galactic  $\{X, Y, Z\}$  coordinates, as well as to a separate deconvolved spatial coordinate distribution discussed in  
 1296 their Section 3.3. The results of their procedure, as well as those of **CG18** and **KC19**, are visible at the website of S. Meingast:  
 1297 <https://homepage.univie.ac.at/stefan.meingast/coronae.html>. Their reported membership probabilities were binary, similar to those  
 1298 of **KC19**. All of the **M21** NGC 2516 members were included in our list of candidate cluster members.




**Figure 11. Different clustering techniques yield different candidate members of NGC 2516.** 3,298 unique candidate cluster members found using three different techniques are considered in this study. **CG18:** Cantat-Gaudin et al. (2018), **KC19:** Kounkel & Covey (2019), **M21:** Meingast et al. (2021).

1299      **B. CONSIDERATIONS FOR REMOVING SYSTEMATIC VARIABILITY FROM TESS LIGHT CURVES**

1300    In detrending before searching for stellar rotation signals, our goal was to preserve astrophysical variability, while removing  
 1301 systematic variability. One particular concern for the TESS light curves is systematic variability at the timescale of the 14-day  
 1302 satellite orbit, mostly induced by scattered light from the Earth and Moon.

1303    We opted to use a variant of the principal component analysis (PCA) approach discussed by Bouma et al. (2019). This PCA  
 1304 approach uses a set of “trend stars” selected from across each CCD according using ad-hoc heuristics that on average lead the  
 1305 trend star light curves to be dominated by systematic variability. The resulting principal component vectors, also referred to as  
 1306 the eigenvectors, are rank-ordered by the degree of variance that they predict in the training set of trend stars.

1307    We then posit that any given target star’s light curve is described as a linear combination of the eigenvectors. We also con-  
 1308 sidered the inclusion of additional systematic vectors that could affect the light curve, discussed below. These can be treated  
 1309 as additional “features” in the linear model. To determine the coefficients of the linear model after the full set of eigenvectors  
 1310 (plus optional “systematic” vectors) had been assembled, we explored two possible methods: ordinary least squares, and ridge  
 1311 regression. Ridge regression is the same as ordinary least squares, except it includes an  $\ell^2$  norm with a regularization coeffi-  
 1312 cient. The regularization coefficient that best applied for any given target light curve was determined using a cross-validation grid  
 1313 search, through `sklearn.linear_modelRidgeCV` (Pedregosa et al. 2011). Each target light curve was mean-subtracted  
 1314 and normalized by its standard deviation, as were the eigenvectors. The linear problem was then solved numerically, and the light



**Figure 12. NGC 2516 light curves sorted in order of stellar color.** Fifty stars are randomly selected from Set  $\mathcal{B}$  (see Section 3.2.1). The light curves were detrended and cleaned as discussed in Appendix B. Each light curve is separated by 7.5% in relative flux.

1315 curve was reconstructed by re-adding the original mean, and re-multiplying by the standard deviation to ensure that the variance  
1316 of the light curve did not change.

1317 We found that the choice of using ordinary least squares versus ridge regression did not significantly affect the resulting light  
1318 curves. In other words, the inclusion (or lack thereof) of a regularization term did not strongly alter the best-fitting coefficients.  
1319 In the spirit of keeping it simple, we opted for ordinary least squares. A few other choices seemed to be more important:

- 1320 • *To smooth, or to not smooth the eigenvectors.* If the systematic trends are smooth in time, the eigenvectors should be  
1321 as well. They should not contain residuals from *e.g.*, eclipsing binaries in the template set, and they should also not be  
1322 intrinsically noisier than the target star. If either of these is the case, extra variability can be introduced into the “detrended”  
1323 light curves. To address this problem, we opted to smooth the eigenvectors using a sliding time-windowed filter (with a  
1324 “biweight” weight scheme, implemented in `wotan` by Hippke et al. 2019). One issue with this is that systematic sharp  
1325 features (“spikes”) are no longer captured, and could be present in the detrended light curves. Since they can be filtered out  
1326 in postprocessing (*e.g.*, using rolling outlier rejection), we prefer this approach to having systematic features being injected  
1327 by the PCA detrending.
- 1328 • *How many eigenvectors to use.* A larger number always leads to greater whitening. In Bouma et al. (2019), we performed  
1329 a factor analysis cross-validation to determine the number of eigenvectors to use. The typical number adopted based on  
1330 this analysis was 10 to 15. While this approach should in theory prevent over-fitting, in our experience, for stellar rotation  
1331 it still often lead to distorts the signals, especially for rotation signals with small amplitudes and periodicities of  $\gtrsim 3$  days.  
1332 Shorter signals typically are not distorted, since the eigenvectors do not contain the high-frequency content that leads to  
1333 the distortions. For the present analysis, we therefore set the maximum number of eigenvectors to be 5.
- 1334 • *Which supplementary systematics vectors to use.* We considered using the BGV, CCDTEMP, XIC, YIC, and BGV vectors,  
1335 packaged with the CDIPS light curves. We found that the background value measured in an annulus centered on the  
1336 aperture, BGV, tended to produce the best independent information from the PCA eigenvectors, and so we adopted it as our  
1337 only “supplementary” trend vector. We opted to not smooth it, assuming that it would provide direct complement to the  
1338 smoothed PCA vectors.

1339 After all these considerations, for every target star, we ultimately decorrelated the raw (image-subtracted and background-  
1340 subtracted) light curve using a linear model with ordinary least squares. The vectors used in the model were the 1 unsmoothed  
1341 background flux vector, and 5 smoothed PCA eigenvectors. Figure 12 shows fifty light curves randomly drawn from the resulting  
1342 Set  $\mathcal{B}$ . A supplementary figure set that enables inspection of each individual star is available at <https://lgouma.com/notes>.


### 1343 C. ROTATION PERIODS FOR COMPARISON FIELD STARS

1344 To verify the significance of the trends seen in Figures 4 and 5, we searched the CDIPS calibration light curves for rotation  
1345 periods. This light curve database comprises all  $G_{RP} < 13$  stars that fell on TESS silicon. The need for separate field and cluster  
1346 star magnitude limits is based on our processing capabilities and is discussed by Bouma et al. (2019). Over the southern sky  
1347 (Sectors 1-13 of TESS), this corresponded to a sample of 9,619,784 stars. Cross-matching these against the 13,843 randomly  
1348 drawn stars in the neighborhood of NGC 2516 yielded 1,987 unique stars. The magnitude cut of  $G_{RP} < 13$  at the distances of the  
1349 neighborhood sample corresponds to an extinction-corrected color cutoff of  $(G_{BP} - G_{RP})_0 \lesssim 0.80$ , or spectral types  $\lesssim G1V$ . This  
1350 reaches sufficiently far down the slow sequence to enable a comparison against the cluster star sample.


1351 We performed the same light curve stitching and period-search procedure discussed in the text on the field stars. Imposing the  
1352 same requirements for crowding resulted in 820 stars for which rotation periods could have been detected. Imposing the same  
1353 Lomb Scargle power cutoff ( $LSP > 0.08$ ) and period upper limit ( $P < 15$  days) yielded 312 period detections (38.0%). Within  
1354 the same brightness cutoff, 374 of 705 cluster stars yielded period detections (53.1%). While the detection fractions are only  
1355 mildly different (likely due to the brightness cutoff), the period–color distributions are very different (Figure 13). This confirms  
1356 our expectation that our period measurement procedure is not biased in any way that could produce the features we observe in  
1357 NGC 2516.

### 1358 D. PROJECTION EFFECTS FOR TANGENTIAL VELOCITIES

1359 To compare any given star’s proper motion against the cluster’s mean, one must consider the position of the star on the sky.  
1360 This is because two stars sharing identical Cartesian velocities will have different velocities when projected on the sky, as was  
1361 emphasized by Meingast et al. (2021). Figure 14 shows the magnitude of the correction in equatorial tangential velocities.



**Figure 13. Rotation in NGC 2516 compared to the field.** Stars with Gaia  $G_{\text{RP}} < 13$  mag in the cluster and field samples are displayed; fainter field stars were not analyzed. Field stars (orange) show a different rotation period distribution than the kinematically selected NGC 2516 members (black). The field and NGC 2516 stars are both subject to the same “Set  $\mathcal{A}$ ” period-measurement and cleaning procedures.



**Figure 14. Projection effects for co-moving populations across the sky.** The maps are colored by the proper motion difference a star co-moving with NGC 2516 would exhibit across the equatorial sphere. Actual positions of candidate NGC 2516 members are shown with points; the star denotes the cluster center. The *left* and *right* panels show the proper motion correction in the declination and right ascension directions.