

## Ages of Young Stars and Planets in the Kepler Field

LUKE G. BOUMA,<sup>1,\*</sup> LYNNE A. HILLENBRAND,<sup>1</sup> ANDREW W. HOWARD,<sup>1</sup> HOWARD ISAACSON,<sup>2</sup> KENTO MASUDA,<sup>3</sup> AND  
ELSA K. PALUMBO<sup>4</sup>

<sup>1</sup>*Department of Astronomy, MC 249-17, California Institute of Technology, Pasadena, CA 91125, USA*

<sup>2</sup>*Astronomy Department, University of California, Berkeley, CA 94720, USA*

<sup>3</sup>*Department of Earth and Space Science, Osaka University, Osaka 560-0043, Japan*

<sup>4</sup>*Department of Statistics & Data Science, Carnegie Mellon University, Pittsburgh, PA 15213, USA*

(Received —; Revised —; Accepted —)

### ABSTRACT

Recent analyses of FGK stars in open clusters have helped clarify the precision with which a star’s rotation rate and lithium content can be used as empirical indicators for its age. Here we apply this knowledge to stars observed by Kepler. Rotation periods are drawn from previous work; lithium is measured from new and archival Keck/HIRES spectra. We report rotation-based ages for 24,426 stars and 810 planets for which our method is applicable. We find that our rotational ages accurately recover the ages of stars in open clusters spanning 0.04–2.5 Gyr; they also agree with  $\gtrsim 90\%$  of the independent lithium ages. The resulting yield includes 60 planets younger than 1 Gyr at  $2\sigma$ , and 108 with median ages below 1 Gyr. This is about half the number expected under the classic assumption of a uniform star formation history. We find that the scarcity of sub-gigayear systems can be attributed to the star formation rate in the Kepler field dropping by a factor of  $2.85 \pm 0.12$  over the past 3 Gyr. We observe this trend both in known planet hosts and in the parent stellar sample. This “demographic cliff” in the Galaxy’s star formation history has been previously reported, and its confirmation helps clarify the age distribution of the known transiting exoplanets.

**Keywords:** Stellar ages (1581), Planet hosting stars (1242), Field stars (2103), Exoplanet evolution (491), Milky Way evolution (1052)

### 1. INTRODUCTION

Exoplanet science is thriving, fueled by the discovery of thousands of worlds orbiting close to their host stars (Borucki et al. 2010; Ricker et al. 2015). However, most known exoplanets are billions of years old. This fact leaves many gaps in our knowledge of the exact physical and dynamical origins of these objects. The reason is that processes such as thermal cooling (Fortney et al. 2007), atmospheric loss (Owen 2019), giant impacts (Raymond et al. 2014), and dynamical instabilities (Izidoro et al. 2017) are expected to be most efficient over timescales of much less than 1 Gyr. For most known exoplanets, these processes have run their course.

Young ( $< 1$  Gyr) exoplanets represent one means of building the timeline for exoplanet evolution. Informative individual exemplars include HIP 67522b, a Jupiter-sized planet with a sub-Neptunian mass (Rizzuto et al. 2020; Thao et al. submitted), and V1298 Tau, a resonant chain of puffy planets that is a likely precursor to the compact multiplanet

systems (David et al. 2019). Population-level analyses have similarly suggested differences in the size distribution of young exoplanets relative to their older counterparts (Berger et al. 2020a; David et al. 2021; Sandoval et al. 2021; Christiansen et al. 2023; Vach et al. 2024).

Discovering a young planet requires solving two problems: finding the planet and measuring the star’s age. Each problem admits a range of solutions (e.g. Marois et al. 2008; Quinn et al. 2012; Tran et al. 2024). In this article we will consider planets whose existence has been previously established using transits, and infer new stellar ages using rotation and lithium.

To begin, imagine that the ages of nearby stars in the Galaxy are uniformly distributed from 0–10 Gyr (Binney et al. 2000; Nordström et al. 2004). This approximation suggests that  $\approx 1\%$  of nearby stars have  $t < 100$  Myr, and  $\approx 10\%$  have  $t < 1$  Gyr. Studies of currently forming protoplanets (Keppler et al. 2018), and of exoplanets evolving just after disk dispersal (e.g. Klein et al. 2022), are thereby capped in their maximum achievable sample sizes by the tyranny of the galactic star formation rate.

Despite this and other observational challenges, young close-in planet discovery has matured over the past decade, primarily due to Kepler, K2, TESS, and Gaia (e.g. Mei-

Corresponding author: Luke G. Bouma  
bouma.luke@gmail.com

\* 51 Pegasi b Fellow

bom et al. 2013; Mann et al. 2016; Curtis et al. 2018; Livingston et al. 2018; David et al. 2019; Bouma et al. 2020; Rizzuto et al. 2020; Plavchan et al. 2020; Newton et al. 2021; Nardiello et al. 2022; Barber et al. 2022; Zhou et al. 2022; Zakhozhay et al. 2022; Wood et al. 2023). The strategy pursued by most groups during the 2010s was to focus on stars with known ages—obvious members of open clusters—and to search them for transiting planets. The resulting stellar, and assumed planetary, ages are precise at the  $\approx 10\%$  level. A recent development facilitated by Gaia is the idea that finding transiting planets is often easier than finding the host star’s birth association (e.g. Tofflemire et al. 2021).

The “cluster-first” approach has a major limitation: most nearby stars are in the field. Only  $\lesssim 1\%$  of stars within  $\approx 500$  pc have been associated with their birth cluster (e.g. Zari et al. 2018; Cantat-Gaudin et al. 2020; Kounkel et al. 2020; Kerr et al. 2021). The implication can be appreciated by querying the NASA Exoplanet Archive (NEA; Akeson et al. 2013) for transiting planets younger than 1 Gyr. Requiring  $t < 1$  Gyr at  $2\sigma$  precision gives  $\approx 50$  such planets at the time of writing. Most of these young planets are in clusters, and were found by Kepler, K2, or TESS. However, these surveys have discovered  $\approx 5,000$  planets in total. Assuming a constant star formation history, we would expect an order of magnitude more sub-gigayear transiting planets.

This study aims to resolve two questions. First, how wrong is it to assume a uniform age distribution for transiting planet host stars? Second, where are the missing young planets? We will find that uniform is wrong by a factor of a few, and that stellar activity may be a less significant bias in young transiting planet detection than the challenge of precisely determining ages for field stars.

Despite Kepler’s main mission ending quite some time ago, the ages of many Kepler planets remain uncertain. While isochrone ages have been calculated for Kepler stars (Berger et al. 2020a) and Kepler Objects of Interest (KOIs; Petigura et al. 2022), such ages are most precise for stars whose luminosities and temperatures separate them from the main sequence. For sub-gigayear stars on the main sequence, isochrone ages are therefore limited.

Stellar rotation periods offer a promising alternative. The idea of using a star’s spin-down as a clock has a rich history (Skumanich 1972; Noyes et al. 1984; Kawaler 1989; Barnes 2003; Mamajek & Hillenbrand 2008; Angus et al. 2015). Empirical models now yield ages precise to  $\lesssim 30\%$  for FGK stars between 1–4 Gyr, and constraining age posteriors for younger ages (Bouma et al. 2023). Physics-based models (Matt et al. 2015; Gallet & Bouvier 2015; Spada & Lanzafame 2020) can connect these empirical relations to the evolution of stellar winds, internal structure, and angular momentum transport.

Rotation-based ages have been reported for various subsets of Kepler stars since the early data releases (e.g. Walkowicz & Basri 2013; McQuillan et al. 2014; Reinhold & Gizon 2015; Angus et al. 2018). More recent work has further explored incorporating information from stellar kinematics (Lu et al. 2021, 2024), and from stellar colors, luminosities, and

starspot amplitudes (Mathur et al. 2023). Our analysis is motivated by a few factors that can yield improvements, particularly for sub-gigayear stars. These factors are as follows.


- i) The Kepler Object of Interest catalog, and our vetting of false positives within it, has now reached maturity (e.g. Thompson et al. 2018).
- ii) Measured rotation periods of FGK stars in open clusters now show not only the average evolution of  $P_{\text{rot}}(T_{\text{eff}}, t)$ , but also how the astrophysical dispersion of stars around this average converges by the  $\approx 700$  Myr age of Praesepe (e.g. Curtis et al. 2019; Gillen et al. 2020; Rampalli et al. 2021; Fritzewski et al. 2021; Rebull et al. 2022; Dungee et al. 2022; Boyle & Bouma 2023).
- iii) Using open cluster data, we can marginalize over the range of ages that might explain any one star’s rotation period (Bouma et al. 2023). This represents an improvement in the accuracy of uncertainty propagation relative to previous calibrations.
- iv) We can identify binary stars with greater fidelity than in the past, which can clarify otherwise problematic estimates of rotation-based ages.
- v) New open clusters in various stages of dissolution have been found in the Kepler field (e.g. Kounkel & Covey 2019; Kerr et al. 2021; Barber et al. 2022). These clusters offer stars that we can use to test the reliability of the available single-star age-dating methods.

Independently, recent years have also yielded improvements in the lithium age scale. Lithium ages include depletion boundary ages for M dwarfs and brown dwarfs in star clusters, and decline-based ages for individual field FGK stars (Soderblom 2010). The latter approach relies on the observed decline of Li abundances in partially-convective stars as they age (e.g. Sestito & Randich 2005). The theoretical explanation for this decline is debated (e.g. Chaboyer et al. 1995; Denissenkov et al. 2010; Carlos et al. 2019). Empirical understanding however has improved due to the work by Jeffries et al. (2023), who modeled the time evolution of the Li I 6708 Å equivalent width (EW) using a set of 6,200 stars in 52 open clusters. Two-sided lithium ages are useful for Kepler (FGK) stars between  $\approx 0.03$ –0.5 Gyr, though with a strong dependence on spectral type. The precision of lithium ages in this regime are now in the range of 0.3–0.5 dex.

We discuss our method for selecting the star and planet samples in Section 2, and describe the origin of our adopted stellar parameters other than ages in Section 3. We describe our age-dating methods in Sections 4 and 5, and test them in Section 6 using clusters in the Kepler field. We discuss population-level trends in Sections 7 and 8, and offer a few conclusions in Section 9.

## 2. SELECTING THE STARS AND PLANETS

This work is focused on Kepler stars for which ages can be inferred using either rotation, lithium, or both. Such stars are a minority of the  $\approx 160,000$  Kepler targets. Rotation periods have been reported for roughly one in three Kepler targets (e.g. McQuillan et al. 2014; Santos et al. 2021).



**Figure 1. The stars.** Our analysis focuses on stars observed by Kepler with previously reported rotation periods (gray points). The rotation periods are primarily drawn from Santos et al. (2019, 2021). About half of the stars with rotation periods are suitable for gyrochronology (dark gray points), based on factors including their non-binarity and proximity to the main sequence (orange line in lower panel; see Section 4.1). Some host “confirmed” or “candidate” KOIs that meet additional planetary quality criteria (yellow points; Section 4.2). Surface gravities and effective temperatures were derived photometrically by Berger et al. (2020b). Isochrones in the lower panel are from MIST (Choi et al. 2016).

High-resolution spectra suitable for measurement of the Li I 6708 Å doublet have only been acquired for the Kepler objects of interest (KOIs), which comprise a few percent of Kepler’s targets. In the following, we will describe the set


of stars that we adopt with measured rotation periods (Section 2.1), the set of objects we adopt as hosting planets (Section 2.2), and the subset of these with high-resolution spectra suitable for lithium analysis (Section 2.3). Figure 1 provides a visualization of the various samples.

### 2.1. Stellar Rotation Periods

To select stars with rotation periods, we turn to previous work. Many investigators have derived rotation periods both specifically for KOIs (McQuillan et al. 2013; Walkowicz & Basri 2013; Mazeh et al. 2015; Angus et al. 2018; David et al. 2021), and also for the broader set of all Kepler target stars (McQuillan et al. 2014; Reinhold & Gizon 2015; Santos et al. 2019, 2021; Reinhold et al. 2023). These studies used a range of detection methods and selection functions. We are interested in understanding the age distribution of all Kepler target stars irrespective of KOI status. Out of these studies, the most homogeneous and precise analyses of both Kepler targets and KOIs appear to be those by Santos et al. (2019, 2021), hereafter S19 and S21. For a discussion of the work by Reinhold et al. (2023), please see Appendix A. S19 and S21 combined a wavelet analysis and autocorrelation-based approach, and cumulatively reported rotation periods for 55,232 main-sequence and subgiant FGKM stars. They included known KOIs and binaries in their analysis, and removed transits and eclipses during the stellar rotation measurement process.

We therefore adopt the results of S19 and S21 as our default rotation periods. S21 provided a comparison against McQuillan et al. (2014) (hereafter M14); the brief summary is that the periods agree for 99.0% of the 31,038 period detections in common between the two studies. S21 classified the 2,992 remaining stars from M14 as not showing rotation periods based on updated knowledge of contaminants (e.g. giant stars and eclipsing binaries) and visual inspection. In addition, S21 reported rotation periods for 24,182 main-sequence and subgiant FGKM stars that were not reported as periodic by M14. Many of these reported detections have lower variability amplitudes and longer periods than those reported by M14.

Analyzing the compilation of S19 and S21 rotation periods for the KOI hosts, we noticed that some known KOIs with rotation periods were missing. This is not surprising, since the rotation periods of KOIs have received more scrutiny than those of ordinary Kepler stars. We therefore decided to split our subsequent analysis into a homogeneous portion that used only the S19 and S21 data, and an inhomogeneous portion that also considered a broader set of available KOI rotation periods. For the latter portion, we first included 32 KOIs with orbital and rotation periods within  $\approx 20\%$  that had been excluded from the S19 and S21 catalogs (A. Santos, private communication). We then incorporated an additional 178 rotation periods for KOI hosts that David et al. (2021) described as either “reliable” or “highly reliable” in their visual analysis of previously reported KOI rotation periods from McQuillan et al. (2013), Walkowicz & Basri (2013), Mazeh et al. (2015) and Angus et al. (2018). Inclusion of these additional KOI rotation periods is a supplementary measure aimed at com-



**Figure 2.** Rotation periods for Kepler target stars (left) and known planet hosts (right). The gray lines are “mean fits” to the rotation sequences of open clusters. The 2-D histogram of the stellar sample (left) includes only apparently single stars near the main sequence with  $\log g > 4.2$ ,  $\text{RUWE} < 1.4$ , and temperatures of 3800–6200 K. The planet-hosts (right) require the same stellar cuts, and include only the confirmed and candidate planets described in Section 4.2. The number of stars  $N_s$  and planets  $N_p$  are noted in the relevant panels. A simple estimate for the number of young planets discovered by Kepler follows by counting points below the mean rotational isochrones.

pleteness in our final KOI age catalog; the provenances of the individual adopted periods are noted in the relevant tables.

Finally, since our scope is focused on rotation-based ages, we restricted our attention to stars with reported  $P_{\text{rot}} < 45$  days. The slowest-rotating FGK stars in the open clusters used to calibrate our gyrochronology model have  $P_{\text{rot}} \approx 35$  days. Figure 2 shows a subset of the resulting 53,663 Kepler stars with rotation periods compiled from S19, S21, and our extended KOI list.

To assess the statistical uncertainties of these rotation periods, we compared our adopted periods with those reported by McQuillan et al. (2014). The details are in Appendix A. We found that for  $P_{\text{rot}} \lesssim 15$  days, the two datasets agree at a precision of  $\lesssim 0.01 P_{\text{rot}}$ . At longer periods of  $P_{\text{rot}} \approx 30$  days, the agreement was typically at the  $\lesssim 0.03 P_{\text{rot}}$  level, and the envelope of the period difference increased roughly linearly with period. Based on this comparison, we adopted a simple prescription for the period uncertainties, such that there are 1% relative uncertainties below  $P_{\text{rot}} = 15$  days, and a linear increase thereafter, with slope set to require 3%  $P_{\text{rot}}$  uncertainties at rotation periods of 30 days.

## 2.2. Kepler Objects of Interest

We considered planets in the NEA cumulative KOI table as of 2023 June 6, which included the best knowledge available on any given planet candidate while also incorporating human-based vetting. These planets represent a superset of those in the fully automated Q1-Q17 DR25 KOI Table (Thompson et al. 2018), which could be adopted in fu-

ture work for planet occurrence rate calculations. This version of the cumulative KOI table included 4,716 objects that are either “confirmed” or “candidate” planets, after excluding known false positives.

## 2.3. High Resolution Spectra

The final piece of our analysis involves assessing ages based on the Li I 6708 Å doublet. We analyzed spectra from the High Resolution Echelle Spectrometer (HIRES; Vogt et al. 1994a) on the Keck I 10m telescope. These spectra were primarily collected through the California Kepler Survey (Pettigura et al. 2017; Johnson et al. 2017; Fulton et al. 2017). We supplemented the existing archive with  $\approx 10$  hours of new observations for 22 stars between Fall 2022 and Spring 2024. These stars were chosen to ensure that confirmed planets with rotational evidence for ages below 1 Gyr had spectra, since this is the age range in which lithium is most likely to yield useful age constraints.

Lithium equivalent widths and abundances for the Kepler Objects of Interest were already analyzed by Berger et al. (2018) for roughly three quarters of the spectra in our sample. However, new spectra have since been acquired, and our approach and selection function are different. We therefore performed our own line width measurements on the reduced HIRES spectra.

We collected all blaze-corrected HIRES spectra from our group’s observations of non-false positive Kepler Objects of Interest with a “multiple event statistic” (MES, *koi\_max\_mult\_ev*) of at least 10. This yielded at least one




spectrum for 1,464 stars hosting 2,174 planets. About half of these stars have measured rotation periods (797 stars and 1,170 planets, respectively). For stars with multiple spectra available, we analyzed only the spectrum with the highest number of counts. The resulting spectra were acquired between 2009 September 6 and 2024 May 16, and cumulatively comprise 304 hours of open-shutter time.

We measured the lithium equivalent widths using a procedure adapted from previous work (Bouma et al. 2021). Our stars of interest are FGK stars, and so the continuum in the vicinity of the Li I 6708 Å doublet is well-defined. We Doppler-corrected the spectra to a common reference wavelength by cross-correlating against a high S/N template for Kepler-1698, chosen because  $|\gamma| < 10 \text{ km s}^{-1}$ ,  $T_{\text{eff}} \approx 5000 \text{ K}$  and  $v \sin i \approx 5 \text{ km s}^{-1}$ , which puts it in the middle of our sample’s temperature range and gives it mild line-broadening. We then trimmed the Doppler-corrected spectra to a local window centered on the lithium line, using a window width of 15 Å (we also considered 10 Å and 20 Å; the results were consistent). We continuum-normalized by fitting a third-degree Chebyshev series, while excluding regions with absorption lines. We then numerically integrated the resulting spectrum using a one-component Gaussian with free amplitude, width, and mean, and estimated uncertainties on the line width through a Monte Carlo procedure that bootstrapped against the local scatter in the continuum. The resulting EWs are shown in Figure 3.

Our EW measurement approach did not correct for the neighboring Fe I 6707.44 Å blend. To evaluate the accuracy and precision of our method, after applying an initial iteration on the 1,464 stars with spectra, we compared our lithium equivalent widths with those reported by Berger et al. (2018). For the stars in both samples, we found broad agreement at  $> 30 \text{ mÅ}$ , and significant differences at  $< 30 \text{ mÅ}$  because Berger et al. (2018) required positive EWs, while we allowed for statistically negative ones. At  $> 30 \text{ mÅ}$ , there is however a small offset in our respective scales caused by our non-treatment of the iron blend, such that (B24-B18) =  $7.50 \pm 8.55 \text{ mÅ}$ . We therefore directly subtracted this constant mean value ( $7.50 \text{ mÅ}$ ) when calculating lithium-based ages. This offset is at worst five times smaller than the astrophysical scatter present in lithium EWs for calibration clusters at any given age (see Jeffries et al. 2023).

### 3. STELLAR PROPERTIES

Our default source for stellar temperatures and surface gravities was the Gaia-Kepler Stellar Properties Catalog (GKSP; Berger et al. 2020b). The GKSP parameters were reported for stars with “AAA” 2MASS photometry, measured parallaxes in Gaia DR2, and g-band photometry available from either SDSS or else Kepler-INT (Greiss et al. 2012). The parameters themselves were derived using *isoclassify* (Huber et al. 2017) to interpolate over the MIST isochrone grids (Choi et al. 2016; Dotter 2016), given the SDSS g and 2MASS  $K_s$  photometry, the Gaia DR2 parallaxes, and V-band extinction from the Green et al. (2019) reddening map. The resulting stellar parameters are available




**Figure 3. Equivalent widths (EW) of the Li I 6708 Å doublet for planet-hosting stars.** These measurements were made from Keck/HIRES spectra collected from 2009–2024. Lines are the “mean” isochrones from Jeffries et al. (2023). The intrinsic dispersion around these isochrones becomes much larger than changes in the mean at  $\geq 1 \text{ Gyr}$  (see Figure 4). Some stars with lithium detections do not have detected rotation periods, and vice-versa.

for  $\approx 94\%$  of the 53,663 Kepler stars with rotation periods. For the remaining  $\approx 6\%$  of stars that lack temperatures and surface gravities from B20, we adopted the values reported by Santos et al. (2019) and Santos et al. (2021), which are primarily derived from the photometric Mathur et al. (2017) DR25 Kepler Stellar Properties Catalog. In the planet sample,  $\approx 92\%$  of the non-false-positive KOIs with rotation periods have parameters from Berger et al. (2020b), and the remainder are drawn from DR25.

David et al. (2021) compared the photometric B20 stellar parameters ( $T_{\text{eff}}$ ,  $R_*$ , [Fe/H]) against the spectroscopic parameters from Fulton & Petigura (2018). The temperature scales showed a few-percent systematic difference, with F18 quoting higher temperatures than B20 for mid K dwarfs, and lower temperatures for early F dwarfs. Our age analysis, described below, adopts the maximum of the two-sided uncertainties reported by B20 as a symmetric Gaussian temperature uncertainty. Systematic uncertainties in the temperature scale generally influence our age uncertainties at a smaller level than the statistical intrinsic scatter in the open cluster rotation sequences.

### 4. AGES FROM ROTATION

We calculated gyrochrone ages using *gyro-interp* (Bouma et al. 2023), which is a method designed to address the fact that stars with the same mass and same rotation period can have a wide range of ages (e.g. Curtis et al. 2019).



**Figure 4. The models.** Points represent  $10^4$  draws from models that have been fitted to rotation periods (Bouma et al. 2023) and lithium equivalent widths (EWs; Jeffries et al. 2023) of stars in open clusters. Lines are the “mean models” at various ages. The intrinsic dispersion around the mean, which is what the models fit, sets the theoretical precision floor for the age-dating methods. Additional sources of uncertainty, including measurement uncertainty, impose further limits on achievable precision. These models are calibrated using data from open clusters. The sizes of the points are the same in each panel, so that low apparent density signifies greater dispersion around the mean.

The gyrochrone age posterior should therefore incorporate the intrinsic population-level scatter into its statistical precision. Figure 4 highlights this problem, particularly in regions where the 0.1–1 Gyr stars overlap.

To estimate the stellar spin-down rate as a function of time and stellar temperature, *gyro-interp* uses measured rotation periods and effective temperatures from reference open clusters, and interpolates between them using cubic Hermite polynomials. We calculated the probability of the rotation-based age,  $t_{\text{gyro}}$ , given the observed periods and temperatures (and their uncertainties) by integrating Equation 1 of Bouma et al. (2023). This procedure marginalizes over the astrophysical scatter that is observed in the open cluster sequences. We adopted a linear age grid spanning 0–5 Gyr with 500 grid points, and interpolated the resulting posteriors to calculate any summary statistics. The oldest cluster for which *gyro-interp* is calibrated is the 4 Gyr M67 cluster (Dunee et al. 2022; Gruner et al. 2023). For Kepler stars with rotation periods above the M67 sequence, our resulting age constraints are quoted as lower limits.

#### 4.1. Stellar Quality Flags

We calculated gyrochrone ages for all 45,229 stars with reported rotation periods that had effective temperatures of 3800–6200 K. To identify stars for which we suspect these ages may not be valid, we then built a set of quality flags which we condensed into a single binary number:  $Q_{\text{star}}$ . When and how this bitmask should be applied depends on the question being asked. If the goal is to construct a false-

positive free sample, all the quality flags could be applied. If the goal is to construct a complete sample, then consider the examples of Kepler 1627Ab ( $\approx 40$  Myr) and Kepler 51c ( $\approx 625$  Myr). The former has a high RUWE due to a resolved binary companion (Bouma et al. 2022a); the latter is on a grazing orbit (Masuda 2014). We leave selection for or against such cases as a decision to the user. For our own analysis, we assume that a star is suitable for gyrochronology if none of bits zero through nine (inclusive) are raised. For analyses that require all stellar rotation periods to come from the same detection pipeline, we further require bit 11 to be zero.

Three assumptions must hold for a rotation-based age dating method like *gyro-interp* to be valid. 1) The evolutionary state of the star must be well-specified by its temperature and its age, 2) the star’s spin-down must not be influenced by binary companions, and 3) the rotation period distribution for field stars of a given temperature and age must be identical to that of equally aged open clusters (metallicity differences, for instance, are ignored). A rephrasing of the first condition is that the star must be “near the main sequence” because during the post main sequence stellar temperatures change. From  $\approx 0.08$  to  $\approx 3$  Gyr, the stars of interest in this work (3800–6200 K,  $0.5$ – $1.2 M_{\odot}$ ) have temperatures constant to  $\approx 1\%$  (Choi et al. 2016). From  $\approx 3$ –4 Gyr, a  $1.2 M_{\odot}$  star’s temperature drops from  $\approx 6200$  K to  $\approx 6000$  K ( $\approx 3\%$ ), because its outer layers expand as it begins hydrogen shell burning as a subgiant. We treat this issue in a manner discussed in “bit 9” below.

*Temperature range (bit 0)*—We require stars to have effective temperatures  $T_{\text{eff}}$  between 3800 and 6200 K in order to report gyrochrone ages (Bouma et al. 2023). Stars hotter than 6200 K spin down very slowly, if at all. Stars cooler than 3800 K do spin down over gigayear timescales (Newton et al. 2016; Engle & Guinan 2023; Chiti et al. 2024), but the currently available open cluster data have yet to clarify when the intrinsic scatter in the population decreases.

*Surface gravity (bit 1)*—We flagged stars as possible subgiants if they had  $\log g < 4.2$ .

*Absolute luminosity (bit 2)*—We calculated the absolute Gaia DR3  $G$ -band luminosity, ignoring reddening, using the reported apparent  $G$ -band magnitude and parallaxes. We flagged stars with  $M_G < 3.9$  or  $M_G > 8.5$ , corresponding to main-sequence spectral types earlier than  $\approx F8V$  or later than  $\approx M0.5V$  (Pecaut & Mamajek 2013).

*Known eclipsing binaries (bit 3)*—We flagged any stars reported to be in the final Kepler eclipsing binary catalog (KEBC; Kirk et al. 2016).

*Kepler-Gaia crossmatch quality (bits 4 and 5)*—To leverage Gaia DR3 data, we used M. Bedell’s 4'' Kepler-to-Gaia crossmatch of the NEA `q1_q17_dr25_stellar` catalog with Gaia DR3 (available at <https://gaia-kepler.fun>). The separation distribution of the Kepler-Gaia DR3 crossmatches is such that 99.2% of candidate matches are within 1''. We nonetheless noted an upturn in the candidate match rate from 3-4''; such sources are flagged using bit 4. For KIC stars with multiple potential Gaia matches within the 4'' radius, we adopted the brightest star as the default match. In most such cases this was unambiguous because there is a large brightness difference between the primary and any apparent neighbors. However cases with multiple stars within 4'' within  $\Delta G < 0.5$  mag are noted using bit flag 5.

*Gaia DR3 non-single-stars (bit 6)*—The Gaia DR3 `non_single_star` column in the `gaia_source` table flags known eclipsing, astrometric, and spectroscopic binaries. We directly included this column.

*RUWE (bit 7)*—We inspected diagrams of the Gaia DR3 renormalized unit weight error (RUWE) as a function of other stellar parameters, and flagged stars with  $\text{RUWE} > 1.4$  as possible binaries. Such astrometric outliers can be either bona fide astrometric binaries, or more often are marginally resolved point sources for which a single-source PSF model provides a poor fit.

*Crowding (bit 8)*—We searched the Gaia DR3 point source catalog for stars within 1 Kepler pixel (4'') of every target star. While such companions may not be physically associated with the target star, their presence can confuse rotation period measurements. We therefore flagged any stars with neighbors down to  $1/10^{\text{th}}$  the brightness of the target star within this region ( $\Delta G < 2.5$ ).

*Near the main sequence (bit 9)*—Figure 1 shows that many stars observed by Kepler are far from the main sequence. Some of the challenges this introduces for rotation-based ages include unresolved binaries, metallicity, reddening, and drivers of rotation other than magnetic braking. After exploring various options, we settled on the orange locus in the

$\log g$ - $T_{\text{eff}}$  plane shown in Figure 1 as a way of flagging unresolved binaries, as well as evolved late-F and early-G stars. While the exact details of how this locus is constructed are arbitrary (see Appendix B), the general aim is to flag stars for which there is evidence based on their location in the HR diagram that our gyrochronology model may not be reliable. While the  $\log g$ - $T_{\text{eff}}$  cut flags about half of Kepler stars with rotation periods as potentially not being suitable for gyrochronology, a few outliers in  $M_G$  vs.  $G_{\text{BP}}-G_{\text{RP}}$  remained after applying it. Such outliers are likely also questionable. We therefore also fitted a polynomial to the KOI main sequence in  $M_G$  vs.  $G_{\text{BP}}-G_{\text{RP}}$ , and flagged stars more than 1 magnitude from this locus as part of the same bitflag.

*Candidate pulsators and close-in binaries (bit 10)*—Santos et al. included in a flag for candidate “classical pulsators” (e.g. Cepheids) and close-in binaries. Visual inspection shows that although this flag selects many bona fide objects in these classes, it also selects most known young planets with  $P_{\text{rot}} < 5$  day. These objects are neither classical pulsators nor close binaries. While we propagated this flag into our set of quality flags, we therefore generally ignore it.

*Not in the homogeneous stellar rotation sample (bit 11)*—This flag was set for the KOIs with rotation periods drawn from a source other than the Santos et al. pipeline.

#### 4.2. Planet Quality Flags

Some planets are more reliably identified than others. We used the following additional quality indicators to assess the reliability and utility of a planet, and assembled them into a separate bitmask,  $Q_{\text{planet}}$ .

*Candidate reliability (bit 0)*—The NEA’s Kepler Objects of Interest Table includes both an overall planet-candidate disposition status (`koi_disposition`), as well as a disposition based only on the Kepler data (`koi_pdisposition`). We required both to include only “planet candidates” and “confirmed planets.”

*Candidate S/N (bit 1)*—In any transit survey, the false positive rate increases greatly toward the noise floor for planet detection (e.g. Jenkins et al. 2002). We required a S/N in excess of Kepler’s usual  $7.1\sigma$  floor, through a cut on the maximum “multiple event statistic” (MES, `koi_max_mult_ev`): we required  $\text{MES} > 10$ .

*Grazing planets (bit 2)*—Grazing objects, for which the impact parameter  $b$  is greater than  $1 - R_p/R_*$ , often yield biased planetary parameters (e.g. Gilbert 2022). For large planets, they also include astrophysical false positives at higher rates (Morton et al. 2016), in part due to the size-impact parameter degeneracy. We flagged planet candidates as potentially grazing if  $b > 0.8$ , using the impact parameters reported by Thompson et al. (2018).

#### 5. AGES FROM LITHIUM

Figure 3 shows our measured equivalent widths for the lithium 6708 Å absorption doublet, plotted over the mean isochrone models from EAGLES (Jeffries et al. 2023). We show an upper-limit for plotting purposes if the  $1\sigma$  lower limit on the equivalent width is below 10 mÅ. Our measured



EWs span -40 to 250 mÅ; all negative EWs have uncertainties that are statistically consistent with zero.

To calculate lithium ages, we used EAGLES (git commit ac09637), which, similar to gyro-interp, is based on an empirical interpolation approach. EAGLES was calibrated on lithium measurements of stars observed by the Gaia-ESO spectroscopic survey in 52 open clusters with ages spanning 2 to 6,000 Myr (Jeffries et al. 2023). We adopted a linear prior in age, spanning  $10^6$  to  $10^{10}$  yr, and symmetrize our EW uncertainties for purposes of interfacing with EAGLES by taking the maximum of the measured positive and negative  $1\sigma$  uncertainty intervals.

In part because many of our EWs are upper limits, many of the lithium ages are lower limits. In such cases, the inferred age posterior is strongly influenced by the assumed intrinsic dispersion in the lithium model, and by the prior. In these instances, we quote the  $3\sigma$  (99.7<sup>th</sup> percentile) limits.

## 6. CLUSTER AGE COMPARISON

Stars that formed in the same birth cluster are the gold standard for the astronomical age scale (Soderblom 2010). Before Gaia, the few open clusters known in the Kepler field had been cataloged by Herschel (1864). Gaia has enabled the discovery of stellar ensembles that are more diffuse, but which nonetheless share a common age based on isochrone, rotation, and lithium dating (e.g. Kounkel & Covey 2019; Bouma et al. 2022b; Barber et al. 2022).

In Figure 5, we compare our rotation-based ages, and available ages from the literature, against the ages of stars in open clusters in the Kepler field. From the literature, we drew ages from Reinhold & Gizon (2015), Berger et al. (2020b), Lu et al. (2021), Mathur et al. (2023), and Lu et al. (2024). We followed any guidance available from each study for removing ages that were unreliable, and plotted stars within  $1\sigma$  of zero as upper limits. For the Reinhold & Gizon ages, we used those calculated using the Mamajek & Hillenbrand (2008) calibration; for the Mathur et al. ages, we show those from kiauhoku (Claytor et al. 2020) rather than STAREVOL (Amard et al. 2019), since the former showed better agreement with the cluster age scale.

Cluster membership is a nuanced subject. Figure 5 is showing reported ages for a set of spatially and kinematically selected stars that could be cluster members, or they could be field interlopers. For NGC6811 and NGC6819, we adopted candidate members from Cantat-Gaudin et al. (2018, 2020) and Kounkel et al. (2020). For Theia 520, we used candidate members from Kounkel et al. (2020). For Melange-3, we used the candidates reported by Barber et al. (2022) and required “offset” tangential velocities below  $2 \text{ km s}^{-1}$ . For Cep-Her, we used candidate members from Kerr et al. 2024 (submitted) with  $P_{\text{fin}} > 0.8$ . Even with contaminants, we can compare the relative age distributions derived by the different studies because in all instances we are comparing reported ages against a fixed list of stars.

There are two main metrics for success in this test. 1) Do the reported ages agree with the cluster age? 2) Do the re-

ported age uncertainties agree with their dispersion around the cluster age?

Figure 5 show that although previous studies reported ages that agree with the cluster scale for  $\gtrsim 1$  Gyr stars, sub-gigayear stars have historically had their ages overestimated by 0.3–2 dex, with severely understated uncertainties. For isochrone and kinematic ages, this is because these methods rely on parameters that do not appreciably change at  $t \lesssim 1$  Gyr. For the rotation-based ages in Mathur et al. (2023) and Lu et al. (2024), the discrepancy is caused by details of the calibration methodologies. The kiauhoku spin-down model for instance has known “stretches” and “compressions” in its age scale relative to observed open clusters (see Mathur et al. 2023, Sec. 7.3). The Mamajek & Hillenbrand (2008) calibration used by Reinhold & Gizon (2015) appears more accurate than the former two models, because it was fitted to reproduce the open cluster sequences known at the time. However, the uncertainties from all of these methods appear to be underestimated. This is probably because they do not marginalize over the range of rotation periods accessible to sub-gigayear stars of a fixed mass and age.

## 7. RESULTS


### 7.1. A Paucity Of Young Stars And Planets

The stellar ages are given in Table 1 for the known planet hosts, and in Table 2 for all Kepler stars. Figure 6 shows the rotation-based age distributions of all the stars (left column) and the KOI hosts (middle column). These “histograms” are sampled from the posterior probability distributions by drawing ten random samples from each posterior, and then computing the normalized histogram of the resulting samples. We considered an alternative approach for constructing these plots using hierarchical Bayesian deconvolution, and found similar results. The uncertainties assume Poisson statistics. We truncated the plots near 4 Gyr since this is the cluster age out to which our rotational ages are calibrated. The overall slope yields a paucity of young stars relative to the expectation of a uniform age distribution.

The age distributions of Kepler stars and KOI hosts in Figure 6 appear similar. One diagnostic for whether the two distributions are drawn from the same underlying distribution is the Kolmogorov-Smirnov test; we calculated this statistic in a manner that accounts for the Poisson uncertainties by performing 1000 random draws of 70% of the stars from each sample when requiring  $t_{\text{gyro}} < 3$  Gyr. The resulting  $\log_{10} p$  values spanned (2.5<sup>th</sup> to 97.5<sup>th</sup> percentiles) -4.5 to -1.8 for the K dwarfs, and -4.5 to -1.9 for all Kepler stars. This agrees with the visual impression that while small differences may be present, they are not drastic.

The similarity of the KOI host and Kepler star age distributions adds some nuance to the argument that young transiting planets are hard to detect due to the photometric variability of their host stars. If true, this statement seems to hold only for the very youngest ( $\lesssim 0.4$  Gyr) stars, where there is a marginal deficit in the KOI host age distribution relative to the parent stellar sample. Examining scatter plots of the planet properties as a function of time, we similarly find that fewer sys-





**Figure 5. Reported ages of individual stars in benchmark open clusters.** Each point denotes a star’s reported age, normalized by the age of a stellar ensemble in which the star is a candidate member. Cluster membership was evaluated without knowledge of rotation. Although some field interlopers may be present in the membership lists, outliers can be compared between different methods on a relative basis. Each study was cross-matched against the same cluster list; certain methods report ages for more stars than others. Horizontal scatter is added to visually clarify the statistical age uncertainties. At  $<1$  Gyr, our ages are generally accurate, and their statistical uncertainties match their dispersion.

tems at  $t_{\text{gyro}} < 0.4$  Gyr are detected with  $P_{\text{orb}} \gtrsim 30$  days than at  $t_{\text{gyro}} > 0.4$  Gyr.

An important separate selection effect concerns Kepler’s ability to detect the rotation signals. Masuda (2022) studied this issue, and found for Sun-like stars that the fraction of rotation signals detectable by Kepler is near unity up to  $\approx 3$  Gyr, and that it drops to almost zero by  $\approx 5$  Gyr. This trend is opposite in functional form to our derived age distributions.

We can quantify the relative counts of stars as a function of age by labelling stars between 0-1 Gyr, 1-2 Gyr, and 2-3 Gyr as “young”, “intermediate-age”, and “old”. A simple counting exercise from Figure 6 tells us that there are 2.1 times as many old stars in the Kepler field as young stars. Similarly, there are 2.7 times as many old planet hosts as young planet hosts. Focusing only on the tails of the distributions (0-0.3 Gyr and 2.7-3 Gyr), the implication is that the formation rate of stars in the Kepler field decreased by a factor of  $2.85 \pm 0.12$  over the past three billion years. If this decrease continues linearly into the future, then a simple extrapolation would imply that in  $\approx 1.4 \pm 0.3$  Gyr the rate of new stars forming in the Galaxy will reach zero.


In terms of detected planet counts, our rotation-based ages for the Kepler sample yield {108, 211, 288} detected planets in the 0-1 Gyr, 1-2 Gyr, and 2-3 Gyr bins.

## 7.2. Planets Younger Than One Billion Years

Sub-gigayear planets can be particularly informative for studies of planet evolution. Our catalog has 108 confirmed and candidate planets with median ages below 1 Gyr. Requiring  $t_{\text{gyro}} < 1$  Gyr at  $2\sigma$  yields 60 planets orbiting 47 stars. The youth claim is most secure for the systems that are either in clusters (Section 7.2.1), or that have independent rotation and lithium-based ages (Section 7.2.2). These samples provide broader context for the planets around field stars that push the boundaries of the current young exoplanet census (Section 7.2.3).

### 7.2.1. Planets in Clusters

Four stellar ensembles in the Kepler field have to date yielded a total of fourteen transiting planets. Our analysis blindly recovers the youth of all of these planets.




**Figure 6. Kepler’s demographic cliff**, visible in the rotation-derived age distributions of its stars (left) and planet hosts (middle). The top row shows all stars with temperatures of 3800–6200 K for which we calculated rotational ages. Opaque lines impose quality cuts on binarity, crowding, and the star’s evolutionary state; transparent lines do not. The bottom row shows stars with temperatures of 4400–5400 K, which have more precise ages due to their fast spin-down. The statistical uncertainty for a mean star at 1, 2, and 3 Gyr is shown, and is identical across each row. Finally, the right panel compares the rotation-derived ages against star formation histories derived using CMD fitting (dashed and dotted lines). Completeness in Kepler’s  $P_{\text{rot}}$  detection sensitivity is near unity until  $t \lesssim 3$  Gyr (Masuda 2022).

*Cep-Her*—The Cep-Her complex ( $\approx 40$  Myr) contains Kepler-1627Ab, Kepler-1643b, Kepler-1974b and Kepler-1975 Ab (Bouma et al. 2022a,b). Different sub-groups of the complex vary in age by  $\approx 50\%$  (Kerr et al., submitted). The planets themselves are all on close-in orbits (5–25 days), with sizes from  $2\text{--}4 R_{\oplus}$ . Our rotation and lithium ages agree with the cluster age for Kepler-1627A, Kepler-1974, and Kepler-1975A. For Kepler-1643, the rotation and cluster ages agree, and the lithium age is  $2.0\sigma$  above the cluster age reported by Bouma et al. (2022b) for RSG-5.

*MELANGE-3*—Barber et al. (2022) reported a  $105 \pm 10$  Myr association in the Kepler field containing two transiting mini-Neptunes: Kepler-970 and Kepler-1928. We find rotation-based ages of  $t_{\text{gyro}} = 176^{+120}_{-40}$  Myr and  $144^{+104}_{-88}$  Myr for Kepler-970 and Kepler-1928 respectively. The former

rotation-based age being slightly older than the suggested Pleiades age for the association agrees with Figure 4 of Barber et al. (2022).

*Theia 520*—We derive rotation-based ages of  $\approx 350$  Myr for Kepler-968 and Kepler-52. Based on spatial and kinematic clustering, these stars were reported by Kounkel & Covey (2019) to be members of Theia 520. The core of this structure is also known as UBC-1 (Castro-Ginard et al. 2018). Based on isochrone and rotation-based age-dating, Theia 520 seems to be  $\approx 230 \pm 70$  Myr old (Kounkel & Covey 2019; Fritzewski et al. 2024), which is within  $1\sigma$  of our  $t_{\text{gyro}}$  measurements. Ongoing work by Curtis (2024) broadly finds that the two planet-hosting stars are indeed members of this diffuse population. Given the cluster-level and individual-star evidence, this makes Kepler-968 and Kepler-52 the



**Figure 7.** Sizes, orbital periods, and rotation-based ages of transiting exoplanets younger than one billion years. Star symbols denote 43 planets that are characterized in this work to have  $t_{\text{gyro}} < 1$  Gyr at  $2\sigma$ ; circles are planets from the literature meeting the same age requirement, but with a heterogeneous set of age provenances. Gray points are transiting planets from the NASA Exoplanet Archive older than 1 Gyr. If one were to instead select for precise ages ( $t/\sigma_t > 3$ ), this would hide  $\approx$ ten  $< 0.5$  Gyr planets, and add  $\approx$ fifteen  $0.5$ – $1$  Gyr planets. For this plot, we require our Kepler stellar hosts to not have  $Q_{\text{star}}$  bits 0–9 raised, and for the planets to similarly not have quality flags raised.

youngest multiplanet systems currently known from the main Kepler mission.

*NGC 6811*—Meibom et al. (2013) reported the discovery of Kepler-66 and -67b, two mini-Neptunes in NGC 6811 ( $\approx 1$  Gyr). We derive rotation-based ages for these systems of  $1417^{+635}_{-353}$  Myr and  $877^{+107}_{-120}$  Myr for Kepler-66 and -67b, respectively. Kepler-66 ( $T_{\text{eff}} \approx 5900$  K,  $P_{\text{rot}} \approx 10.4$  days) has a lower precision and a more asymmetric posterior because of its slow spin-down rate. Kepler-67 ( $T_{\text{eff}} \approx 5100$  K,  $P_{\text{rot}} \approx 10.3$  days) is marginally hotter than the later K dwarfs that are in “stalled” spin-down at this time, enabling its rotation period to be diagnostic of its age.

### 7.2.2. Rotation vs. Lithium Ages

We compared the rotation and lithium ages for the known planet hosts in the “Consistent?” column of Table 1. There were three cases of interest. *i*) If two-sided posteriors for both  $t_{\text{gyro}}$  and  $t_{\text{Li}}$  existed, and their median values were consistent within  $2\sigma$ , we listed them as consistent; if they were consistent within  $2$ – $3\sigma$ , we listed them as “maybe” consistent. Otherwise, they disagreed. *ii*) If  $t_{\text{Li}}$  was a lower limit, we compared this lower limit with the  $1\sigma$  upper limit from  $t_{\text{gyro}}$ ; if the two overlapped, we judged the age estimates to be consistent. *iii*) If  $t_{\text{Li}}$  provided a two-sided posterior, and no rotation period was found, we judged the age estimates to be inconsistent. This is because a two-sided lithium constraint can only be provided for G and K dwarfs  $\lesssim 2$  Gyr old,

and Kepler should have been sensitive to the rotation periods of such stars.

In the entire planet sample (i.e. without imposing any quality cuts) this yielded 697 consistent cases, 12 maybe consistent cases, and 22 inconsistent cases. Rephrased, in the overall sample, 95% of stars have consistent rotation and lithium-based ages (97% potentially consistent).

Appendix C describes all cases of “discrepant” confirmed planets for which a sub-gigayear age is reported by at least one age indicator. All but two systems are either evolved stars or else unresolved binaries, and are automatically flagged as such. The more interesting of the two is Kepler-786, an early K dwarf with  $t_{\text{Li}} = 228^{+168}_{-87}$  Myr, but with a  $\approx 33$  day rotation period that implies  $t_{\text{gyro}} = 4390^{+243}_{-241}$  Myr. Other age indicators in the spectrum similarly suggest that the star is not young.

### 7.2.3. Notable New Young Planets

Let us focus on blemish-free stars with blemish-free planets:  $Q_{\text{star}}$  must not have any of bits zero through nine raised, and  $Q_{\text{planet}}$  must similarly have no quality flags raised. Under this constraint, our results include 14 confirmed planets younger than 1 Gyr with two-sided  $t_{\text{gyro}}$  and  $t_{\text{Li}}$ , and 39 confirmed planets with comparable ages for which  $t_{\text{gyro}}$  is two-sided while  $t_{\text{Li}}$  is one-sided.

Figure 7 shows the planets and planet candidates with  $t_{\text{gyro}} < 1$  Gyr at  $2\sigma$ . This figure includes 34 confirmed Kepler planets, and nine candidate planets, all of which are listed in Table 1. These include six objects Earth-sized or smaller, two Jovian-sized planet candidates, four super-Neptunes ( $4$ – $10 R_{\oplus}$ ), 19 mini-Neptunes above the radius valley as parametrized by Van Eylen et al. 2018, and 12 super-Earth sized planets below the radius valley. While most of these planets are on orbital periods below 50 days, two are on more distant orbits.

Among these systems, four highlights are Kepler-1529, Kepler-1565, Kepler-1312, and Kepler-1629. Kepler-1529b is a  $\approx 100 \pm 50$  Myr mini-Neptune with well-measured rotation and lithium ages. Kepler-1565b is an analogous  $\approx 170$ – $230$  Myr super-Earth. Kepler-1312 is a  $t_{\text{gyro}} = 357^{+75}_{-109}$  Myr near-Solar analog with an Earth-sized planet on a one-day orbit, and a mini-Neptune on a five-day orbit. The Earth-sized planet, Kepler-1312 c, along with the near-identical Kepler-1561 b ( $t_{\text{gyro}} = 426^{+74}_{-78}$  Myr), rank among the youngest Earth-sized planets currently known, comparable to planets such as HD 63433d ( $1.1 R_{\oplus}$ ,  $414 \pm 23$  Myr; Capistrant et al. 2024) and TOI-1807 b ( $1.3 R_{\oplus}$ ,  $180 \pm 40$  Myr; Hedges et al. 2021). Kepler-1629 b, with a size just two-thirds that of Earth ( $0.67 \pm 0.05 R_{\oplus}$ ), is marginally older, also based on rotation ( $529^{+62}_{-62}$  Myr). The lithium age from a reconnaissance TRES spectrum confirms the point.

## 8. DISCUSSION

### 8.1. The Thin Disk’s Star Formation History

Most of the stars observed by Kepler are in the thin disk. This can be verified by following Gaia Collaboration et al. (2018), and labeling stars with 2D tangential velocities  $v_T < 40 \text{ km s}^{-1}$  as thin disk members, and those with



60  $< v_T < 150 \text{ km s}^{-1}$  as thick disk members. For all stars in the KIC, this yields 95,694 thin and 54,039 thick disk stars respectively. The thin disk stars have a threefold larger detected rotation period fraction: 35,675 rotators are in the thin disk, while 7,312 are in the thick disk. Further imposing the gyrochronology quality flags discussed in Section 4.1 yields 17,755 thin-disk stars for which gyrochronology is applicable, and 2,462 thick-disk stars.

Classifying Kepler stars as thin vs. thick disk members helps connect them to previous work on the Galaxy’s star formation history. On extragalactic scales, the star formation rate in spiral galaxies with mass similar to the Milky Way peaked  $\approx 10 \text{ Gyr}$  ago, and has since decreased by an order of magnitude (e.g. Heavens et al. 2004; Hopkins & Beacom 2006). In our own galaxy, previous studies have measured star formation histories through CMD fitting of resolved stellar populations (Mor et al. 2019; Ruiz-Lara et al. 2020; Alzate et al. 2021; Xiang & Rix 2022) and by modeling the local white dwarf luminosity function (e.g. Isern 2019). Mor et al. (2019) and Isern (2019) for instance focused on stars (and white dwarfs) within a few hundred parsecs, and both found a peak in the local star formation rate 2–3 Gyr ago. Ruiz-Lara et al. (2020) focused on stars in a wider 2 kpc bubble, and additionally reported three local maxima in the galactic star formation rate, which they associated with pericenter passages of the Sagittarius satellite galaxy.

The right column of Figure 6 compares our derived age distribution against star formation histories (SFHs) reported by Mor et al. (2019) and Ruiz-Lara et al. (2020) based on CMD fitting. The overall slope of both SFHs broadly agrees with what we find from rotation-based ages. The SFH from Mor et al. (2019) seems entirely consistent, particularly after accounting for the statistical uncertainties of that study. Regarding the episodic star formation bursts reported by Ruiz-Lara et al. (2020), they are not apparent in our overall FGK star sample (top row). However, rotation-based ages are most precise for  $\approx \text{G8V-K4V}$  dwarfs older than  $\approx 1.5 \text{ Gyr}$  (Bouma et al. 2023). The lower row of Figure 6 selects these stars with a temperature cut, and may show a hint of the  $\approx 2 \text{ Gyr}$  spike reported by Ruiz-Lara et al. (2020). The  $\approx 1 \text{ Gyr}$  spike however is not recovered. This  $\approx 2.5 \text{ Gyr}$  local maximum is similarly present in the isochrone ages derived by Berger et al. (2020b) for the Kepler field, and in the red giant asteroseismic ages derived by Silva Aguirre et al. (2018). The main novelty of our ages relative to those studies is their improved accuracy at  $< 1 \text{ Gyr}$ .

## 8.2. Caveats & Limitations

Our most constraining ages tend to come from one source of information: rotation. Factors other than stellar age and mass can influence stellar rotation rates. Binarity is one example: even intermediate-separation ( $\sim 10 \text{ AU}$ ) binaries are biased toward rapid rotation (e.g. Meibom et al. 2007, and many studies thereafter). Metallicity may also be relevant (Amard et al. 2020; See et al. 2024). An additional caveat is that our rotation-based ages used photometric effective temperatures (Section 3), even though spectroscopic tempera-

tures are available for the planet-hosting subset. This decision was driven by a desire for homogeneity irrespective of planet-hosting status. However it implies that the estimated planet ages might shift if one were to account for the spectroscopic information.

We were interested in not only the overall age distribution of the Kepler field, but also in the reliability of individual ages for young stars known to host planets. We assessed this reliability using quality bitmasks (Table 1), a comparison against open cluster ages (Section 6), and a comparison between rotation and lithium ages. The open cluster comparison suggested that our ages were generally accurate at  $\ll 0.5 \text{ dex}$  across  $0.04\text{--}2.5 \text{ Gyr}$ , comparable to their stated precision level. Nonetheless, the upper-left panel of Figure 5 shows that outliers do exist, typically at most 10% of the population in each cluster. These stars could be either field star contaminants, or else anomalous stars whose physical rotation histories were altered by processes not captured by our statistical uncertainties. Similarly, although we found  $t_{\text{gyro}}$  and  $t_{\text{Li}}$  to agree for 14 planets orbiting “high-quality” stars, we did find one system, Kepler-786, with radically different lithium ( $228^{+168}_{-87} \text{ Myr}$ ) and rotational ( $4390^{+243}_{-241} \text{ Myr}$ ) ages. Planetary mergers one process that could produce such a signal, but testing this would require a way of measuring differential abundances across a large number of elements.

Other age indicators may help in verifying the ages of the  $\lesssim 3 \text{ Gyr}$  stars that are the focus of this work. Specifically, chromospheric emission in the X-ray, Ca II HK line, and the UV can serve as an age tracer (Mamajek & Hillenbrand 2008; Vidotto et al. 2014; Engle 2024). However, these age indicators are all in a sense “rotation-powered”. The dynamo converts kinetic energy into magnetic energy, which is emitted through these chromospheric pathways. We did not attempt to incorporate these indicators into this study due to concerns regarding sensitivity, homogeneity, and the question of whether they in fact provide age information that is truly independent from rotation.

A broader question, beyond the scope of this work, concerns the ages of stars  $\gtrsim 4 \text{ Gyr}$  old. For age-dating models based only on rotation rates, an important challenge is that the detectability of the rotation signals for 3–10 Gyr stars, especially Sun-like stars, is at the limits of the Kepler data (Masuda 2022). In the “old star” regime, methods using either asteroseismology (van Saders et al. 2016; Saunders et al. 2024), or else using both the evolution of stellar luminosity and stellar rotation (Angus et al. 2019; Claytor et al. 2020; Mathur et al. 2023) seem the most capable of providing useful age constraints for single stars. Incorporating kinematics (Lu et al. 2021) seems likely to remain most useful for stellar ensembles, due to the stochasticity of kinematic heating.

## 8.3. Future Directions

**Occurrence Rates:** The age-dependent trends predicted for exoplanet populations, such as the Kelvin-Helmholtz cooling of mini-Neptunes (Gupta & Schlichting 2019), time-dependent carving of the photoevaporation desert (Owen & Lai 2018), and the time evolution of the radius valley (Rogers

& Owen 2021) can be explored using our data. We defer this analysis to a separate publication. Some care is required since beyond age, exoplanet demographics also depend on stellar metallicity and mass (e.g. Petigura et al. 2018; Miyazaki & Masuda 2023).

*Field star ages from other surveys:* While this study focused on Kepler, other photometric surveys (e.g. K2, TESS, HAT, WASP, NGTS, ZTF, ATLAS) open opportunities for age-dating a far broader set of stars and planets. Future prospects also include PLATO (Rauer et al. 2014), Earth 2.0 (Ge et al. 2022), and the Roman Galactic Bulge Time Domain Survey (Wilson et al. 2023). Rotation-based age-dating from these surveys could yield new insight into whether the star formation history that we see in the Kepler field is universal across the thin disk.

*Searches for new associations:* Some of the youngest stars in Tables 1 and 2 have been linked to their birth clusters, while others have not. A systematic search for the birth clusters of the youngest “field stars” could combine positions and velocities from Gaia with rotation measurements from TESS. Given the  $\sim 100$  Myr decoherence time of young clusters, the youngest “field stars” may have easily identifiable young neighbors.

## 9. CONCLUSIONS

We began this work with two questions: how wrong is the assumption of a uniform age distribution for stars in the galactic thin disk? And why are only  $\approx 50$  sub-gigayear transiting planets known, rather than the  $\approx 500$  that would be expected under the assumption of a uniform star formation history for the  $\approx 5,000$  known planets?

Our approach to answering these questions was to curate a sample of stellar rotation periods, lithium equivalent widths, and temperatures using archival and new data from the Kepler field. We derived new ages using empirical interpolation-based methods, and assessed the reliability of these ages by comparing them against benchmark open clusters (Figure 5).

Tables 1 and 2 summarize the results for planets and stars, respectively. Our recovered ages are accurate for all 14 known Kepler planets in clusters. While lithium provided minimal added information for most of the sample,  $t_{\text{Li}}$  and  $t_{\text{gyro}}$  agreed in  $\gtrsim 90\%$  of cases for which comparison was possible. Our resulting ages included two-sided  $t_{\text{gyro}}$  and  $t_{\text{Li}}$  for 14 sub-gigayear confirmed planets, and two-sided  $t_{\text{gyro}}$  with one-sided  $t_{\text{Li}}$  for 39 sub-gigayear confirmed planets. Allowing for “candidate” planets, grazing transit geometries, and stars with RUWE far from unity expands the counts by a factor of two.

The sizes and periods for the most secure set of planets are shown in Figure 7. While the new young planets are mostly mini-Neptunes, some are near the lower boundary of the “sub-Jovian desert” (Owen & Lai 2018), which could have an evolutionary connection to their youth. Other discoveries include Earth-sized planets with new ages of only a few hundred million years.

Our main conclusions with respect to our original questions are as follows.

1. Rather than being uniform, the age distributions of both the Kepler target stars and the known Kepler planet-hosts show a demographic cliff. There are twice as many “old” (2-3 Gyr) stars in the Kepler field as “young” (0-1 Gyr) stars. The star formation rate today is  $2.85 \pm 0.12$  times lower than it was three billion years ago. This result from rotation-based ages broadly agrees with recent reports of a declining star-formation rate from CMD fitting and white-dwarf chronology, though with the advantage of better sensitivity for FGK stars at  $t < 1$  Gyr (e.g. Figure 5).
2. Rather than expecting  $\approx 500$  exoplanets younger than one billion years, the age distribution of stars in the Kepler field implies that we should instead expect  $\approx 250$ .
3. We have derived rotation-based ages for 60 Kepler planets younger than 1 Gyr at  $2\sigma$ , and 108 planets with median ages below 1 Gyr. Concatenating these planets against the existing literature yields  $\approx 170$  known sub-gigayear planets. This lessens the original factor of ten discrepancy to a factor of at most two.

## ACKNOWLEDGMENTS

The work itself was supported by the Heising-Simons 51 Pegasi b Fellowship (LGB, EKP) and the Arthur R. Adams SURF Fellowship (EKP). The HIRES spectra were obtained at the W. M. Keck Observatory. We recognize the importance that the summit of Maunakea has always had within the indigenous Hawaiian community, and we are deeply grateful for the opportunity to conduct observations from this mountain.

*Contributions:* Per <https://credit.niso.org/>: Conceptualization: LGB. Data curation: LGB, AWH, HI. Formal analysis: LGB, KM. Funding acquisition: LGB. Investigation: LGB, EKP. Methodology: LGB, EKP, KM. Project administration: LGB, LAH, AWH. Resources: LGB, AWH. Software: LGB. Supervision: LGB, LAH. Validation: LGB, KM. Visualization: LGB. Writing – original draft: LGB. Writing – review & editing: all authors.

*Facilities:* Gaia (Gaia Collaboration et al. 2022), Kepler (Borucki et al. 2010), Keck:I (HIRES) (Vogt et al. 1994b), 2MASS (Skrutskie et al. 2006), SDSS (York et al. 2000).

*Software:* astropy (Price-Whelan et al. 2018), claude (Anthropic 2024), eagles (Jeffries et al. 2023), gyro-interp (Bouma et al. 2023), matplotlib (Hunter et al. 2007), numpy (Van Der Walt et al. 2011), scipy (Virtanen et al. 2020).

**Table 1.** Ages of Kepler planets and planet candidates. This version of the table is truncated to include the youngest systems, sorted by the minimum of either the rotation or lithium-based age. The full machine-readable table contains ages and age limits for 2,461 non-false positive KOIs with  $MES > 10$ . A bash script to decode the  $Q_{\text{star}}$  quality flag is [available online](#). A python script to select stars with specific bit flags is [also available](#). All quoted age uncertainties are statistical.

KOI	Kepler	$T_{\text{eff}}$	$P_{\text{rot}}$	$EW_{\text{Li}}^*$	$t_{\text{gyro}}$	$t_{\text{Li}}$	Consistent?	$R_p$	$P_{\text{orb}}$	$Q_{\text{planet}}$	$Q_{\text{star}}$	Spec?	Comment
–	–	K	days	mÅ	Myr	Myr	str	Earths	days	int	int	bool	–
K05245.01	Kepler-1627 b	5357	2.62	$225 \pm 7$	$81_{-55}^{+158}$	$51_{-27}^{+38}$	Yes	3.79	7.2	0	1152	1	Cep-Her
K07368.01	Kepler-1974 b	5068	2.56	$248 \pm 4$	$88_{-60}^{+183}$	$54_{-25}^{+47}$	Yes	2.22	6.84	0	1536	1	Cep-Her
K06228.01	Kepler-1644 b	5521	1.43	$-2 \pm 13$	$77_{-53}^{+144}$	$> 767$	No	1.88	21.09	4	1666	1	Unres. Binary
K06186.01	Kepler-1643 b	4918	5.05	$120 \pm 6$	$79_{-54}^{+182}$	$191_{-76}^{+92}$	Yes	2.11	5.34	0	0	1	Cep-Her
K03933.01	Kepler-1699 b	5496	4.16	$-11 \pm 7$	$85_{-58}^{+106}$	$> 889$	No	1.32	3.49	0	1664	1	Unres. Binary
K03916.01	Kepler-1529 b	4974	6.43	$200 \pm 6$	$109_{-71}^{+117}$	$90_{-39}^{+53}$	Yes	2.01	5.34	0	64	1	
K01804.01	Kepler-957 b	4947	4.52	$24 \pm 9$	$96_{-66}^{+196}$	$> 241$	Yes	6.9	5.91	0	0	1	✓
K07913.01	Kepler-1975 b	4450	3.36	$56 \pm 9$	$96_{-66}^{+223}$	$> 57$	Yes	2.03	24.28	0	1792	1	Cep-Her
K03936.02	Kepler-1930 b	4906	7.1	$170 \pm 4$	$174_{-65}^{+106}$	$115_{-49}^{+55}$	Yes	1.52	13.03	4	0	1	
K03876.01	Kepler-1928 b	5577	4.64	$137 \pm 4$	$148_{-87}^{+102}$	$189_{-94}^{+150}$	Yes	1.86	19.58	0	1024	1	MELANGE-3
K04069.01	Kepler-1938 b	4617	7.82	$6 \pm 16$	$152_{-42}^{+112}$	$> 208$	Yes	1.47	13.06	0	1088	1	
K02678.01	Kepler-1313 b	5236	6.13	$142 \pm 3$	$197_{-91}^{+112}$	$174_{-72}^{+96}$	Yes	1.71	3.83	4	1024	1	
K04194.01	Kepler-1565 b	4958	7.4	$133 \pm 7$	$230_{-85}^{+111}$	$174_{-72}^{+81}$	Yes	1.27	1.54	0	0	1	✓✓
K03835.01	Kepler-1521 b	4806	7.82	$117 \pm 5$	$207_{-68}^{+98}$	$176_{-69}^{+79}$	Yes	2.3	47.15	0	1024	1	✓✓
K01838.01	Kepler-970 b	4314	9.23	$36 \pm 14$	$177_{-39}^{+124}$	$> 92$	Yes	2.15	16.74	4	0	1	MELANGE-3
K00063.01	Kepler-63 b	5486	5.49	$89 \pm 4$	$223_{-92}^{+98}$	$542_{-256}^{+475}$	Yes	5.64	9.43	0	2048	1	✓✓
K01199.01	Kepler-786 b	4680	33.06	$83 \pm 6$	$4390_{-241}^{+243}$	$228_{-87}^{+168}$	No	2.31	53.53	0	0	1	Mystery
K03316.01	Kepler-1467 b	5252	6.31	$122 \pm 6$	$230_{-98}^{+112}$	$236_{-95}^{+151}$	Yes	3.11	47.06	0	64	1	
K01074.01	Kepler-762 b	5921	4.01	$-27 \pm 25$	$244_{-100}^{+113}$	$> 548$	Maybe	15.19	3.77	0	512	1	
K01839.01	Kepler-971 b	5447	6.22	$105 \pm 6$	$306_{-115}^{+95}$	$366_{-164}^{+290}$	Yes	3.93	9.59	0	128	1	
K01833.01	Kepler-968 b	4413	10.46	$10 \pm 18$	$328_{-87}^{+108}$	$> 159$	Yes	1.85	3.69	0	0	1	Theia-520
K01833.03	Kepler-968 c	4413	10.46	$10 \pm 18$	$328_{-87}^{+108}$	$> 159$	Yes	1.63	5.71	0	0	1	Theia-520
K01833.02	Kepler-968 d	4413	10.46	$10 \pm 18$	$328_{-87}^{+108}$	$> 159$	Yes	2.28	7.68	4	0	1	Theia-520
K02675.01	Kepler-1312 b	5584	6.13	$86 \pm 4$	$357_{-109}^{+75}$	$642_{-318}^{+617}$	Yes	2.07	5.45	0	2112	1	
K02675.02	Kepler-1312 c	5584	6.13	$86 \pm 4$	$357_{-109}^{+75}$	$642_{-318}^{+617}$	Yes	0.94	1.12	0	2112	1	
K00775.02	Kepler-52 b	4164	11.85	$22 \pm 18$	$359_{-96}^{+200}$	$> 100$	Yes	2.19	7.88	0	0	1	Theia-520
K00775.01	Kepler-52 c	4164	11.85	$22 \pm 18$	$359_{-96}^{+200}$	$> 100$	Yes	2.04	16.38	0	0	1	Theia-520
K00775.03	Kepler-52 d	4164	11.85	$22 \pm 18$	$359_{-96}^{+200}$	$> 100$	Yes	2.03	36.45	0	0	1	Theia-520
K04004.01	Kepler-1933 b	5576	6.21	$85 \pm 3$	$366_{-109}^{+74}$	$642_{-318}^{+603}$	Yes	1.01	4.94	4	0	1	
K02174.03	Kepler-1802 b	4245	11.45	–	$378_{-109}^{+198}$	–	–	1.71	7.73	4	884	0	
K02174.02	Kepler-1802 c	4245	11.45	–	$378_{-109}^{+198}$	–	–	2.05	33.14	0	884	0	
K03935.01	Kepler-1532 b	5554	6.48	$90 \pm 6$	$397_{-102}^{+71}$	$567_{-278}^{+545}$	Yes	1.26	1.09	0	0	1	✓✓
K01801.01	Kepler-955 b	5221	7.5	$79 \pm 4$	$397_{-131}^{+99}$	$536_{-237}^{+481}$	Yes	2.69	14.53	0	0	1	✓✓
K01800.01	Kepler-447 b	5648	6.4	$103 \pm 3$	$420_{-78}^{+64}$	$405_{-203}^{+355}$	Yes	18.49	7.79	4	1024	1	
K03370.02	Kepler-1481 b	4832	9.11	$22 \pm 7$	$407_{-118}^{+126}$	$> 210$	Yes	1.09	5.94	0	0	1	✓
K04156.01	Kepler-1943 b	6002	–	$99 \pm 5$	–	$409_{-254}^{+520}$	No	1.29	4.85	4	518	1	
K00448.01	Kepler-159 b	4511	10.5	$19 \pm 10$	$415_{-110}^{+159}$	$> 161$	Yes	2.3	10.14	0	2048	1	✓
K00448.02	Kepler-159 c	4511	10.5	$19 \pm 10$	$415_{-110}^{+159}$	$> 161$	Yes	2.75	43.59	0	2048	1	✓
K00046.01	Kepler-101 b	5498	–	$100 \pm 5$	–	$419_{-196}^{+349}$	No	5.9	3.49	0	518	1	
K04169.01	Kepler-1561 b	5742	6.18	$66 \pm 3$	$426_{-78}^{+74}$	$1409_{-788}^{+1718}$	Maybe	0.94	1.01	0	1024	1	✓✓
K02708.01	Kepler-1320 b	4536	10.46	$25 \pm 32$	$429_{-113}^{+172}$	$> 140$	Yes	1.39	0.87	0	0	1	✓
K00119.01	Kepler-108 b	5626	–	$100 \pm 5$	–	$438_{-220}^{+402}$	No	8.2	49.18	0	418	1	
K00119.02	Kepler-108 c	5626	–	$100 \pm 5$	–	$438_{-220}^{+402}$	No	7.78	190.32	4	418	1	
K00323.01	Kepler-523 b	5267	7.6	$49 \pm 4$	$444_{-114}^{+87}$	$1824_{-1047}^{+2706}$	Maybe	1.9	5.84	0	0	1	✓✓
K02115.01	Kepler-67 b	5126	10.39	$83 \pm 10$	$877_{-120}^{+107}$	$458_{-206}^{+351}$	Yes	2.96	15.73	0	0	1	✓✓
K00002.01	Kepler-2 b	6436	–	$83 \pm 4$	–	$485_{-353}^{+924}$	–	16.42	2.2	0	7	1	
K03010.01	Kepler-1410 b	3808	14.19	$-21 \pm 24$	$489_{-166}^{+539}$	$> 80$	Yes	1.39	60.87	0	512	1	

**Table 1 continued**



Table 1 (continued)

K03371.02	Kepler-1482 b	5330	7.7	52 ± 3	491 <sup>+76</sup> <sub>-91</sub>	1630 <sup>+2285</sup> <sub>-904</sub>	Maybe	1.0	12.25	0	640	1	
K03497.01	Kepler-1512 b	4894	9.33	14 ± 8	505 <sup>+135</sup> <sub>-115</sub>	> 295	Yes	0.8	20.36	4	2690	1	
K03864.01	Kepler-1698 b	4866	9.49	2 ± 8	518 <sup>+151</sup> <sub>-123</sub>	> 358	Yes	0.9	1.21	0	0	1	✓
K05447.02	Kepler-1629 b	5585	7.45	–	529 <sup>+62</sup> <sub>-62</sub>	–	–	0.67	3.88	0	0	0	
K04246.01	Kepler-1576 b	5794	7.09	17 ± 8	559 <sup>+102</sup> <sub>-71</sub>	> 613	Yes	0.9	6.98	0	0	1	✓
K03324.01	Kepler-1469 b	5356	8.15	–5 ± 25	562 <sup>+75</sup> <sub>-82</sub>	> 535	Yes	2.53	21.86	0	0	1	✓
K01779.01	Kepler-318 b	5799	7.09	65 ± 3	562 <sup>+105</sup> <sub>-72</sub>	1507 <sup>+1876</sup> <sub>-858</sub>	Yes	3.97	4.66	0	0	1	✓✓
K01779.02	Kepler-318 c	5799	7.09	65 ± 3	562 <sup>+105</sup> <sub>-72</sub>	1507 <sup>+1876</sup> <sub>-858</sub>	Yes	3.1	11.82	4	0	1	
K02035.01	Kepler-1066 b	5847	7.0	60 ± 4	587 <sup>+152</sup> <sub>-84</sub>	2018 <sup>+2616</sup> <sub>-1187</sub>	Maybe	1.96	1.93	4	0	1	
K02084.01	Kepler-1792 b	4942	9.49	13 ± 11	587 <sup>+155</sup> <sub>-144</sub>	> 312	Yes	2.15	4.2	0	0	1	✓
K03274.01	Kepler-1451 b	5675	7.82	42 ± 4	597 <sup>+79</sup> <sub>-65</sub>	> 316	Yes	2.33	35.62	0	0	1	✓
K01615.01	Kepler-908 b	5670	7.88	67 ± 4	602 <sup>+77</sup> <sub>-64</sub>	1317 <sup>+1607</sup> <sub>-724</sub>	Yes	1.36	1.34	0	2560	1	
K02022.01	Kepler-349 b	5756	7.71	64 ± 6	617 <sup>+102</sup> <sub>-72</sub>	1686 <sup>+2185</sup> <sub>-976</sub>	Yes	1.99	5.93	0	0	1	✓✓
K02022.02	Kepler-349 c	5756	7.71	64 ± 6	617 <sup>+102</sup> <sub>-72</sub>	1686 <sup>+2185</sup> <sub>-976</sub>	Yes	1.97	12.25	0	0	1	✓✓
K00620.01	Kepler-51 b	5635	8.14	48 ± 8	623 <sup>+75</sup> <sub>-65</sub>	> 258	Yes	6.62	45.16	0	0	1	✓
K00620.03	Kepler-51 c	5635	8.14	48 ± 8	623 <sup>+75</sup> <sub>-65</sub>	> 258	Yes	5.49	85.32	4	0	1	
K00620.02	Kepler-51 d	5635	8.14	48 ± 8	623 <sup>+75</sup> <sub>-65</sub>	> 258	Yes	9.04	130.18	0	0	1	✓
K02803.01	Kepler-1877 b	5506	8.37	3 ± 12	624 <sup>+69</sup> <sub>-65</sub>	> 726	Maybe	0.55	2.38	0	0	1	✓
K00720.04	Kepler-221 b	5070	9.3	21 ± 5	636 <sup>+120</sup> <sub>-115</sub>	> 341	Yes	1.51	2.8	0	0	1	✓
K00720.01	Kepler-221 c	5070	9.3	21 ± 5	636 <sup>+120</sup> <sub>-115</sub>	> 341	Yes	2.86	5.69	0	0	1	✓
K00720.02	Kepler-221 d	5070	9.3	21 ± 5	636 <sup>+120</sup> <sub>-115</sub>	> 341	Yes	2.57	10.04	0	0	1	✓
K00720.03	Kepler-221 e	5070	9.3	21 ± 5	636 <sup>+120</sup> <sub>-115</sub>	> 341	Yes	2.58	18.37	4	0	1	
K03097.02	Kepler-431 b	6259	16.16	80 ± 3	–	671 <sup>+1092</sup> <sub>-458</sub>	–	0.93	6.8	4	2055	1	
K03097.03	Kepler-431 c	6259	16.16	80 ± 3	–	671 <sup>+1092</sup> <sub>-458</sub>	–	0.93	8.7	4	2055	1	
K03097.01	Kepler-431 d	6259	16.16	80 ± 3	–	671 <sup>+1092</sup> <sub>-458</sub>	–	1.08	11.92	4	2055	1	
K01982.01	Kepler-1781 b	5363	9.17	–	708 <sup>+80</sup> <sub>-78</sub>	–	–	1.95	4.89	0	0	0	
K03375.01	Kepler-1918 b	5522	9.11	–	722 <sup>+75</sup> <sub>-71</sub>	–	–	2.19	47.06	0	0	0	
K01835.02	Kepler-326 b	5142	9.56	8 ± 8	724 <sup>+108</sup> <sub>-103</sub>	> 506	Yes	1.25	2.25	0	640	1	
K01835.01	Kepler-326 c	5142	9.56	8 ± 8	724 <sup>+108</sup> <sub>-103</sub>	> 506	Yes	1.38	4.58	0	640	1	
K01835.03	Kepler-326 d	5142	9.56	8 ± 8	724 <sup>+108</sup> <sub>-103</sub>	> 506	Yes	1.31	6.77	0	640	1	
K01797.01	Kepler-954 b	4736	10.68	4 ± 11	726 <sup>+220</sup> <sub>-183</sub>	> 270	Yes	2.16	16.78	0	0	1	✓
K01821.01	Kepler-963 b	5383	9.42	–	748 <sup>+80</sup> <sub>-78</sub>	–	–	2.64	9.98	0	0	0	
K03681.01	Kepler-1514 b	5852	7.87	69 ± 2	752 <sup>+372</sup> <sub>-138</sub>	1302 <sup>+1622</sup> <sub>-754</sub>	Yes	11.94	217.83	0	516	1	
K03681.02	Kepler-1514 c	5852	7.87	69 ± 2	752 <sup>+372</sup> <sub>-138</sub>	1302 <sup>+1622</sup> <sub>-754</sub>	Yes	1.17	10.51	4	516	1	
K00647.01	Kepler-634 b	6272	–	77 ± 3	–	768 <sup>+1250</sup> <sub>-527</sub>	–	2.13	5.17	0	7	1	
K02037.01	Kepler-1995 b	4746	10.8	9 ± 21	770 <sup>+211</sup> <sub>-193</sub>	> 223	Yes	3.46	73.76	0	2048	1	✓
K01781.02	Kepler-411 b	4920	10.32	4 ± 3	775 <sup>+163</sup> <sub>-166</sub>	> 396	Yes	2.2	3.01	4	0	1	
K01781.01	Kepler-411 c	4920	10.32	4 ± 3	775 <sup>+163</sup> <sub>-166</sub>	> 396	Yes	3.47	7.83	0	0	1	✓
K07375.01	–	4212	3.88	117 ± 9	104 <sup>+235</sup> <sub>-72</sub>	70 <sup>+37</sup> <sub>-21</sub>	Yes	1.73	4.85	0	1600	1	
K03991.01	–	5226	5.22	98 ± 3	71 <sup>+91</sup> <sub>-48</sub>	354 <sup>+253</sup> <sub>-145</sub>	Maybe	1.37	1.57	0	1152	1	
K01546.01	–	5639	0.9	33 ± 24	75 <sup>+134</sup> <sub>-51</sub>	> 316	Maybe	11.92	0.92	0	1152	1	
K05482.01	–	5519	0.81	–	77 <sup>+144</sup> <sub>-53</sub>	–	–	2.96	31.71	4	1666	0	
K00064.01	–	5306	2.23	3 ± 4	81 <sup>+129</sup> <sub>-55</sub>	> 567	No	10.49	1.95	4	2566	1	
K06188.01	–	5209	1.62	6 ± 11	84 <sup>+171</sup> <sub>-58</sub>	> 554	Maybe	2.75	1.65	0	1024	1	
K02695.01	–	5174	2.88	–	86 <sup>+176</sup> <sub>-59</sub>	–	–	20.23	2.5	4	640	0	
K07449.01	–	4928	1.31	–	92 <sup>+194</sup> <sub>-63</sub>	–	–	20.43	1.32	4	1024	0	
K06130.01	–	4560	3.02	–1 ± 25	94 <sup>+217</sup> <sub>-65</sub>	> 186	Yes	1.45	1.54	0	1952	1	
K06195.01	–	4677	1.42	–8 ± 30	94 <sup>+210</sup> <sub>-65</sub>	> 208	Yes	2.28	1.44	0	1024	1	

NOTE—EW<sub>Li</sub><sup>\*</sup> is the lithium equivalent width *after* subtracting a constant 7.5 mÅ to account for the Fe I 6707.44 Å blend (see Section 2.3). Two checkmarks (✓✓) denote confirmed planets with two-sided  $t_{\text{gyro}}$  and  $t_{\text{Li}}$  for which both the age and the planet are expected to be reliable. One checkmark (✓) denotes confirmed planets with two-sided  $t_{\text{gyro}}$  only. “Confirmed” planets appear in the machine-readable version before “candidate” planets. Planetary sizes are mostly drawn from (in order of precedence) [Petigura et al. \(2022\)](#), [Berger et al. \(2020a\)](#), and [Thompson et al. \(2018\)](#), and might be unphysical for grazing planets. The bit quality flags for the rotation-based ages,  $Q_{\text{star}}$ , are described in Section 4.1. Concisely summarized, they are: Bit 0:  $T_{\text{eff}}/\text{K} \in [3800 - 6200]$ ? Bit 1:  $\log g < 4.2$ ? Bit 2:  $M_G < 3.9$  or  $M_G > 8.5$ ? Bit 3: In KEBC? Bit 4: Large  $d_{\text{km, Kep-Gaia}}$ ? Bit 5: Confused Kep-Gaia crossmatch? Bit 6: Gaia DR3 non-single star? Bit 7: RUWE > 1.4? Bit 8: Crowded? Bit 9: Far from main sequence? Bit 10: S21 CP/CB? Bit 11:  $P_{\text{rot}}$  not in the homogeneous S19 S21 sample? As an example, Kepler-1627 has  $Q_{\text{star}}$  flagged with bit 10 and bit 7. The analogous planet quality bitmask,  $Q_{\text{planet}}$ , has the following meaning. Bit 0: Candidate not reliable? Bit 1: MES < 10? Bit 2: Grazing? For a star to have a high likelihood of being “reliable for gyrochronology” we suggest a  $Q_{\text{star}}$  bitmask with bits 0–9 not raised, and for a planet to be “reliable”, we suggest  $Q_{\text{planet}}$  to be zero.

**Table 2.** Ages of Kepler target stars derived from rotation periods. The full machine-readable table includes 53,663 stars with reported rotation periods from S19 and S21, 45,229 of which have finite reported gyrochrone ages. A random set of these stars is shown to give an idea for form and content. The quality bitmask is as in Table 1; requiring bits zero to nine to be null yields 24,426 stars for which gyrochronology is likely to be valid.

KIC	Gaia DR3	$T_{\text{eff}}$	$P_{\text{rot}}$	$t_{\text{gyro}}$	$Q_{\text{star}}$
–	–	K	days	Myr	int
5091688	2101261695298774528	6044	6.52	$1053^{+749}_{-418}$	518
5893899	2076809140819657984	6001	16.52	$3218^{+358}_{-431}$	0
10148324	2128545320427520256	5838	7.89	$872^{+991}_{-276}$	646
1572433	2051763709044322944	4632	35.43	> 4000	0
10864531	2128902558627267200	5342	21.66	$2578^{+154}_{-136}$	512

## REFERENCES

- 1009 Akeson, R. L., Chen, X., Ciardi, D., et al. 2013, *PASP*, 125, 989
- 1010 Alzate, J. A., Bruzual, G., & Díaz-González, D. J. 2021, *MNRAS*,  
1011 501, 302
- 1012 Amard, L., Palacios, A., Charbonnel, C., et al. 2019, *A&A*, 631,  
1013 A77
- 1014 Amard, L., Roquette, J., & Matt, S. P. 2020, *MNRAS*, 499, 3481
- 1015 Angus, R., Aigrain, S., Foreman-Mackey, D., & McQuillan, A.  
1016 2015, *MNRAS*, 450, 1787
- 1017 Angus, R., Morton, T., Aigrain, S., Foreman-Mackey, D., &  
1018 Rajpaul, V. 2018, *MNRAS*, 474, 2094
- 1019 Angus, R., Morton, T. D., Foreman-Mackey, D., et al. 2019, *AJ*,  
1020 158, 173
- 1021 Anthropic. 2024, Claude, v3, Anthropic, Conversational AI model  
1022 used for editing manuscript and generating testable code; the  
1023 authors wrote all of the original manuscript text.  
1024 <https://claude.ai/>
- 1025 Barber, M. G., Mann, A. W., Bush, J. L., et al. 2022, *AJ*, 164, 88
- 1026 Barnes, S. A. 2003, *ApJ*, 586, 464
- 1027 —. 2007, *ApJ*, 669, 1167
- 1028 Berger, T. A., Howard, A. W., & Boesgaard, A. M. 2018, *ApJ*, 855,  
1029 115
- 1030 Berger, T. A., Huber, D., Gaidos, E., van Saders, J. L., & Weiss,  
1031 L. M. 2020a, *AJ*, 160, 108
- 1032 Berger, T. A., Huber, D., van Saders, J. L., et al. 2020b, *AJ*, 159,  
1033 280
- 1034 Binney, J., Dehnen, W., & Bertelli, G. 2000, *MNRAS*, 318, 658
- 1035 Bonomo, A. S., Sozzetti, A., Lovis, C., et al. 2014, *A&A*, 572, A2
- 1036 Borucki, W. J., Koch, D., Basri, G., et al. 2010, *Science*, 327, 977
- 1037 Bouma, L. G., Curtis, J. L., Hartman, J. D., Winn, J. N., & Bakos,  
1038 G. Á. 2021, *AJ*, 162, 197
- 1039 Bouma, L. G., Palumbo, E. K., & Hillenbrand, L. A. 2023, *ApJL*,  
1040 947, L3
- 1041 Bouma, L. G., Hartman, J. D., Brahm, R., et al. 2020, *AJ*, 160, 239
- 1042 Bouma, L. G., Curtis, J. L., Masuda, K., et al. 2022a, *AJ*, 163, 121
- 1043 Bouma, L. G., Kerr, R., Curtis, J. L., et al. 2022b, *AJ*, 164, 215
- 1044 Boyle, A. W., & Bouma, L. G. 2023, *AJ*, 166, 14
- 1045 Cantat-Gaudin, T., Jordi, C., Vallenari, A., et al. 2018, *A&A*, 618,  
1046 A93
- 1047 Cantat-Gaudin, T., Anders, F., Castro-Ginard, A., et al. 2020,  
1048 *A&A*, 640, A1
- 1049 Capistrant, B. K., Soares-Furtado, M., Vanderburg, A., et al. 2024,  
1050 *AJ*, 167, 54
- 1051 Carlos, M., Meléndez, J., Spina, L., et al. 2019, *MNRAS*, 485,  
1052 4052
- 1053 Castro-Ginard, A., Jordi, C., Luri, X., et al. 2018, *A&A*, 618, A59
- 1054 Chaboyer, B., Demarque, P., & Pinsonneault, M. H. 1995, *ApJ*,  
1055 441, 865
- 1056 Chiti, F., van Saders, J. L., Heintz, T. M., et al. 2024, *arXiv*  
1057 e-prints, *arXiv:2403.12129*
- 1058 Choi, J., Dotter, A., Conroy, C., et al. 2016, *ApJ*, 823, 102
- 1059 Christiansen, J. L., Zink, J. K., Hardegree-Ullman, K. K., et al.  
1060 2023, *AJ*, 166, 248
- 1061 Claytor, Z. R., van Saders, J. L., Santos, Â. R. G., et al. 2020, *ApJ*,  
1062 888, 43
- 1063 Curtis, J. 2024, in *American Astronomical Society Meeting*  
1064 Abstracts, Vol. 243, American Astronomical Society Meeting  
1065 Abstracts, 458.08

- Curtis, J. L., Agüeros, M. A., Douglas, S. T., & Meibom, S. 2019, *ApJ*, 879, 49
- Curtis, J. L., Vanderburg, A., Torres, G., et al. 2018, *AJ*, 155, 173
- David, T. J., Petigura, E. A., Luger, R., et al. 2019, *ApJL*, 885, L12
- David, T. J., Contardo, G., Sandoval, A., et al. 2021, *AJ*, 161, 265
- Denissenkov, P. A., Pinsonneault, M., Terndrup, D. M., & Newsham, G. 2010, *ApJ*, 716, 1269
- Dotter, A. 2016, *ApJS*, 222, 8
- Dungee, R., van Saders, J., Gaidos, E., et al. 2022, *ApJ*, 938, 118
- Engle, S. G. 2024, *ApJ*, 960, 62
- Engle, S. G., & Guinan, E. F. 2023, *ApJL*, 954, L50
- Fortney, J. J., Marley, M. S., & Barnes, J. W. 2007, *ApJ*, 659, 1661
- Fritzewski, D. J., Barnes, S. A., James, D. J., & Strassmeier, K. G. 2021, *A&A*, 652, A60
- Fritzewski, D. J., Van Reeth, T., Aerts, C., et al. 2024, *A&A*, 681, A13
- Fulton, B. J., & Petigura, E. A. 2018, *AJ*, 156, 264
- Fulton, B. J., Petigura, E. A., Howard, A. W., et al. 2017, *AJ*, 154, 109
- Gaia Collaboration, Babusiaux, C., van Leeuwen, F., et al. 2018, *A&A*, 616, A10
- Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2022, *arXiv e-prints*, arXiv:2208.00211
- Gallet, F., & Bouvier, J. 2015, *A&A*, 577, A98
- Ge, J., Zhang, H., Zang, W., et al. 2022, *arXiv e-prints*, arXiv:2206.06693
- Gilbert, G. J. 2022, *AJ*, 163, 111
- Gillen, E., Briegal, J. T., Hodgkin, S. T., et al. 2020, *MNRAS*, 492, 1008
- Green, G. M., Schlafly, E., Zucker, C., Speagle, J. S., & Finkbeiner, D. 2019, *ApJ*, 887, 93
- Greiss, S., Steeghs, D., Gänsicke, B. T., et al. 2012, *AJ*, 144, 24
- Gruner, D., Barnes, S. A., & Weingrill, J. 2023, *A&A*, 672, A159
- Gupta, A., & Schlichting, H. E. 2019, *MNRAS*, 487, 24
- Heavens, A., Panter, B., Jimenez, R., & Dunlop, J. 2004, *Nature*, 428, 625
- Hedges, C., Hughes, A., Zhou, G., et al. 2021, *AJ*, 162, 54
- Herschel, J. F. W. 1864, *Philosophical Transactions of the Royal Society of London Series I*, 154, 1
- Hopkins, A. M., & Beacom, J. F. 2006, *ApJ*, 651, 142
- Huber, D., Chaplin, W. J., Christensen-Dalsgaard, J., et al. 2013, *ApJ*, 767, 127
- Huber, D., Zinn, J., Bojsen-Hansen, M., et al. 2017, *ApJ*, 844, 102
- Hunter, J. D., et al. 2007, *Computing in science and engineering*, 9, 90
- Isern, J. 2019, *ApJL*, 878, L11
- Izidoro, A., Ogihara, M., Raymond, S. N., et al. 2017, *MNRAS*, 470, 1750
- Jeffries, R. D., Jackson, R. J., Wright, N. J., et al. 2023, *MNRAS*, 523, 802
- Jenkins, J. M., Caldwell, D. A., & Borucki, W. J. 2002, *ApJ*, 564, 495
- Johnson, J. A., Petigura, E. A., Fulton, B. J., et al. 2017, *AJ*, 154, 108
- Kawaler, S. D. 1989, *ApJL*, 343, L65
- Keppler, M., Benisty, M., Müller, A., et al. 2018, *A&A*, 617, A44
- Kerr, R. M. P., Rizzuto, A. C., Kraus, A. L., & Offner, S. S. R. 2021, *ApJ*, 917, 23
- Kipping, D., & Bakos, G. 2011, *ApJ*, 733, 36
- Kirk, B., Conroy, K., Prša, A., et al. 2016, *AJ*, 151, 68
- Klein, B., Zicher, N., Kavanagh, R. D., et al. 2022, *MNRAS*, 512, 5067
- Kolbl, R., Marcy, G. W., Isaacson, H., & Howard, A. W. 2015, *AJ*, 149, 18
- Kounkel, M., & Covey, K. 2019, *AJ*, 158, 122
- Kounkel, M., Covey, K., & Stassun, K. G. 2020, *AJ*, 160, 279
- Livingston, J. H., Dai, F., Hirano, T., et al. 2018, *AJ*, 155, 115
- Lu, Y., Angus, R., Foreman-Mackey, D., & Hattori, S. 2024, *AJ*, 167, 159
- Lu, Y. L., Angus, R., Curtis, J. L., David, T. J., & Kiman, R. 2021, *AJ*, 161, 189
- Mamajek, E. E., & Hillenbrand, L. A. 2008, *ApJ*, 687, 1264
- Mann, A. W., Gaidos, E., Mace, G. N., et al. 2016, *ApJ*, 818
- Marois, C., Macintosh, B., Barman, T., et al. 2008, *Science*, 322, 1348
- Masuda, K. 2014, *ApJ*, 783, 53
- . 2022, *ApJ*, 937, 94
- Mathur, S., Huber, D., Batalha, N. M., et al. 2017, *ApJS*, 229, 30
- Mathur, S., Claytor, Z. R., Santos, Â. R. G., et al. 2023, *ApJ*, 952, 131
- Matt, S. P., Brun, A. S., Baraffe, I., Bouvier, J., & Chabrier, G. 2015, *ApJL*, 799, L23
- Mazeh, T., Perets, H. B., McQuillan, A., & Goldstein, E. S. 2015, *ApJ*, 801, 3
- McQuillan, A., Mazeh, T., & Aigrain, S. 2013, *ApJL*, 775, L11
- . 2014, *ApJS*, 211, 24
- Meibom, S., Mathieu, R. D., & Stassun, K. G. 2007, *ApJL*, 665, L155
- Meibom, S., Torres, G., Fressin, F., et al. 2013, *Nature*, 499, 55
- Mills, S. M., & Fabrycky, D. C. 2017, *AJ*, 153, 45
- Miyazaki, S., & Masuda, K. 2023, *AJ*, 166, 209
- Mor, R., Robin, A. C., Figueras, F., Roca-Fàbrega, S., & Luri, X. 2019, *A&A*, 624, L1
- Morton, T. D., Bryson, S. T., Coughlin, J. L., et al. 2016, *ApJ*, 822, 86
- Nardiello, D., Malavolta, L., Desidera, S., et al. 2022, *A&A*, 664, A163
- Newton, E. R., Irwin, J., Charbonneau, D., et al. 2016, *ApJ*, 821, 93
- Newton, E. R., Mann, A. W., Kraus, A. L., et al. 2021, *AJ*, 161, 65



- 1165 Nordström, B., Mayor, M., Andersen, J., et al. 2004, *A&A*, 418,  
1166 989
- 1167 Noyes, R. W., Hartmann, L. W., Baliunas, S. L., Duncan, D. K., &  
1168 Vaughan, A. H. 1984, *ApJ*, 279, 763
- 1169 O'Donovan, F. T., Charbonneau, D., Mandushev, G., et al. 2006,  
1170 *ApJL*, 651, L61
- 1171 Owen, J. E. 2019, *Annual Review of Earth and Planetary Sciences*,  
1172 47, 67
- 1173 Owen, J. E., & Lai, D. 2018, *MNRAS*, 479, 5012
- 1174 Pecaú, M. J., & Mamajek, E. E. 2013, *ApJS*, 208, 9
- 1175 Petigura, E. A., Howard, A. W., Marcy, G. W., et al. 2017, *AJ*, 154,  
1176 107
- 1177 Petigura, E. A., Marcy, G. W., Winn, J. N., et al. 2018, *AJ*, 155, 89
- 1178 Petigura, E. A., Rogers, J. G., Isaacson, H., et al. 2022, *AJ*, 163,  
1179 179
- 1180 Plavchan, P., Barclay, T., Gagné, J., et al. 2020, *Nature*, 582, 497
- 1181 Price-Whelan, A. M., Sipőcz, B. M., Günther, H. M., et al. 2018,  
1182 *AJ*, 156, 123
- 1183 Quinn, S. N., White, R. J., Latham, D. W., et al. 2012, *ApJL*, 756,  
1184 L33
- 1185 Rampalli, R., Agüeros, M. A., Curtis, J. L., et al. 2021, *ApJ*, 921,  
1186 167
- 1187 Rauer, H., Catala, C., Aerts, C., et al. 2014, *Experimental*  
1188 *Astronomy*, 38, 249
- 1189 Raymond, S. N., Kokubo, E., Morbidelli, A., Morishima, R., &  
1190 Walsh, K. J. 2014, in *Protostars and Planets VI*, ed. H. Beuther,  
1191 R. S. Klessen, C. P. Dullemond, & T. Henning, 595–618
- 1192 Rebull, L. M., Stauffer, J. R., Hillenbrand, L. A., et al. 2022, *AJ*,  
1193 164, 80
- 1194 Reinhold, T., Bell, K. J., Kuzlewicz, J., Hekker, S., & Shapiro,  
1195 A. I. 2019, *A&A*, 621, A21
- 1196 Reinhold, T., & Gizon, L. 2015, *A&A*, 583, A65
- 1197 Reinhold, T., Shapiro, A. I., Solanki, S. K., & Basri, G. 2023,  
1198 *A&A*, 678, A24
- 1199 Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, *Journal of*  
1200 *Astronomical Telescopes, Instruments, and Systems*, 1, 014003
- 1201 Rizzuto, A. C., Newton, E. R., Mann, A. W., et al. 2020, *AJ*, 160,  
1202 33
- 1203 Rogers, J. G., & Owen, J. E. 2021, *MNRAS*, 503, 1526
- 1204 Ruiz-Lara, T., Gallart, C., Bernard, E. J., & Cassisi, S. 2020,  
1205 *Nature Astronomy*, 4, 965
- 1206 Sandoval, A., Contardo, G., & David, T. J. 2021, *ApJ*, 911, 117
- 1207 Santos, A. R. G., Breton, S. N., Mathur, S., & García, R. A. 2021,  
1208 *ApJS*, 255, 17
- 1209 Santos, A. R. G., García, R. A., Mathur, S., et al. 2019, *ApJS*, 244,  
1210 21
- 1211 Saunders, N., van Saders, J. L., Lyttle, A. J., et al. 2024, *ApJ*, 962,  
1212 138
- 1213 See, V., Yuxi, Lu, Amard, L., & Roquette, J. 2024, *arXiv e-prints*,  
1214 *arXiv:2405.00779*
- 1215 Sestito, P., & Randich, S. 2005, *A&A*, 442, 615
- 1216 Silva Aguirre, V., Bojsen-Hansen, M., Slumstrup, D., et al. 2018,  
1217 *MNRAS*, 475, 5487
- 1218 Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131,  
1219 1163
- 1220 Skumanich, A. 1972, *ApJ*, 171, 565
- 1221 Soderblom, D. R. 2010, *ARA&A*, 48, 581
- 1222 Southworth, J. 2011, *MNRAS*, 417, 2166
- 1223 Sozzetti, A., Torres, G., Charbonneau, D., et al. 2007, *ApJ*, 664,  
1224 1190
- 1225 Spada, F., & Lanzafame, A. C. 2020, *A&A*, 636, A76
- 1226 Sun, L., Ioannidis, P., Gu, S., et al. 2019, *A&A*, 624, A15
- 1227 Thompson, S. E., Coughlin, J. L., Hoffman, K., et al. 2018, *ApJS*,  
1228 235, 38
- 1229 Tofflemire, B. M., Rizzuto, A. C., Newton, E. R., et al. 2021, *AJ*,  
1230 161, 171
- 1231 Tran, Q. H., Bowler, B. P., Cochran, W. D., et al. 2024, *AJ*, 167,  
1232 193
- 1233 Vach, S., Zhou, G., Huang, C. X., et al. 2024, *arXiv e-prints*,  
1234 *arXiv:2403.03261*
- 1235 Van Der Walt, S., Colbert, S. C., & Varoquaux, G. 2011,  
1236 *Computing in Science & Engineering*, 13, 22
- 1237 Van Eylen, V., Agentoft, C., Lundkvist, M. S., et al. 2018,  
1238 *MNRAS*, 479, 4786
- 1239 van Saders, J. L., Ceillier, T., Metcalfe, T. S., et al. 2016, *Nature*,  
1240 529, 181
- 1241 Vidotto, A. A., Gregory, S. G., Jardine, M., et al. 2014, *MNRAS*,  
1242 441, 2361
- 1243 Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *Nature*  
1244 *Methods*, 17, 261
- 1245 Vogt, S. S., Allen, S. L., Bigelow, B. C., et al. 1994a, *SPIE*  
1246 *Conference Series*, ed. D. L. Crawford & E. R. Craine, Vol. 2198
- 1247 Vogt, S. S., Allen, S. L., Bigelow, B. C., et al. 1994b, in *Society of*  
1248 *Photo-Optical Instrumentation Engineers (SPIE) Conference*  
1249 *Series*, Vol. 2198, *Society of Photo-Optical Instrumentation*  
1250 *Engineers (SPIE) Conference Series*, ed. D. L. Crawford & E. R.  
1251 Craine, 362
- 1252 Walkowicz, L. M., & Basri, G. S. 2013, *MNRAS*, 436, 1883
- 1253 Wilson, R. F., Barclay, T., Powell, B. P., et al. 2023, *ApJS*, 269, 5
- 1254 Winn, J. N., Johnson, J. A., Narita, N., et al. 2008, *ApJ*, 682, 1283
- 1255 Wood, M. L., Mann, A. W., Barber, M. G., et al. 2023, *AJ*, 165, 85
- 1256 Xiang, M., & Rix, H.-W. 2022, *Nature*, 603, 599
- 1257 York, D. G., Adelman, J., Anderson, John E., J., et al. 2000, *AJ*,  
1258 120, 1579
- 1259 Zakhochay, O. V., Launhardt, R., Trifonov, T., et al. 2022, *A&A*,  
1260 667, L14
- 1261 Zari, E., Hashemi, H., Brown, A. G. A., Jardine, K., & de Zeeuw,  
1262 P. T. 2018, *A&A*, 620, A172
- 1263 Zhou, G., Wirth, C. P., Huang, C. X., et al. 2022, *AJ*, 163, 289
- 1264 Ziegler, C., Law, N. M., Morton, T., et al. 2017, *AJ*, 153, 66

## APPENDIX

## A. ROTATION PERIOD CATALOG COMPARISON


While we opted for the S19 and S21 rotation period catalogs, the recent analysis by Reinhold et al. (2023) bears comment. Reinhold et al. (2023) used a similar selection function as Santos et al. (all Kepler stars), and considered a novel period measurement approach based on the gradient of the power spectrum (GPS). Their method likely provides greater completeness, due to its greater sensitivity to signals that are only weakly periodic.

Figure 8 shows histograms of the differences between reported periods from these various studies. The left panel compares overlapping stars from McQuillan et al. (2014) and the Santos et al. studies. The periods agree at a precision of  $\lesssim 0.01 P_{\text{rot}}$  for  $P_{\text{rot}} \lesssim 15$  days, and at  $\lesssim 0.03 P_{\text{rot}}$  for  $P_{\text{rot}} \approx 30$  days. A small bias exists at  $P_{\text{rot}} \approx 10$  days, in the sense that the Santos periods tend to be  $\approx 1\%$  faster for such stars.

The right panel compares overlapping periods from the Reinhold et al. (2023) GPS method and the Santos et al. studies. The periods agree at a precision of  $\lesssim 0.20 P_{\text{rot}}$  for  $P_{\text{rot}} \lesssim 15$  days, and at  $\lesssim 0.30 P_{\text{rot}}$  for  $P_{\text{rot}} \approx 30$  days. A bias develops past  $P_{\text{rot}} \gtrsim 30$  days, in the sense that the Santos et al. periods tend to be 10-20% longer than the GPS periods. The origin of this larger scatter could be connected to the need for the GPS method to determine the “ $\alpha$  calibration factor”, which is known only at a statistical level for large stellar populations (see Reinhold et al. 2023). While we considered deriving independent rotation-based ages using the Reinhold et al. (2023) GPS periods with an inflated empirical uncertainty, we ultimately opted against this path. A rotation period difference of 20-30% away from the “true period” could sufficiently skew a star’s age that we prefer to restrict our attention to the stars for which precision is a possible outcome.

## B. DEFINING AN “ISOCRONALLY FAR FROM THE MAIN SEQUENCE” BOUNDARY

The (arbitrary) method used to construct the orange line in the lower panel of Figure 1 was as follows. The line is defined as the maximum of two numerically-determined functions,  $f_0(T_{\text{eff}})$  and  $f_1(T_{\text{eff}})$ . We defined  $f_0$  by fitting an  $N^{\text{th}}$ -order polynomial to the stars with rotation periods for which  $t_{\text{B20,iso}}/(\sigma_{t,\text{B20,iso}}) \approx 3$  and  $T_{\text{eff}} \in [3700, 5900]$  K, where  $t_{\text{B20,iso}}$  was the isochronal age reported by Berger et al. (2020b). We let the order of the fit vary from  $N=1$  to 10, minimized the Bayesian Information Criterion, and found  $N=6$ . By eye, this yielded a plausible locus for  $T_{\text{eff}} \lesssim 5300$  K. However, for F and G dwarfs, the resulting locus allowed for stars that were “too evolved”; the density of KOIs was anomalously low in this region of the  $\log g$  vs.  $T_{\text{eff}}$  plane. We therefore visually selected KOIs that appeared to be near the main sequence – i.e. most of the yellow points in Figure 1 – and used them to fit a separate polynomial  $f_1$  through a similar BIC-minimization procedure, which yielded  $N=3$ . This polynomial,  $f_1$ , is shown in very faint opacity in Figure 1. The portion of the orange locus from  $\approx 5300$ – $6200$  K is set as  $f_1 + c$ , for  $c$  a constant offset that we defined to be 0.1 dex above the “KOI main sequence”. The exact break-point in temperature is automatically set by



**Figure 8. Comparison of reported literature stellar rotation periods.** “Santos” refers to the concatenation of Santos et al. (2019) and Santos et al. (2021). “M14” refers to McQuillan et al. (2014). “R23GPS” refers to the Gradient of Power Spectrum periods from Reinhold et al. (2023). The background is a 2-D histogram with a logarithmic color stretch that counts overlapping stars between these studies. The black errorbars show the median and  $\pm 1\sigma$  range of period the differences in 1-day bins. The solid gray “empirical  $\sigma_P/P$ ” line in the left panel shows the expected  $\pm 1\sigma$  range that would be spanned if the independent  $P_{\text{rot}}$  measurements had gaussian uncertainties drawn following the empirical estimate described in Section 2.1. Note the different vertical scales.

max( $f_0, f_1 + c$ ). While we do report rotation-based ages for stars above this orange locus, they are flagged as not being near the main sequence in the  $\log g$  vs.  $T_{\text{eff}}$  plane.

### C. CASES WITH “INCONSISTENT” ROTATION AND LITHIUM-BASED AGES

In this section we discuss the systems with confirmed planets that have nominally discrepant rotation and lithium-based ages. We find that only one of them, Kepler-786, is interesting.

*Kepler-1* — TrES-2/Kepler-1 (O’Donovan et al. 2006) is a  $\approx 2.5$  day near-grazing hot Jupiter orbiting a G0 V primary which has been studied in detail by multiple investigators (e.g. Sozzetti et al. 2007; Winn et al. 2008; Kipping & Bakos 2011; Southworth 2011). Our lithium-based age of this system ( $785^{+845}_{-419}$  Myr), derived from  $\text{EW}_{\text{Li}}^* = 81 \pm 3$  mÅ, qualitatively agrees with the previously noted lithium abundance (Sozzetti et al. 2007). However, we detect no photometric rotation signal, and the spectroscopic  $v \sin i$  is low. The lithium age is strongly asymmetric because of the scatter in  $\text{EW}_{\text{Li}}$  vs.  $T_{\text{eff}}$  over ages  $t \gtrsim 1$  Gyr for early G dwarfs. We calculate a  $2\sigma$  upper limit on  $t_{\text{Li}}$  of 2.8 Gyr, and a  $3\sigma$  upper limit of 6.1 Gyr. The non-detection of rotation, while mildly surprising, is not shocking if the system’s true age is in this long tail.

*Kepler-101* — This  $\approx 5500$  K star has  $\text{EW}_{\text{Li}}^* = 100 \pm 5$  mÅ, nominally implying  $t_{\text{Li}} \approx 200$ –800 Myr, but no rotation detection. Our automated flags noted that this star has a low surface gravity, high luminosity, and is far from the main sequence. Bonomo et al. (2014) concur: this star is a sub-giant slightly more massive than the Sun; no rotation detection is expected; the lithium likely simply survived over the star’s main sequence lifetime.

*Kepler-108* — This system of two mutually inclined giant planets (Mills & Fabrycky 2017) has a  $T_{\text{eff}} \approx 5600$  K host star with a strong lithium detection, and no rotation detection. The host star is entering the red giant branch; this has been previously noted through asteroseismology (Huber et al. 2013), and our analysis of the Keck/HIRES spectrum with SpecMatch-Synth (Petigura et al. 2017) concurs, yielding  $T_{\text{eff}} = 5668 \pm 100$  K,  $\log g = 3.8 \pm 0.1$ ,  $[\text{Fe}/\text{H}] = 0.38 \pm 0.06$ , and  $v \sin i = 3.2 \pm 1.0$  km s $^{-1}$ . Given the star’s high mass, the high lithium content and lack of rotation detection is not surprising.

*Kepler-786* — This K3 V star has a surprisingly high lithium content. The spectrum is single-lined, with SpecMatch-Synth derived parameters of  $T_{\text{eff}} = 4769 \pm 100$  K,  $\log g = 4.5 \pm 0.1$ ,  $[\text{Fe}/\text{H}] = 0.11 \pm 0.06$ , and  $v \sin i = 2.4 \pm 0.6$  km s $^{-1}$ . Yet with  $\text{EW}_{\text{Li}}^* = 83 \pm 6$  mÅ, the lithium age of  $228^{+168}_{-87}$  Myr would predict an obvious rotation signal. None is present in the Kepler light curve, consistent with the low  $v \sin i$ . Out of all “apparently discrepant” lithium and rotation measurements discussed in this appendix, this is the only one that seems to remain discrepant after scrutiny. The Ca II doublet is in emission, with  $R'_{\text{HK}} = -4.7 \pm 0.5$ , which suggests an age at least as old as the Hyades (Mamajek & Hillenbrand 2008). The Balmer lines are in absorption, and display no obvious signatures of youth.


*Kepler-1644* — The rotation-based age of this system ( $77^{+144}_{-53}$  Myr) is nominally much lower than the lithium limit ( $> 767$  Myr). The Kepler light curve shows a  $\approx 1\%$  amplitude 1.4 day rotation signal with many flares. However the automated quality flags note that the star has low (photometric) surface gravity, a high RUWE ( $\text{RUWE}_{\text{DR3}} = 7.0$ ), and that it is far from the main sequence. The Keck/HIRES spectrum also shows visually narrow lines, with a SpecMatch-Synth  $v \sin i \leq 2$  km s $^{-1}$ . However, we performed a cross-correlation between the HIRES spectrum and the nearest matches in the Keck/HIRES template library (Kolbl et al. 2015), and found that on 31 July 2022 (UT) the system displayed a broad CCF with a blended second component at  $\approx +26$  km s $^{-1}$  relative to the primary, with a best-fit flux-ratio of  $\approx 3.4\%$ , and a preferred  $T_{\text{eff,B}} \approx 4400$  K. The spectrum and astrometric excess noise therefore point to this system being an unresolved binary, which calls the reliability of the rotation-based age into question. The non-detection of the companion from Robo-AO imaging at Palomar (Ziegler et al. 2017) suggests that the companion(s) are likely within  $\rho \lesssim 0.3''$ .

*Kepler-1699* — This system is in a similar qualitative regime as Kepler-1644, with an apparently young  $t_{\text{gyro}} = 85^{+106}_{-58}$  Myr derived from a  $\approx 2\%$  amplitude 4.2 day rotation signal, and no evidence for lithium. The system has  $\text{RUWE}_{\text{DR3}} = 19.1$ . From the same style of CCF analysis from the Keck/HIRES spectrum acquired on 31 Aug 2022 (UT), we also find a double-peaked CCF, in this case with a secondary component at  $\approx -16$  km s $^{-1}$  relative to the primary with  $T_{\text{eff,B}} \approx 4900$  K. This putative companion is similarly not detected in high-resolution imaging (Ziegler et al. 2017). The rotation-based age is questionable, given that we do not know which source the rotation signal is from, or whether the stars have interacted.

*Kepler-1943* — This system nominally has a  $409^{+520}_{-254}$  Myr lithium age, and no reported rotation detection. However, the star is flagged as being over-luminous, low-surface gravity, and far from the main sequence. In other words, it is a subgiant.

*Kepler-639, Kepler-320, Kepler-1719, Kepler-1876, Kepler-1072, Kepler-1743, Kepler-1929, Kepler-1488* — These eight systems all have nominally two-sided lithium age posteriors between 1 and 3 Gyr, and yet lack rotation period detections. All are subgiants, flagged with  $Q_{\text{star}}$  under various combinations of bits 1, 2, and 9.





**Figure 9. Rotation-based age vs. effective temperature.** The entire Kepler sample is shown on the left. KOI hosts are shown on the right. Each bar shows the  $\pm 1\sigma$  uncertainty for a star’s age as derived from its rotation period. Notable features are discussed in Appendix E.

#### D. KEPLER-411

Kepler-411, a  $t_{\text{gyro}} \approx 780 \pm 170$  Myr early K dwarf with three transiting planets at 3.0, 7.8, and 58 days, has a rare architecture. [Sun et al. \(2019\)](#) conducted a transiting timing analysis of the system, confirming Kepler-411d and arguing for the presence of a fourth non-transiting planet, Kepler-411e, based on the TTVs seen in Kepler-411d and assuming that the system is nearly coplanar. Regarding the stellar age, [Sun et al. \(2019\)](#) used the [Barnes \(2007\)](#) relation and the measured rotation period, which yielded an estimate of  $212 \pm 31$  Myr. However, the star’s temperature ( $\approx 4900$  K) puts it in a regime where the [Barnes](#) relation has been known to not fit observations for a while (e.g. [Mamajek & Hillenbrand 2008](#), Fig. 9). Our adopted age of  $\approx 780 \pm 170$  Myr agrees with the open cluster calibrators.

#### E. AGE DIAGNOSTICS

Figure 9 is a visualization of  $t_{\text{gyro}}$  vs.  $T_{\text{eff}}$  for our rotation-based age catalog. Each bar denotes the  $\pm 1\sigma$  uncertainty for a star’s rotation-based age: this plot can be viewed as the transformation of the left panel of Figure 2. The overdensity of  $P_{\text{rot}} \approx 20$  day G dwarfs corresponds to the large overdensity at  $t_{\text{gyro}} \approx 3$  Gyr. The (non-physical) deficit of  $\sim 22$  day rotators appears as a deficit between 5500–6000 K. A second deficit is also visible from 3800–5000 K; it is associated with the “intermediate period gap”, which may be associated with a transition from spot-dominated to faculae-dominated light curves (e.g. [Reinhold et al. 2019](#)). At  $\approx 1$  Gyr, stalled spin-down yields larger uncertainties for K dwarfs. The pile-up near  $\approx 5$  Gyr is imposed by our choice of prior; given that `gyro_interp` returns non-calibrated ages in this regime, we truncated our analysis at 4 Gyr.