

## Ages of Stars and Planets in the Kepler Field Over the First Four Billion Years

LUKE G. BOUMA,<sup>1,2,\*</sup> LYNNE A. HILLENBRAND,<sup>1</sup> ANDREW W. HOWARD,<sup>1</sup> HOWARD ISAACSON,<sup>3</sup> KENTO MASUDA,<sup>4</sup> AND  
ELSA K. PALUMBO<sup>1,5</sup>

<sup>1</sup>*Department of Astronomy, California Institute of Technology, Pasadena, CA 91125, USA*

<sup>2</sup>*Observatories of the Carnegie Institution for Science, Pasadena, CA 91101, USA*

<sup>3</sup>*Astronomy Department, University of California, Berkeley, CA 94720, USA*

<sup>4</sup>*Department of Earth and Space Science, Osaka University, Osaka 560-0043, Japan*

<sup>5</sup>*Department of Statistics & Data Science, Carnegie Mellon University, Pittsburgh, PA 15213, USA*

(Received 2024 June 21; Revised 2024 September 4; Accepted —)

### ABSTRACT

Recent analyses of FGK stars in open clusters have helped clarify the precision with which a star’s rotation rate and lithium content can be used as empirical indicators for its age. Here we apply this knowledge to stars observed by Kepler. Rotation periods are drawn from previous work; lithium is measured from new and archival Keck/HIRES spectra. We report rotation-based ages for 23,813 stars (harboring 795 known planets) for which our method is applicable. We find that our rotational ages (~~Deleted: accurately~~) recover the ages of stars in open clusters spanning 0.04–2.5 Gyr; they also agree with  $\gtrsim 90\%$  of the independent lithium ages. The resulting yield includes 63 planets younger than 1 Gyr at  $2\sigma$ , and 109 with median ages below 1 Gyr. This is about half the number expected under the classic assumption of a uniform star formation history. (~~Deleted: We find that the scarcity of sub-gigayear systems can be attributed to the star formation rate in the Kepler field dropping by a factor of  $2.81 \pm 0.12$  over the past 3 Gyr. We observe this trend both for known planet hosts and for the parent stellar sample. This “demographic cliff” in the Galaxy’s star formation history has been previously reported, and its confirmation helps clarify the age distribution of the known transiting exoplanets.~~) (Added: The age distribution that we observe, rather than being uniform, shows that the youngest stars in the Kepler field are 3–5 times rarer than stars 3 Gyr old. This trend holds for both known planet hosts and for the parent stellar sample. We attribute this “demographic cliff” to a combination of kinematic heating and a declining star formation rate in the Galaxy’s thin disk, and highlight its impact on the age distribution of known transiting exoplanets.)

**Keywords:** Stellar ages (1581), Planet hosting stars (1242), Field stars (2103), Exoplanet evolution (491), Milky Way evolution (1052)

### 1. INTRODUCTION

Exoplanet science is thriving, fueled by the discovery of thousands of worlds orbiting close to their host stars (Borucki et al. 2010; Ricker et al. 2015). However, most known exoplanets are billions of years old. This fact leaves many gaps in our knowledge of the exact physical and dynamical origins of these objects. The reason is that processes such as thermal cooling (Fortney et al. 2007), atmospheric loss (Owen 2019), giant impacts (Raymond et al. 2014), and dynamical instabilities (Izidoro et al. 2017) are expected to be most efficient over timescales of much less than 1 Gyr. For most known exoplanets, these processes have run their course.

Young ( $<1$  Gyr) exoplanets represent one means of building the timeline for exoplanet evolution. Informative individual exemplars include HIP 67522 b, a Jupiter-sized planet with a sub-Neptunian mass (Rizzuto et al. 2020; Thao et al. submitted), and V1298 Tau, a resonant chain of puffy planets that is a likely precursor to the compact multiplanet systems (David et al. 2019). Population-level analyses have similarly suggested differences in the size distribution of young exoplanets relative to their older counterparts (Berger et al. 2020a; David et al. 2021; Sandoval et al. 2021; Christiansen et al. 2023; Vach et al. 2024).

Discovering a young planet requires solving two problems: finding the planet and measuring the star’s age. Each problem admits a range of solutions (e.g. Marois et al. 2008; Quinn et al. 2012; Tran et al. 2024). In this article we will consider planets whose existence has been previously established using transits, and infer new stellar ages using rotation and lithium.

Corresponding author: Luke G. Bouma  
bouma.luke@gmail.com

\* 51 Pegasi b Fellow; Carnegie Fellow

To begin, imagine that the ages of nearby stars in the Galaxy are uniformly distributed from 0–10 Gyr (Binney et al. 2000; Nordström et al. 2004). This approximation suggests that  $\approx 1\%$  of nearby stars have  $t < 100$  Myr, and  $\approx 10\%$  have  $t < 1$  Gyr. Studies of currently forming protoplanets (Keppler et al. 2018), and of exoplanets evolving just after disk dispersal (e.g. Klein et al. 2022), are thereby capped in their maximum achievable sample sizes by the tyranny of the galactic star formation rate.

Despite this and other observational challenges, young close-in planet discovery has matured over the past decade, primarily due to Kepler, K2, TESS, and Gaia (e.g. Meibom et al. 2013; Mann et al. 2016; Curtis et al. 2018; Livingston et al. 2018; Bouma et al. 2020; Plavchan et al. 2020; Newton et al. 2021; Nardiello et al. 2022; Barber et al. 2022; Zhou et al. 2022; Zakhozhay et al. 2022; Wood et al. 2023). The strategy pursued by most groups during the 2010s was to focus on stars with known ages—obvious members of open clusters—and to search them for transiting planets. The resulting stellar, and assumed planetary, ages are precise at the  $\approx 10\%$  level. An alternative strategy, facilitated by Gaia, is to select (~~Deleted: transiting~~) planets based on empirical youth indicators, and to then search their local neighborhoods for comoving (~~Added: stellar~~) companions (e.g. Tofflemire et al. 2021).

While stellar ensembles provide the gold standard for astronomical ages, this “cluster-first” approach has a major limitation: most stars are in the field. Only  $\lesssim 1\%$  of stars within  $\approx 500$  pc have been associated with their birth cluster (e.g. Zari et al. 2018; Cantat-Gaudin et al. 2020; Kounkel et al. 2020; Kerr et al. 2021). The implication for known planetary ages can be appreciated by querying the NASA Exoplanet Archive (NEA; Akeson et al. 2013) for transiting planets younger than 1 Gyr. Requiring  $t < 1$  Gyr at  $2\sigma$  precision gives  $\approx 50$  (~~Deleted: such~~) planets at the time of writing. Most of these young planets are in clusters, and were found by Kepler, K2, or TESS. However, these surveys have cumulatively discovered (~~Deleted: a total of~~)  $\approx 5,000$  planets. Assuming a constant star formation history, we would expect an order of magnitude more sub-gigayear transiting planets (~~Added: than the 50 currently known~~).

This study aims to resolve two questions. First, how wrong is it to assume a uniform age distribution for transiting planet host stars? Second, where are the missing young planets? We will find that uniform is wrong by a factor of a few, and that stellar activity may be a less significant bias in young transiting planet detection than the challenge of precisely determining ages for field stars.

Despite Kepler’s main mission ending quite some time ago, the ages of many Kepler planets remain uncertain. While isochrone ages have been calculated for Kepler stars (Berger et al. 2020a) and Kepler Objects of Interest (KOIs; Petigura et al. 2022), such ages are most precise for stars whose luminosities and temperatures separate them from the main sequence. For sub-gigayear stars on the main sequence, isochrone ages are therefore limited.

Stellar rotation periods offer a promising alternative. The idea of using a star’s spin-down as a clock has a rich history (Skumanich 1972; Noyes et al. 1984; Kawaler 1989; Barnes 2003; Mamajek & Hillenbrand 2008; Angus et al. 2015; Engle & Guinan 2023). Empirical models now yield ages precise to  $\lesssim 30\%$  for FGK stars between 1–4 Gyr, and constraining age posteriors for younger ages (Bouma et al. 2023). Physics-based models (Matt et al. 2015; Gallet & Bouvier 2015; Spada & Lanzafame 2020) can connect these empirical relations to the evolution of stellar winds, internal structure, and angular momentum transport.

Rotational ages have been reported for various subsets of Kepler stars since the early data releases (e.g. Walkowicz & Basri 2013; McQuillan et al. 2014; Reinhold & Gizon 2015; Angus et al. 2018). More recent work has further explored incorporating information from stellar kinematics (Lu et al. 2021, 2024), and from stellar colors, luminosities, and starspot amplitudes (Mathur et al. 2023). Our analysis is motivated by a few factors that can yield improvements, particularly for sub-gigayear stars. These factors are as follows.

- i) The Kepler Object of Interest catalog, and our vetting of false positives within it, has now reached maturity (e.g. Thompson et al. 2018).
- ii) Measured rotation periods of FGK stars in open clusters now show not only the average evolution of  $P_{\text{rot}}(T_{\text{eff}}, t)$ , but also how the astrophysical dispersion of stars around this average converges by the  $\approx 700$  Myr age of Praesepe (e.g. Curtis et al. 2019; Gillen et al. 2020; Rampalli et al. 2021; Fritzewski et al. 2021; Rebull et al. 2022; Dungee et al. 2022; Boyle & Bouma 2023).
- iii) Using open cluster data, we can marginalize over the range of ages that might explain any one star’s rotation period (Bouma et al. 2023). This represents an improvement in the accuracy of uncertainty propagation relative to previous calibrations.
- iv) We can identify (~~Replaced: binary stars~~ replaced with: **stars that are binaries and that have high metallicity**) with greater fidelity (~~Deleted: -than-in-the-past~~), which can clarify otherwise problematic estimates of rotation-based ages.
- v) New open clusters in various stages of dissolution have been found in the Kepler field (e.g. Kounkel & Covey 2019; Barber et al. 2022; Kerr et al. 2024). These clusters offer stars that we can use to test the reliability of (~~Replaced: the available single-star~~ replaced with: **our**) age-dating methods.

Recent years have also yielded improvements in the lithium age scale. Lithium ages include depletion boundary ages for M dwarfs and brown dwarfs in star clusters, and decline-based ages for individual field FGK stars (Soderblom 2010). The latter approach relies on the observed decrease of Li abundances in partially-convective stars as they age (e.g. Sestito & Randich 2005). The theoretical explanation for these observations is debated (e.g. Chaboyer et al. 1995; Denissenkov et al. 2010; Carlos et al. 2019). Empirical understanding however has improved due to work by Jeffries

et al. (2023), who modeled the time evolution of the Li I 6708 Å equivalent width (EW) using a set of 6,200 stars in 52 open clusters. Two-sided lithium ages are useful for Kepler (FGK) stars between  $\approx 0.03$ –0.5 Gyr, though with a strong dependence on spectral type. The precision of lithium ages in this regime are now in the range of 0.3–0.5 dex.

We discuss our method for selecting the star and planet samples in Section 2, and describe the origin of our adopted stellar parameters other than ages in Section 3. We present our age-dating methods in Sections 4 and 5, and test them in Section 6 using clusters in the Kepler field. We discuss population-level trends in Sections 7 and 8, and offer a few conclusions in Section 9.


## 2. SELECTING THE STARS AND PLANETS

This work is focused on Kepler stars for which ages can be inferred using either rotation, lithium, or both. Such stars are a minority of the  $\approx 160,000$  Kepler targets. Rotation periods have been reported for roughly one in three Kepler targets (e.g. McQuillan et al. 2014; Santos et al. 2021). High-resolution spectra suitable for measurement of the Li I 6708 Å doublet have only been acquired for the Kepler objects of interest (KOIs), which comprise a few percent of Kepler’s targets. In the following, we will describe the set of stars that we adopt with measured rotation periods (Section 2.1), the set of objects we adopt as hosting planets (Section 2.2), and the subset of these with high-resolution spectra suitable for lithium analysis (Section 2.3). Figure 1 provides a visualization of the various samples.

### 2.1. Stellar Rotation Periods


To select stars with rotation periods, we turn to previous work. Many investigators have derived rotation periods both specifically for KOIs (McQuillan et al. 2013; Walkowicz & Basri 2013; Mazeh et al. 2015; Angus et al. 2018; David et al. 2021), and also for the broader (Replaced: set of all replaced with: sample of) Kepler target stars (McQuillan et al. 2014; Reinhold & Gizon 2015; Santos et al. 2019, 2021; Reinhold et al. 2023). These studies used a range of detection methods and selection functions. We are interested in understanding the age distribution of all Kepler target stars irrespective of KOI status. Out of these studies, the most homogeneous (Deleted: and precise-) analyses of both Kepler targets and KOIs appear to be those by Santos et al. (2019, 2021), hereafter S19 and S21. For discussion of the work by Reinhold et al. (2023), see Appendix A. S19 and S21 combined a wavelet analysis and autocorrelation-based approach, and cumulatively reported rotation periods for 55,232 main-sequence and subgiant FGKM stars. They included known KOIs and binaries in their analysis, and removed transits and eclipses during the stellar rotation measurement process.

We therefore adopt the results of S19 and S21 as our default rotation periods. S21 provided a comparison against McQuillan et al. (2014) (hereafter M14); the brief summary is that the periods agree for 99.0% of the 31,038 period detections in common between the two studies. S21 classified the 2,992 remaining stars from M14 as not showing rotation



**Figure 1. The stars.** Our analysis focuses on stars observed by Kepler with previously reported rotation periods (gray points). The rotation periods are primarily drawn from Santos et al. (2019, 2021). About half of the stars with rotation periods are suitable for gyrochronology (dark gray (Deleted: -points)), based on factors including their (Added: apparent) non-binarity and proximity to the main sequence (orange line in (Replaced: lower replaced with: upper) panel; see Section 4.1). Some host “confirmed” or “candidate” KOIs that meet additional planetary quality criteria (yellow points; Section 4.2). Surface gravities and effective temperatures were derived photometrically by Berger et al. (2020b). Isochrones in the (Replaced: lower replaced with: upper) panel are from MIST (Choi et al. 2016).

periods based on updated knowledge of contaminants (such as giant stars and eclipsing binaries) and visual inspection. In addition, S21 reported rotation periods for 24,182 main-



**Figure 2.** Rotation periods for Kepler target stars (left) and known planet hosts (right). The gray lines are (Replaced: “mean fits” to the rotation sequences of open clusters replaced with: polynomials fitted to the slow rotation sequences of open clusters). The 2-D histogram of the stellar sample (left) shows only apparently single stars near the main sequence with  $\log g > 4.2$ ,  $\text{RUWE} < 1.4$ , and temperatures of 3800–6200 K. The planet hosts (right) require the same stellar cuts, and include only the confirmed and candidate planets described in Section 4.2. The total number of stars  $N_s$  and planets  $N_p$  are noted in the relevant panels. A simple estimate for the number of young planets discovered by Kepler follows by counting points below the (Deleted: mean-) rotational isochrones.

sequence and subgiant FGKM stars that were not reported as periodic by M14. Many of these reported signals have lower variability amplitudes and longer periods than those reported by M14.

Analyzing the compilation of S19 and S21 rotation periods for the KOI hosts, we noticed that some known KOIs with rotation periods were missing. This is not surprising, since the rotation periods of KOIs have received more scrutiny than those of ordinary Kepler stars. We therefore decided to split our subsequent analysis into a homogeneous portion that used only the S19 and S21 data, and an inhomogeneous portion that also considered a broader set of available KOI rotation periods. For the latter portion, we first included 32 KOIs with orbital and rotation periods within  $\approx 20\%$  that had been excluded from the S19 and S21 catalogs (A. Santos, private communication). We then incorporated an additional 178 rotation periods for KOI hosts that David et al. (2021) described as either “reliable” or “highly reliable” in their visual analysis of previously reported KOI rotation periods from McQuillan et al. (2013), Walkowicz & Basri (2013), Mazeh et al. (2015) and Angus et al. (2018). Inclusion of these additional KOI rotation periods is a supplementary measure aimed at completeness in our final KOI age catalog; the provenances of the individual adopted periods are noted in the relevant tables.

Finally, since our scope is focused on rotation-based ages, we restricted our attention to stars with reported  $P_{\text{rot}} < 45$  days. The slowest-rotating FGK stars in the open clusters used to calibrate our gyrochronology model have  $P_{\text{rot}} \approx 35$  days. Figure 2 shows a subset of the resulting 53,663

Kepler stars with rotation periods compiled from S19, S21, and our extended KOI list. (Added: The gray lines in this figure are the seventh-order polynomials described by Bouma et al. (2023) that fit the slow rotation sequences of the Pleiades (Rebull et al. 2016), Praesepe (Rampalli et al. 2021), NGC-6811 (Curtis et al. 2019), NGC-6819 (Meibom et al. 2015), Ruprecht-147 (Curtis et al. 2020), and M67 (Barnes et al. 2016; Dungee et al. 2022; Gruner et al. 2023).)

To assess the statistical uncertainties of (Replaced: these replaced with: our adopted) rotation periods, we compared our (Deleted: adopted) periods with those reported by McQuillan et al. (2014). The details are in Appendix A. We found that for  $P_{\text{rot}} \lesssim 15$  days, the two datasets agree at a precision of  $\lesssim 0.01 P_{\text{rot}}$ . At longer periods of  $P_{\text{rot}} \approx 30$  days, the agreement was typically at the  $\lesssim 0.03 P_{\text{rot}}$  level, and the envelope of the period difference increased roughly linearly with period. Based on this comparison, we adopted a simple prescription for the period uncertainties, such that there are 1% relative uncertainties below  $P_{\text{rot}} = 15$  days, and a linear increase thereafter, with slope set to require 3%  $P_{\text{rot}}$  uncertainties at rotation periods of 30 days.

## 2.2. Kepler Objects of Interest

We considered planets in the NEA cumulative KOI table as of 2023 June 6, which included the best knowledge available on any given planet candidate while also incorporating human-based vetting. These planets represent a superset of those in the fully automated Q1-Q17 DR25 KOI Ta-



ble (Thompson et al. 2018), which could be adopted in future work for planet occurrence rate calculations. This version of the cumulative KOI table included 4,716 objects that are either “confirmed” or “candidate” planets, after excluding known false positives.


### 2.3. High Resolution Spectra

The final piece of our analysis involves assessing ages based on the Li I 6708 Å doublet. We analyzed spectra from the High Resolution Echelle Spectrometer (HIRES; Vogt et al. 1994) on the Keck I 10m telescope. These spectra were primarily collected through the California Kepler Survey (Petrigura et al. 2017; Johnson et al. 2017; Fulton et al. 2017). We supplemented the existing archive with  $\approx 10$  hours of new observations for 22 stars between Fall 2022 and Spring 2024. These stars were chosen to ensure that confirmed planets with rotational evidence for ages below 1 Gyr had spectra, since this is the age range in which lithium is most likely to yield useful age constraints.

Lithium equivalent widths and abundances for the Kepler Objects of Interest were already analyzed by Berger et al. (2018) for roughly three quarters of the spectra in our sample. However, new spectra have since been acquired, and our approach and selection function are different. We therefore performed our own line width measurements on the reduced HIRES spectra.

We collected all blaze-corrected HIRES spectra from our group’s observations of non-false positive Kepler Objects of Interest with a “multiple event statistic” (MES, *koi\_max\_mult\_ev*) of at least 10. This yielded at least one spectrum for 1,464 stars hosting 2,174 planets. About half of these stars have measured rotation periods (797 stars and 1,170 planets, respectively). For stars with multiple spectra available, we analyzed only the spectrum with the highest number of counts. The resulting spectra were acquired between 2009 September 6 and 2024 May 16, and cumulatively comprise 304 hours of open-shutter time.

We measured the lithium equivalent widths using a procedure adapted from previous work (Bouma et al. 2021). Our stars of interest are FGK stars, and so the continuum in the vicinity of the Li I 6708 Å doublet is well-defined. We Doppler-corrected the spectra to a common reference wavelength by cross-correlating against a high S/N template for Kepler-1698, chosen because  $|\gamma| < 10 \text{ km s}^{-1}$ ,  $T_{\text{eff}} \approx 5000 \text{ K}$  and  $v \sin i \approx 5 \text{ km s}^{-1}$ , which puts it in the middle of our sample’s temperature range and gives it mild line-broadening. We then trimmed the Doppler-corrected spectra to a local window centered on the lithium line, using a window width of 15 Å (we also considered 10 Å and 20 Å; the results were consistent). We continuum-normalized by fitting a third-degree Chebyshev series, while excluding regions with absorption lines. We then numerically integrated the resulting spectrum using a one-component Gaussian with free amplitude, width, and mean, and estimated uncertainties on the line width through a Monte Carlo procedure that bootstrapped against the local scatter in the continuum. The resulting EWs are shown in Figure 3.



**Figure 3. Equivalent widths (EW) of the Li I 6708 Å doublet for planet-hosting stars.** These measurements were made from Keck/HIRES spectra collected from 2009–2024. Lines are the “mean” isochrones from Jeffries et al. (2023). The intrinsic dispersion around these isochrones becomes much larger than changes in the mean at  $\gtrsim 1$  Gyr (see Figure 4). Some stars with lithium detections do not have detected rotation periods, and vice-versa.

Our EW measurement approach did not correct for the neighboring Fe I 6707.44 Å blend. To evaluate the accuracy and precision of our method, after applying an initial iteration on the 1,464 stars with spectra, we compared our lithium equivalent widths with those reported by Berger et al. (2018). For the stars in both samples, we found broad agreement at  $> 30 \text{ mÅ}$ , and significant differences at  $< 30 \text{ mÅ}$  because Berger et al. (2018) required positive EWs, while we allowed for statistically negative ones. At  $> 30 \text{ mÅ}$ , there is however a small offset in our respective scales caused by our non-treatment of the iron blend, such that  $(B24-B18) = 7.50 \pm 8.55 \text{ mÅ}$ . We therefore directly subtracted this constant mean value ( $7.50 \text{ mÅ}$ ) when calculating lithium-based ages. This offset is at worst five times smaller than the astrophysical scatter present in lithium EWs for calibration clusters at any given age (see Jeffries et al. 2023).

### 3. STELLAR PROPERTIES

Our default source for stellar temperatures and surface gravities was the Gaia-Kepler Stellar Properties Catalog (GKSP; Berger et al. 2020b). The GKSP parameters were reported for stars with “AAA” 2MASS photometry, measured parallaxes in Gaia DR2, and g-band photometry available from either SDSS or else Kepler-INT (Greiss et al. 2012). The parameters themselves were derived using isoclassify (Huber et al. 2017) to interpolate over the MIST isochrone grids (Choi et al. 2016; Dotter 2016), given the SDSS g and 2MASS  $K_s$  photometry, the Gaia DR2 par-

allaxes, and V-band extinction from the Green et al. (2019) reddening map. The resulting stellar parameters are available for  $\approx 94\%$  of the 53,663 Kepler stars with rotation periods. For the remaining  $\approx 6\%$  of stars that lack temperatures and surface gravities from B20, we adopted the values reported by Santos et al. (2019) and Santos et al. (2021), which are primarily derived from the photometric Mathur et al. (2017) DR25 Kepler Stellar Properties Catalog. In the planet sample,  $\approx 92\%$  of the non-false-positive KOIs with rotation periods have parameters from Berger et al. (2020b), and the remainder are drawn from DR25.

David et al. (2021) compared the photometric B20 stellar parameters ( $T_{\text{eff}}$ ,  $R_*$ , [Fe/H]) against the spectroscopic parameters from Fulton & Petigura (2018). The temperature scales showed a few-percent systematic difference, with F18 quoting higher temperatures than B20 for mid K dwarfs, and lower temperatures for early F dwarfs. Our age analysis, described below, adopts the maximum of the two-sided uncertainties reported by B20 as a symmetric Gaussian temperature uncertainty. Systematic uncertainties in the temperature scale generally influence our age uncertainties at a smaller level than the statistical intrinsic scatter in the open cluster rotation sequences.

#### 4. AGES FROM ROTATION

We calculated gyrochrone ages using `gyro-interp` ((Added: **v0.6**;) Bouma et al. 2023), which is a method designed to address the fact that stars with the same mass and same rotation period can have a wide range of ages (e.g. Gallet & Bouvier 2015). The gyrochrone age posterior should therefore incorporate the intrinsic population-level scatter into its statistical precision. Figure 4 highlights this problem, particularly in regions where the 0.1–1 Gyr stars overlap.

To estimate the stellar spin-down rate as a function of time and stellar temperature, `gyro-interp` uses measured rotation periods and effective temperatures from reference open clusters, and interpolates between them using cubic Hermite polynomials. We calculated the probability of the rotation-based age,  $t_{\text{gyro}}$ , given the observed periods and temperatures (and their uncertainties) by integrating Equation 1 of Bouma et al. (2023). This procedure marginalizes over the astrophysical scatter that is observed in the open cluster sequences. We adopted a linear age grid spanning 0–5 Gyr with 500 grid points, and interpolated the resulting posteriors to calculate any summary statistics. The oldest cluster for which `gyro-interp` is calibrated (Added: **in v0.6**) is the 4 Gyr M67 cluster (Dunee et al. 2022; Gruner et al. 2023). For (Deleted: **Kepler**–) stars with rotation periods above the M67 sequence, our resulting age constraints are (Added: **therefore**) quoted as lower limits.

(Added: **Finally, we opted to modify `gyro-interp` in v0.6 to incorporate an age-dependent population-level Gaussian scatter, as detailed in the change log at [gyro-interp.readthedocs.io](https://github.com/bouma/gyro-interp/blob/main/README.md). While age precisions at  $\leq 1$  Gyr are unaffected, this update accounts for the observed intrinsic scatter in Ruprecht-147 and M67 being**

**larger than in Praesepe or NGC-6811, potentially due to differential rotation (e.g., Epstein & Pinsonneault 2014). For  $\approx 2$ –4 Gyr old stars with rotation periods measured at  $\approx 5\%$  precision, this change yields a  $\lesssim 50\%$  increase in age uncertainties.)**

#### 4.1. Stellar Quality Flags


We calculated gyrochrone ages for all 45,229 stars with reported rotation periods that had effective temperatures of 3800–6200 K. To identify stars for which we suspect these ages may not be valid, we then built a set of quality flags which we condensed into a single binary number:  $Q_{\text{star}}$ . When and how this bitmask should be applied depends on the question being asked. If the goal is to construct a false-positive free sample, all the quality flags could be applied. If the goal is to construct a complete sample, then consider the examples of Kepler 1627Ab ( $\approx 40$  Myr) and Kepler 51c ( $\approx 625$  Myr). The former has a high RUWE due to a resolved binary companion (Bouma et al. 2022a); the latter is on a grazing orbit (Masuda 2014). We leave selection for or against such cases as a decision to the user. For our own analysis, we assume that a star is suitable for gyrochronology if none of bits zero through (Replaced: **nine** replaced with: **ten**) (inclusive) are raised. For analyses that require all stellar rotation periods to come from the same detection pipeline, we further require bit (Replaced: **11** replaced with: **12**) to be zero.

Three assumptions must hold for a rotation-based age dating method like `gyro-interp` to be valid. 1) The evolutionary state of the star must be well-specified by its temperature and its age, 2) the star’s spin-down must not be influenced by binary companions, and 3) the rotation period distribution for field stars of a given temperature and age must be identical to that of equally aged open clusters (metallicity differences, for instance, are ignored). A rephrasing of the first condition is that the star must be “near the main sequence” because during the post main sequence stellar temperatures change. From  $\approx 0.08$  to  $\approx 3$  Gyr, the stars of interest in this work (3800–6200 K,  $0.5$ – $1.2 M_{\odot}$ ) have temperatures constant to  $\approx 1\%$  (Choi et al. 2016). From  $\approx 3$ –4 Gyr, a  $1.2 M_{\odot}$  star’s temperature drops from  $\approx 6200$  K to  $\approx 6000$  K ( $\approx 3\%$ ), because its outer layers expand as it begins hydrogen shell burning as a subgiant. We treat this issue in a manner discussed in “bit 9” below.

*Temperature range (bit 0)*—We require stars to have effective temperatures  $T_{\text{eff}}$  between 3800 and 6200 K in order to report gyrochrone ages (Bouma et al. 2023). Stars hotter than 6200 K spin down very slowly, if at all. Stars cooler than 3800 K do spin down over gigayear timescales (Newton et al. 2016; Engle & Guinan 2023; Chiti et al. 2024), but the currently available open cluster data have yet to clarify when the intrinsic scatter in the population decreases.

*Surface gravity (bit 1)*—We flagged stars as possible subgiants if they had  $\log g < 4.2$ .

*Absolute luminosity (bit 2)*—We calculated the absolute Gaia DR3  $G$ -band luminosity, ignoring reddening, using the reported apparent  $G$ -band magnitude and parallaxes. We



**Figure 4. The models.** Points represent  $10^4$  draws from models that have been fitted to rotation periods (Bouma et al. 2023) and lithium equivalent widths (EWs; Jeffries et al. 2023) of stars in open clusters. Lines are the “mean models” at various ages. The intrinsic dispersion around the mean, which is what the models fit, sets the theoretical precision floor for the age-dating methods. Additional sources of uncertainty, including measurement uncertainty, impose further limits (Deleted: ~~on achievable precision~~). These models are calibrated using data from open clusters. The sizes of the points are the same in each panel, so that low apparent density signifies greater dispersion around the mean.

flagged stars with  $M_G < 3.9$  or  $M_G > 8.5$ , corresponding to main-sequence spectral types earlier than  $\approx F8V$  or later than  $\approx M0.5V$  (Pecaut & Mamajek 2013).

*Known eclipsing binaries (bit 3)*—We flagged any stars reported to be in the final Kepler eclipsing binary catalog (KEBC; Kirk et al. 2016).

*Kepler-Gaia crossmatch quality (bits 4 and 5)*—To leverage Gaia DR3 data, we used M. Bedell’s  $4''$  Kepler-to-Gaia crossmatch of the NEA q1\_q17\_dr25\_stellar catalog with Gaia DR3 (available at <https://gaia-kepler.fun>). The separation distribution of the Kepler-Gaia DR3 crossmatches is such that 99.2% of candidate matches are within  $1''$ . We nonetheless noted an upturn in the candidate match rate from  $3-4''$ ; such sources are flagged using bit 4. For KIC stars with multiple potential Gaia matches within the  $4''$  radius, we adopted the brightest star as the default match. In most such cases this was unambiguous because there is a large brightness difference between the primary and any apparent neighbors. However cases with multiple stars within  $4''$  within  $\Delta G < 0.5$  mag are noted using bit flag 5.

*Gaia DR3 non-single-stars (bit 6)*—The Gaia DR3 non\_single\_star column in the gaia\_source table flags known eclipsing, astrometric, and spectroscopic binaries. We directly included this column.

*RUWE (bit 7)*—We inspected diagrams of the Gaia DR3 renormalized unit weight error (RUWE) as a function of other stellar parameters, and flagged stars with  $RUWE > 1.4$  as possible binaries. Such astrometric outliers can be either bona fide astrometric binaries, or more often are marginally

resolved point sources for which a single-source PSF model provides a poor fit.

*Crowding (bit 8)*—We searched the Gaia DR3 point source catalog for stars within 1 Kepler pixel ( $4''$ ) of every target star. While such companions may not be physically associated with the target star, their presence can confuse rotation period measurements. We therefore flagged any stars with neighbors down to  $1/10^{\text{th}}$  the brightness of the target star within this region ( $\Delta G < 2.5$ ). (Added: We also considered a deeper cut ( $\Delta G < 5$ ), and found it had negligible impact on our conclusions.)

*Near the main sequence (bit 9)*—Figure 1 shows that many stars observed by Kepler are far from the main sequence. Some of the challenges this introduces for rotational ages include unresolved binaries, metallicity, reddening, and drivers of rotation other than magnetic braking. After exploring various options, we settled on the orange locus in the  $\log g - T_{\text{eff}}$  plane shown in Figure 1 as a way of flagging unresolved binaries, as well as evolved late-F and early-G stars. While the exact details of how this locus is constructed are arbitrary (see Appendix B), the general aim is to flag stars for which there is evidence based on their location in the (Replaced: HR replaced with: HR or Kiel) diagram (Added: s) that our gyrochronology model may not be reliable. While the  $\log g - T_{\text{eff}}$  cut flags about half of Kepler stars with rotation periods as potentially not being suitable for gyrochronology, a few outliers in  $M_G$  vs.  $G_{\text{BP}} - G_{\text{RP}}$  remained after applying it. Such outliers are likely also questionable. We therefore also fitted a polynomial to the KOI main sequence in  $M_G$  vs.  $G_{\text{BP}} - G_{\text{RP}}$ ,



and flagged stars more than 1 magnitude from this locus as part of the same bitflag.

**(Added: Metallicity (bit 10))—Models suggest that stars with non-solar metallicities spin down at different rates than solar-metallicity stars (Amard & Matt 2020). This expectation has been hard to verify due to a dearth of nearby non-solar metallicity open clusters. However, if true, this could cause systematic biases for gyrochrone ages (e.g. Figure 9 of Clayton et al. 2020). We therefore flagged stars with spectroscopic  $[\text{Fe}/\text{H}] > 0.3$  from either LAMOST DR5 (Zong et al. 2018) or CKS (Petigura et al. 2022) as being outside the range of our gyrochronology calibration. We discuss limitations of this approach in Section 8.)**

**Candidate pulsators and close-in binaries (bit 10 replaced with: 11))—Santos et al.** included in a flag for candidate “classical pulsators” (e.g. Cepheids) and close-in binaries. Visual inspection shows that although this flag selects many bona fide objects in these classes, it also selects known young planets with  $P_{\text{rot}} \lesssim 5$  day. These objects are neither classical pulsators nor close binaries. While we propagated this flag into our set of quality flags, we therefore generally ignore it.

**Not in the homogeneous stellar rotation sample (bit 10 replaced with: 12))—**This flag was set for the KOIs with rotation periods drawn from a source other than the Santos et al. pipeline.

#### 4.2. Planet Quality Flags

Some planets are more reliably identified than others. We used the following additional quality indicators to assess the reliability and utility of a planet, and assembled them into a separate bitmask,  $Q_{\text{planet}}$ .

**Candidate reliability (bit 0)**—The NEA’s Kepler Objects of Interest Table includes both an overall planet-candidate disposition status (*koi\_disposition*), as well as a disposition based only on the Kepler data (*koi\_pd disposition*). We required both to include only “planet candidates” and “confirmed planets.”

**Candidate S/N (bit 1)**—In any transit survey, the false positive rate increases greatly toward the noise floor for planet detection (e.g. Jenkins et al. 2002). We required a S/N in excess of Kepler’s usual  $7.1\sigma$  floor, through a cut on the maximum “multiple event statistic” (MES, *koi\_max\_mult\_ev*): we required  $\text{MES} > 10$ .

**Grazing planets (bit 2)**—Grazing objects, for which the impact parameter  $b$  is greater than  $1 - R_p/R_*$ , often yield biased planetary parameters (e.g. Gilbert 2022). For large planets, they also include astrophysical false positives at higher rates (Morton et al. 2016), in part due to the size-impact parameter degeneracy. We flagged planet candidates as potentially grazing if  $b > 0.8$ , using the impact parameters reported by Thompson et al. (2018).

#### 5. AGES FROM LITHIUM

Figure 3 shows our measured equivalent widths for the lithium 6708 Å absorption doublet, plotted over the mean isochrone models from EAGLES (Jeffries et al. 2023). We show an upper-limit for plotting purposes if the  $1\sigma$  lower limit on the equivalent width is below 10 mÅ. Our measured EWs span -40 to 250 mÅ; all negative EWs have uncertainties that are statistically consistent with zero.

To calculate lithium ages, we used EAGLES (git commit ac09637), which, similar to *gyro-interp*, is based on an empirical interpolation approach. EAGLES was calibrated on lithium measurements of stars observed by the Gaia-ESO spectroscopic survey in 52 open clusters with ages spanning 2 to 6,000 Myr (Jeffries et al. 2023). We adopted a linear prior in age, spanning  $10^6$  to  $10^{10}$  yr, and symmetrize our EW uncertainties for purposes of interfacing with EAGLES by taking the maximum of the measured positive and negative  $1\sigma$  uncertainty intervals.

In part because many of our EWs are upper limits, many of the lithium ages are lower limits. In such cases, the inferred age posterior is strongly influenced by the assumed intrinsic dispersion in the lithium model, and by the prior. In these instances, we quote the  $3\sigma$  (99.7<sup>th</sup> percentile) limits.


#### 6. CLUSTER AGE COMPARISON

Stars that formed in the same birth cluster are the gold standard for the astronomical age scale (Soderblom 2010). Before Gaia, the few open clusters known in the Kepler field had been cataloged by Herschel (1864). Gaia has enabled the discovery of new stellar ensembles that are more diffuse, but which nonetheless share a common age based on isochrone, rotation, and lithium dating. Specifically in the Kepler field, NGC6811 (1 Gyr; Curtis et al. 2019) and NGC6819 (2.5 Gyr; Meibom et al. 2015) have been known for centuries, while Theia 520 ( $\approx 300$  Myr; Kounkel & Covey 2019), Melange-3 ( $\approx 150$  Myr; Barber et al. 2022), and Cep-Her ( $\approx 40$  Myr; Bouma et al. 2022b and Kerr et al. 2024) are more recent discoveries.

In Figure 5, we compare our rotational ages, and available ages from the literature, against the ages of stars in these open clusters. From the literature, we drew ages from Reinhold & Gizon (2015), Berger et al. (2020b), Lu et al. (2021), Mathur et al. (2023), and Lu et al. (2024). We followed any guidance available from each study for removing ages that were unreliable, and plotted stars within  $1\sigma$  of zero as upper limits. For the Reinhold & Gizon ages, we used those calculated using the Mamajek & Hillenbrand (2008) calibration; for the Mathur et al. ages, we show those from *kiauhoku* (Clayton et al. 2020) rather than STAREVOL (Amard et al. 2019), since the former showed better agreement with the cluster age scale.

Cluster membership is a nuanced subject. Figure 5 is showing reported ages for a set of spatially and kinematically selected stars that could be cluster members, or they could be field interlopers. For NGC6811 and NGC6819, we adopted candidate members from Cantat-Gaudin et al. (2018, 2020) and Kounkel et al. (2020). For Theia 520, we used candidate members from Kounkel et al. (2020). For Melange-3,





**Figure 5. Reported ages of individual stars in benchmark open clusters.** Each point denotes a star’s reported age, normalized by the age of a stellar ensemble in which the star is a candidate member. Cluster membership was evaluated without knowledge of rotation(Added: , and for this plot, no attempt was made to exclude known binaries). Although some field interlopers may be present in the membership lists, outliers can be compared between different methods on a relative basis. Each study was cross-matched against the same cluster list; certain methods report ages for more stars than others. Horizontal scatter is added to visually clarify the statistical age uncertainties. Below 1 Gyr, our ages are generally accurate, and their statistical uncertainties match their dispersion.

we used the candidates reported by Barber et al. (2022) and required “offset” tangential velocities below  $2 \text{ km s}^{-1}$ . For Cep-Her, we used candidate members from Kerr et al. (2024) with  $P_{\text{lin}} > 0.8$ . Even with contaminants, we can compare the relative age distributions derived by the different studies because in all instances we are comparing reported ages against a fixed list of stars.

There are two main metrics for success in this test. 1) Do the reported ages agree with the cluster age? 2) Do the reported age uncertainties agree with their dispersion around the cluster age?

Figure 5 show that although previous studies reported ages that agree with the cluster scale for  $\gtrsim 1$  Gyr stars, sub-gigayear stars have historically had their ages overestimated by 0.3–2 dex, with severely understated uncertainties. For isochrone and kinematic ages, this is because these methods rely on parameters that do not appreciably change at  $t \lesssim 1$  Gyr. For the rotational ages in Mathur et al. (2023) and Lu et al. (2024), the discrepancy is caused by details of the calibration methodologies. The kiauohoku spin-down model for in-


stance has known “stretches” and “compressions” in its age scale relative to observed open clusters (see Mathur et al. 2023, Sec. 7.3). The Mamajek & Hillenbrand (2008) calibration used by Reinhold & Gizon (2015) appears more accurate than the former two models, because it was fitted to reproduce the open cluster sequences known at the time. However, the uncertainties from all of these methods appear to be underestimated. This is probably because they do not marginalize over the range of rotation periods accessible to sub-gigayear stars of a fixed mass and age.

## 7. RESULTS

### 7.1. Rotation-Based Age Distribution

#### 7.1.1. Kepler’s Demographic Cliff

The stellar ages are given in Table 1 for the known planet hosts, and in Table 2 for all Kepler stars. Figure 6 shows the rotation-based age distributions of all the stars (left column) and the KOI hosts (middle column). These “histograms” are sampled from the posterior probability distributions by



**Figure 6. Kepler’s demographic cliff**, visible in the rotation-derived age distributions of its stars (left) and planet hosts (middle). The top row shows (Replaced: all stars with temperatures of 3800–6200 K for which we calculated rotational ages replaced with: stars with temperatures of 3800–6200 K). Opaque lines impose quality cuts on binarity, (Added: metallicity, ) crowding, and the star’s evolutionary state; transparent lines do not (Added: (see Section 4.1)). The bottom row shows stars with temperatures of 4400–5400 K, which (Replaced: have more precise ages due to their fast spin-down replaced with: have more precise ages). The statistical uncertainty for an average star at 1, 2, and 3 Gyr is shown, and is identical across each row. Finally, the right panel compares the (Replaced: rotation-derived replaced with: rotational) ages against star formation histories derived (Replaced: using CMD fitting replaced with: from CMD fits to other regions of the Galaxy) (dashed and dotted lines). Completeness in Kepler’s  $P_{\text{rot}}$  detection sensitivity is near unity until  $t \lesssim 3$  Gyr (Added: , and decreases rapidly at older ages) (Masuda 2022a).

686 drawing ten random samples from each posterior, and then  
 687 computing the normalized histogram of the resulting samples. We also considered an alternative approach for con-  
 688 structing these plots using hierarchical Bayesian deconvolu-  
 689 tion (Masuda et al. 2022), and found similar results. The  
 690 plotted uncertainties (Replaced: assume Poisson statistics  
 691 replaced with: are Poissonian). We truncated the plots at  
 692 4 Gyr, which is the upper bound for our rotational age cali-  
 693 bration. The overall slope (Replaced: yields replaced with:  
 694 shows) a paucity of young stars relative to the expectation of  
 695 a uniform age distribution.

697 The age distributions of Kepler stars and KOI hosts in Fig-  
 698 ure 6 appear similar. One diagnostic for whether the two dis-  
 699 tributions are drawn from the same underlying distribution is  
 700 the Kolmogorov-Smirnov test; we calculated this statistic in  
 701 a manner that accounts for the Poisson uncertainties by per-  
 702 forming 1000 random draws of 70% of the stars from each  
 703 sample when requiring  $t_{\text{gyro}} < 3$  Gyr. The resulting  $\log_{10} p$  val-  
 704 ues spanned (2.5<sup>th</sup> to 97.5<sup>th</sup> percentiles) -4.5 to -1.8 for the  
 705 K dwarfs, and -4.5 to -1.9 for all Kepler stars. This agrees  
 706 with the visual impression that while small differences may  
 707 be present, they are not drastic.

The similarity of the KOI host and Kepler star age distributions adds some nuance to the argument that young transiting planets are hard to detect due to the photometric variability of their host stars. If true, this statement (~~Replaced: seems to be replaced with: might~~) hold only for the very youngest ( $\lesssim 0.4$  Gyr) stars, where there is a marginal deficit in the KOI host age distribution relative to the parent stellar sample. Examining scatter plots of the planet properties as a function of age, we similarly find that fewer systems at  $t_{\text{gyro}} < 0.4$  Gyr are detected with orbital periods beyond 30 days than at  $t_{\text{gyro}} > 0.4$  Gyr.

We can quantify the relative counts of stars as a function of age by labelling stars between 0-1 Gyr, 1-2 Gyr, and 2-3 Gyr as “young”, “intermediate-age”, and “old”. A simple counting exercise from Figure 6 tells us that there are 2.1 times as many old stars in the Kepler field as young stars. Similarly, there are 2.6 times as many old planet hosts as young planet hosts. Focusing only on the tails of the distributions (0-0.3 Gyr and 2.7-3 Gyr), the implication is that the formation rate of stars in the Kepler field decreased by a factor of  $2.81 \pm 0.12$  over the past three billion years. If this decrease continues linearly into the future, then a simple extrapolation would imply that in  $\approx 1.4 \pm 0.3$  Gyr the abundance of newly-formed stars in this region of the Galaxy will reach zero.

In terms of detected planet counts, our rotation-based ages for the Kepler sample yield {109, 201, 281} detected planets in the 0-1 Gyr, 1-2 Gyr, and 2-3 Gyr bins.

#### 7.1.2. Nuances in the Age Distribution

(~~Added: Completeness—Figure 6 has a gray overlay beyond  $\gtrsim 3$  Gyr because Kepler’s sensitivity to rotation signals diminishes at old ages.~~) Masuda (2022a) studied (~~Deleted: this issue~~) (~~Added: Kepler’s completeness to rotation signals using MIST-derived isochrone ages~~), and found for Sun-like stars that the fraction of rotation signals (~~Added: that are~~) detectable (~~Deleted: by Kepler~~) is near unity up to  $\approx 3$  Gyr, and that it drops to almost zero by  $\approx 5$  Gyr. (~~Deleted: This trend is opposite in functional form to our derived age distributions.~~) (~~Added: This estimate suggests that while aspects of Figure 6 might be interpretable in terms of the Galaxy’s star formation history at  $\lesssim 3$  Gyr, at older ages incompleteness precludes any such interpretation.~~)

(~~Added: Temperature sensitivity—There are quantitative differences in Figure 6 between the K dwarfs (4400–5400 K) and the FGK stars (3800–6200 K). For instance, the distribution peaks at 2.3 Gyr for K dwarfs, and at 2.9 Gyr for the FGK sample. The peak is also sharper for the K dwarfs; its decrease from 2.3-3 Gyr is in an age range in which we believe the rotation period catalog completeness to be near unity. We show further details of the  $t_{\text{gyro}}-T_{\text{eff}}$  distributions in Appendix D, and propose possible explanations for these differences in Section 8.2.~~)

## 7.2. Planets Younger Than One Billion Years

(~~Deleted: Sub-gigayear planets can be particularly informative for studies of planet evolution.~~) (~~Added: This analysis provides an important path toward expanding the population of young and evolving exoplanets.~~) Our catalog has 109 confirmed and candidate planets with median ages below 1 Gyr. Requiring  $t_{\text{gyro}} < 1$  Gyr at  $2\sigma$  yields 63 planets orbiting 50 stars. The youth claim is most secure for the systems that are either in clusters (Section 7.2.1), or that have independent rotation and lithium-based ages (Section 7.2.2). These samples provide broader context for the planets around field stars that push the boundaries of the current young exoplanet census (Section 7.2.3).

### 7.2.1. Planets in Clusters

Four stellar ensembles in the Kepler field have to date yielded (~~Deleted: a total of~~) fourteen transiting planets. Our analysis blindly recovers the youth of all of these planets.


*Cep-Her*—The Cep-Her complex ( $\approx 40$  Myr) contains Kepler-1627Ab, Kepler-1643b, Kepler-1974b and Kepler-1975 Ab (Bouma et al. 2022a,b). Different sub-groups of the complex vary in age by  $\approx 50\%$  (Kerr et al. 2024). The planets themselves are all on close-in orbits (5-25 days), with sizes from  $2-4 R_{\oplus}$ . Our rotation and lithium ages agree with the cluster age for Kepler-1627A, Kepler-1974, and Kepler-1975A. For Kepler-1643, the rotation and cluster ages agree, and the lithium age is  $2.0\sigma$  above the cluster age (~~Replaced: reported by~~ replaced with: from) Bouma et al. (2022b) for RSG-5.

*MELANGE-3*—Barber et al. (2022) reported a  $105 \pm 10$  Myr association in the Kepler field containing two transiting mini-Neptunes: Kepler-970 and Kepler-1928. We find rotational ages of  $t_{\text{gyro}} = 176^{+120}_{-40}$  Myr and  $144^{+104}_{-88}$  Myr for Kepler-970 and Kepler-1928 respectively. The former (~~Deleted: rotation-based~~) age being slightly older than the suggested Pleiades age for the association agrees with Figure 4 of Barber et al. (2022).

*Theia 520*—We find rotational ages of  $\approx 350$  Myr for Kepler-968 and Kepler-52. Based on spatial and kinematic clustering, these stars are candidate members of Theia 520 (Kounkel & Covey 2019). The core of this cluster is also known as UBC-1 (Castro-Ginard et al. 2018). Based on isochrone and rotational age-dating, Theia 520 seems to be  $\approx 230 \pm 70$  Myr old (Fritzewski et al. 2024), which is within  $1\sigma$  of our  $t_{\text{gyro}}$  measurements. Ongoing work by Curtis (2024) broadly finds that the two planet-hosting stars are indeed members of this diffuse population. Given the cluster-level and individual-star evidence, this makes Kepler-968 and Kepler-52 the youngest multiplanet systems currently known from the main Kepler mission.

*NGC 6811*—Meibom et al. (2013) reported the discovery of Kepler-66 and -67b, two mini-Neptunes in NGC 6811 ( $\approx 1$  Gyr). We derive rotational ages for these systems of  $1431^{+675}_{-364}$  Myr and  $878^{+108}_{-121}$  Myr for Kepler-66 and -67b, respectively. Kepler-66 ( $T_{\text{eff}} \approx 5900$  K,  $P_{\text{rot}} \approx 10.4$  days) has





**Figure 7.** Sizes, orbital periods, and rotation-based ages of transiting exoplanets younger than one billion years. (Replaced: Star symbols denote 46 planets that are characterized in this work to have  $t_{\text{gyro}} < 1$  Gyr at  $2\sigma$ ; circles are planets from the literature meeting the same age requirement, but with a heterogeneous set of age provenances. Gray points are transiting planets from the NASA Exoplanet Archive older than 1 Gyr. If one were to instead select for precise ages ( $t/\sigma_t > 3$ ), this would hide  $\approx 10$   $< 0.5$  Gyr planets, and add  $\approx 15$   $0.5$ – $1$  Gyr planets. For this plot, we require our Kepler stellar hosts to not have  $Q_{\text{star}}$  bits 0–9 raised, and for the planets to similarly no have quality flags raised. replaced with: Systems are selected to have  $t < 1$  Gyr at  $2\sigma$ . Star symbols denote 46 planets characterized in this work to have such ages at the highest confidence, meaning that none of  $Q_{\text{star}}$  bits 0–10 are raised, and the planets have no quality flags raised. Transparent crosses denote 17 planets for which either the planet is grazing, or the star has  $\text{RUWE} > 1.4$ . Gray points are from the NASA Exoplanet Archive. )

a lower precision and a more asymmetric posterior because of its slow spin-down rate. Kepler-67 ( $T_{\text{eff}} \approx 5100$  K,  $P_{\text{rot}} \approx 10.3$  days) is marginally hotter than the later K dwarfs that are in “stalled” spin-down at this time, enabling its rotation period to be diagnostic of its age.

### 7.2.2. Rotation vs. Lithium Ages

We compared the rotation and lithium ages for the known planet hosts in the “Consistent?” column of Table 1. There were three cases of interest. *i*) If two-sided posteriors for both  $t_{\text{gyro}}$  and  $t_{\text{Li}}$  existed, and their median values were consistent within  $2\sigma$ , we listed them as consistent; if they were consistent within  $2$ – $3\sigma$ , we listed them as “maybe” consistent. Otherwise, they disagreed. *ii*) If  $t_{\text{Li}}$  was a lower limit, we compared this lower limit with the  $1\sigma$  upper limit from  $t_{\text{gyro}}$ ; if the two overlapped, we judged the age estimates to be consistent. *iii*) If  $t_{\text{Li}}$  provided a two-sided posterior, and no rotation period was found, we judged the age estimates to be inconsistent. This is because a two-sided lithium constraint can only be provided for G and K dwarfs  $\lesssim 2$  Gyr old,

and Kepler should have been sensitive to the rotation periods of such stars.

In the entire planet sample (i.e., without imposing any quality cuts) this yielded 697 consistent cases, 12 maybe consistent cases, and 22 inconsistent cases. Rephrased, in the overall sample, 95% of stars have consistent rotation and lithium-based ages (97% potentially consistent).

Appendix C describes all cases of “discrepant” confirmed planets for which a sub-gigayear age is reported by at least one age indicator. All but two systems are either evolved stars or else unresolved binaries, and are automatically flagged as such. The more interesting of the two is Kepler-786, an early K dwarf with  $t_{\text{Li}} = 228^{+168}_{-87}$  Myr, but with a  $\approx 33$  day rotation period that implies  $t_{\text{gyro}} \approx 4.4$  Gyr. Other age indicators in the spectrum similarly suggest that the star is not young. It therefore appears to be anomalously lithium-rich.

### 7.2.3. Notable New Young Planets


Let us focus on blemish-free stars with blemish-free planets:  $Q_{\text{star}}$  must not have any of bits zero through (Replaced: **nine** replaced with: **ten**) raised, and  $Q_{\text{planet}}$  must similarly have no quality flags raised. Under this constraint, our results include 18 confirmed planets younger than 1 Gyr with two-sided  $t_{\text{gyro}}$  and  $t_{\text{Li}}$ , and 37 confirmed planets with comparable ages for which  $t_{\text{gyro}}$  is two-sided while  $t_{\text{Li}}$  is one-sided.

Figure 7 shows the planets and planet candidates with  $t_{\text{gyro}} < 1$  Gyr at  $2\sigma$ . (Replaced: This replaced with: The stars in this figure include (Deleted: s) 37 confirmed Kepler planets, and nine candidate planets, all (Deleted: of which are) listed in Table 1. These include four objects Earth-sized or smaller, six Jovian-sized planet candidates, seven super-Neptunes ( $4$ – $10 R_{\oplus}$ ), 28 mini-Neptunes above the radius valley as parametrized by Van Eylen et al. 2018, and 18 super-Earth sized planets below the radius valley. While most of these planets have orbital periods below 50 days, four are on more distant orbits.

Among these systems, four highlights are Kepler-1529, Kepler-1565, Kepler-1312, and Kepler-1629. Kepler-1529 b is a  $\approx 100 \pm 50$  Myr mini-Neptune with well-measured rotation and lithium ages. Kepler-1565 b is an analogous  $\approx 170$ – $230$  Myr super-Earth. Kepler-1312 is a  $t_{\text{gyro}} = 357^{+75}_{-109}$  Myr near-Solar analog with an Earth-sized planet on a one-day orbit, and a mini-Neptune on a five-day orbit. The Earth-sized planet, Kepler-1312 c, along with the near-identical Kepler-1561 b ( $t_{\text{gyro}} = 426^{+74}_{-78}$  Myr), (Replaced: rank among replaced with: **could be**) the youngest Earth-sized planets currently known, (Added: and are) comparable to planets such as HD 63433 d ( $1.1 R_{\oplus}$ ,  $414 \pm 23$  Myr; Capistrant et al. 2024) and TOI-1807 b ( $1.3 R_{\oplus}$ ,  $180 \pm 40$  Myr; Hedges et al. 2021). Kepler-1629 b, with a size just two-thirds that of Earth ( $0.67 \pm 0.05 R_{\oplus}$ ), is marginally older, also based on rotation ( $529^{+62}_{-62}$  Myr). The lithium age from a reconnaissance TRES spectrum confirms the point.

## 8. DISCUSSION

### 8.1. Kinematic Heating and Star Formation in the Thin Disk



**Figure 8. Impact of star formation rate (SFR) and kinematic heating on age distribution.** Most Kepler stars are a few hundred parsecs above the Galactic plane ( $\hat{Z}$  points to the north galactic pole;  $\hat{Y}$  toward the direction of galactic rotation). This is comparable to the  $\approx 300$  pc mass-weighted scale height. The dotted gray and solid orange lines show synthetic age distributions from TRILEGAL assuming a constant SFR (middle) and a two-step SFR (right), described in the text. A combination of both kinematic heating and a decreasing SFR are needed to qualitatively match the observed paucity of the youngest stars (blue line; as in Figure 6).

(Added: Two separate astrophysical effects might explain our observed age distribution (Figure 6): a declining star formation rate (SFR), and kinematic heating. Kinematic heating is associated with an increase in both stellar velocity dispersions and the disk's vertical scale height over time (e.g. Villumsen 1983; Schmidt et al. 2024). The Kepler field points toward  $(l, b) = (76.3^\circ, 13.5^\circ)$ . A typical Kepler star at 1 kpc is therefore  $\approx 230$  pc above the galactic plane, a significant fraction of the thin disk's mass-weighted  $\approx 300$  pc scale height (Bland-Hawthorn & Gerhard 2016). )

(Added: We assessed the relative importance of these two effects using the TRILEGAL Monte Carlo population synthesis tool (v1.6; Girardi et al. 2005). (Added: TRILEGAL treats the Galaxy as a linear combination of a thin disk, thick disk, halo, and bulge, and allows for two possible SFRs: constant, and a "two-step" SFR in which the thin disk SFR is 50% larger from 1-4 Gyr than at other ages. This two-step SFR was introduced to improve the model's ability to match star counts from Hipparcos and 2MASS (see Section 5.1 of Girardi et al. 2005). (Added: For kinematic heating, TRILEGAL assumes that the disk scale height grows as  $h_z = z_0(1 + t/t_0)^\alpha$ , for  $\alpha = 5/3$ ,  $z_0 = 94.7$  pc, and  $t_0 = 5.55$  Gyr, similar to Rana (1991). We queried the web interface toward the Kepler field, and toward the galactic plane (at  $l = 76.3^\circ$ ) to negate the effects of kinematic heating. We limited the output to stars with apparent Kepler magnitudes brighter than 16, and set the field of view to  $10 \text{ deg}^2$ , the maximum available through the web interface. )

(Added: Figure 8 compares the results against our observed age distribution. The middle panel shows that assuming a constant SFR, kinematic heating alone can yield

a two-fold difference in the abundance of the youngest stars relative to those  $\approx 3$  Gyr old. The right-most panel shows that additionally including the time-variable SFR yields the best agreement with the slope of our observed age distribution. )

(Deleted: Most stars observed by Kepler are in the thin disk. This can be verified by following Gaia Collaboration et al. (2018), and labeling stars with 2D tangential velocities  $v_T < 40 \text{ km s}^{-1}$  as thin disk members, and those with  $60 < v_T < 150 \text{ km s}^{-1}$  as thick disk members. These criteria yield 95,694 thin and 54,039 thick disk stars observed by Kepler. The thin disk stars have a threefold larger detected rotation period fraction: 35,675 rotators are in the thin disk, while 7,312 are in the thick disk. Further imposing the quality flags discussed in Section 4.1 yields 17,755 thin disk stars for which gyrochronology is applicable, and 2,462 thick disk stars. )

(Deleted: Classifying Kepler stars as thin vs. thick disk members helps connect them to previous work on the Galaxy's star formation history. On extragalactic scales, the star formation rate in spiral galaxies with mass similar to the Milky Way peaked  $\approx 10$  Gyr ago, and has since decreased by an order of magnitude. ) (Added: Despite the expected differences between pencil-beam surveys and volume-complete samples, our observed age distribution qualitatively agrees with a number of previous studies.) (Replaced: In our own galaxy, previous studies have replaced with: Previous work has) measured star formation histories through CMD fitting of resolved stellar populations (Mor et al. 2019; Ruiz-Lara et al. 2020; Alzate et al. 2021; Xiang & Rix 2022), and by modeling the local white dwarf luminosity function (e.g. Isern 2019). Mor et al. (2019) (Replaced: and Isern (2019) for instance focused on stars

(and white dwarfs) replaced with: and [Mazzi et al. \(2024\)](#) for instance focused on stars) within a few hundred parsecs, and both found a peak in the local star formation rate 2-3 Gyr ago. [Ruiz-Lara et al. \(2020\)](#) focused on stars in a wider 2 kpc bubble, and additionally reported three local maxima in the SFR, which they associated with pericenter passages of the Sagittarius satellite galaxy.

The right column of Figure 6 compares our derived age distribution against star formation histories (SFHs) reported by [Mor et al. \(2019\)](#) and [Ruiz-Lara et al. \(2020\)](#) based on CMD fitting. The overall slope of both SFHs broadly agrees with what we find from rotation-based ages. The SFH from [Mor et al. \(2019\)](#) seems entirely consistent, particularly after accounting for the statistical uncertainties of that study. Regarding the episodic star formation bursts reported by [Ruiz-Lara et al. \(2020\)](#), they are not apparent in our overall FGK star sample (top row). However, rotational ages are most precise for  $\approx$ G8V-K4V dwarfs older than  $\approx$ 1.5 Gyr ([Bouma et al. 2023](#)). The lower row of Figure 6 selects these stars with a temperature cut, and may show a (Added: weak) hint of the (Added:  $\approx$ 1 Gyr and  $\approx$ 2 Gyr spike(Added: s) reported by [Ruiz-Lara et al. \(2020\)](#). (Deleted: The  $\approx$ 1 Gyr spike however is not recovered.— ) This  $\approx$ 2.5 Gyr local maximum is similarly present in the isochrone ages derived by [Berger et al. \(2020b\)](#) for the Kepler field, and in the red giant asteroseismic ages derived by [Silva Aguirre et al. \(2018\)](#). The main novelty of our ages relative to (Replaced: those studies is their improved accuracy at  $<$  replaced with: these previous studies is their accuracy below) 1 Gyr(Added: (see Figure 5)).

## 8.2. Caveats & Limitations

Our most constraining ages tend to come from one source of information: rotation. Factors other than (Deleted: stellar) age and mass can influence (Deleted: stellar-) rotation rates. (Added: Astrophysical factors include metallicity, latitudinal differential rotation, mergers, and tidal spin-up. Observational limitations include the difficulty of measuring small-amplitude, long-period signals that are only quasi-coherent due to the starspot lifetime approaching the stellar rotation period. There is also the issue that our age-dating approach is not calibrated beyond 4 Gyr. In the following, we attempt to quantify the importance of these effects in our analysis, ordered by what we perceive to be their descending importance.)

*Binaries:* Stars in binaries are often biased toward rapid rotation, even beyond the typical realm of tidal synchronization (see [Meibom et al. 2007](#); [Gruner et al. 2023](#)). (Added: [Simonian et al. \(2019\)](#) for instance found that 59% of Kepler stars with rotation periods between 1.5 and 7 days lie 0.3 mag above the main sequence, compared with 28% of their full rotation sample. We attempted to mitigate the effects of binarity with five distinct quality flags (Section 4.1). This attempt was at least somewhat effective, based on the peak near zero in the raw distributions of Figure 6 being flattened in the cleaned distributions. However, the completeness of our quality flags is hard to


assess. One specific concern is that these flags might have minimal sensitivity to binaries with separations of  $\approx$ 0.05-0.5 AU, for all but the highest mass ratios. This inner limit is set by the onset of detectable ellipsoidal variations in [Kirk et al. \(2016\)](#); the outer limit is set by the typical orbital periods of astrometric binaries in Gaia DR3 (see Figure 3 of [Gaia Collaboration et al. 2023](#)).

(Added: If we were not at all sensitive to binaries with separations of 0.05-0.5 AU, then based on the binary fraction and separation distribution from [Raghavan et al. \(2010\)](#), as much as  $\approx$ 5% of our “gyro applicable” sample (1,190/23,813 stars) might be contaminated. In a worst case scenario, all of these unidentified binaries might induce  $P_{\text{rot}} < 10$  days, which would then be mixed in with the bona fide young, single stars. This would constitute a major source of contamination, given that our nominal sample of apparently clean stars had 2,706 stars with  $P_{\text{rot}} < 10$  days. )

(Added: Figure 9 shows the rotation-based age distribution calculated by randomly dropping 1,190 of the  $P_{\text{rot}} < 10$  day stars. Relative to the nominal age distribution, there are three important differences. There is a much larger drop in the star formation rate over the past three gigayears, by a factor of  $4.81 \pm 0.26$ , rather than  $2.81 \pm 0.12$ . The rate of decline in the age distribution is also constant, rather than having a “kink” at 1.5 Gyr. And finally, the corrected distributions show closer agreement with both the SFHs derived from CMD fitting, and also with the KOI host distribution, which is almost devoid of close binaries within 1 AU (e.g. [Moe & Kratter 2021](#), and references therein). We caution however that this correction assumes that none of our quality flags were able to identify any of these unresolved binaries. Since we did in fact include flags based on e.g., location in the HR diagram, this correction is best interpreted as an upper limit on the possible impact of binarity.)

(Added: Rotational outliers: ) We were interested in not only the overall age distribution of the Kepler field, but also in the reliability of individual ages for (Replaced: young stars known to host planets replaced with: each star). We attempted to assess this reliability using quality bitmasks (Table 1), a comparison against (Deleted: open-) cluster ages (Section 6), and a comparison between rotation and lithium ages. The (Deleted: open-) cluster comparison suggested that our ages were mostly accurate at  $\ll 0.5$  dex across 0.04–2.5 Gyr, similar to their quoted precision. Nonetheless, the upper-left panel of Figure 5 does show outliers, typically  $\lesssim 10\%$  of the population in any cluster. (Added: While most of these outliers have known bad quality flags, some do not.) These stars could be either field star contaminants, or else anomalous stars whose physical rotation histories were altered by processes not captured by our statistical uncertainties. Similarly, although we found consistent  $t_{\text{gyro}}$  and  $t_{\text{Li}}$  for 18 planets orbiting “high-quality” stars,





**Figure 9. Potential impact of unresolved binaries.** This figure shows the same raw (thinnest line) and quality-flag cleaned (“gyro applicable”) distributions as in Figure 6. However, even after applying our quality cuts, up to 1,190 of 2,706 apparently clean rotators with  $P_{\text{rot}} < 10$  days could be unresolved  $\lesssim 0.5$  AU binaries (see Section 8.2). The dark blue “5% binary contamination” distribution corrects for this possibility by randomly removing what we estimate (estm.) to be the maximum possible number of such systems; the true correction might be smaller. We expect such a correction to be more important in the sample of all Kepler stars than in the KOI host sample.

we did find one system, Kepler-786, with radically different lithium ( $228^{+168}_{-87}$  Myr) and rotational ( $\approx 4.4$  Gyr) ages. Planetary mergers are one process that could produce such a signal, but testing this would require a method for measuring differential abundances across a large number of elements.

(Added: A separate plausible origin for rotational outliers is that they come from erroneously reported rotation periods. While the large surveys do show good internal consistency (Appendix E), the referee correctly pointed out that this is only a comparative test of methods, rather than an ability to recover ground truth (e.g. Aigrain et al. 2015). One could imagine an adversarial scenario in which, say, 10-20% of stars with  $P_{\text{rot}} > 20$  days are erroneously reported, and in fact have much longer rotation periods. In this scenario, a similar analysis to that in Figure 9 showed that our analysis would overestimate the degree by which the SFR in the Kepler field has decreased, by 7-14%.)

(Added: Weakened magnetic braking: If stellar rotation rates and spot-induced photometric amplitudes remained constant after  $\approx 3$  Gyr, then older stars could contaminate the peaks in Figure 6. This possibility warrants consideration, given asteroseismic evidence suggesting slowed spin-down rates after  $\approx 4$  Gyr for Sun-like stars (van Saders et al. 2016; Hall et al. 2021; Saunders et al. 2024), and the “long-period edge” in the  $P_{\text{rot}}-T_{\text{eff}}$  distribution which may pile up over similar timescales (David et al. 2022). However, this stalling is thought to happen at  $> 5$  Gyr for late-G dwarfs and early K dwarfs (e.g. Figure 4 by Saunders et al. 2024), which means that

it cannot account for the 2.3 Gyr peak in the 4400-5400 K age distribution shown in Figure 6.)

(Added: Additionally, Masuda (2022a) found that the McQuillan et al. (2014) sample, discussed in Appendix E, includes few stars older than  $\approx 4$  Gyr, based on isochrone-fitting to a high-quality subset of Kepler planet hosts. Some solar twins have also been shown to have rotation periods of 30-50 days (Lorenzo-Oliveira et al. 2019, 2020; do Nascimento et al. 2020), indicating that a completely stalled spin-down may be unlikely. These arguments suggest that while the peak in the age distribution is not primarily caused by an accumulation of much older stars with slowed spin-down rates, contamination by such stars beyond  $\approx 3$  Gyr remains possible, particularly for stars hotter than the Sun.)

(Added: Metallicity: Stars with higher metallicity are predicted to have longer convective turnover times, which may slow their rate of spin-down (Amard & Matt 2020). While recent observational work has empirically confirmed that such an effect is present in the Kepler data (See et al. 2024), the anti-correlation between close, unresolved binaries and stellar metallicity (Moe et al. 2019) could be a complicating factor. Nonetheless, based on these studies, we flagged stars with known spectroscopic  $[\text{Fe}/\text{H}] > 0.3$ , using either LAMOST or Keck/HIRES spectra (Zong et al. 2018; Petigura et al. 2022). The main limitation of this approach is that only 53.3% of the stars in Table 2 with  $t_{\text{gyro}} < 4$  Gyr had such spectra available. For the remainder of our sample, our quality flags were not sensitive to metallicity outliers. The metallicity distribution of stars for which we do have spectra suggests

that at most  $\approx 5\%$  of stars in our apparently clean sample might have  $|\text{[Fe/H]}| > 0.3$  and not be flagged, with a slight bias toward lower metallicity stars over higher metallicity. Based on work by Claytor et al. (2020), this will be most important for stars near the Kraft break, which could have ages over-estimated by  $\approx 50\text{--}100\%$  if they are in fact metal-poor, and under-estimated in the converse case.)

(Added: *Differential rotation*: The Sun’s surface rotates once per  $\approx 25\text{--}30$  days at latitudes between the equator and  $\approx \pm 60^\circ$  (e.g. Snodgrass & Ulrich 1990). This latitudinal surface differential rotation yields two separate limitations on rotational ages. The first is on the accuracy with which any one star’s rotation period can be measured over a few-year mission like Kepler; the second is on the degree to which the intrinsic scatter in open cluster rotation sequences can be measured (e.g. Epstein & Pinsonneault 2014). In this study, we accounted for the latter intrinsic scatter based on available measurements from Ruprecht-147 (Curtis et al. 2020) and M67 data (Barnes et al. 2016; Dungee et al. 2022; Gruner et al. 2023). This means that irrespective of the assumed  $P_{\text{rot}}$  measurement precision, our model imposed a  $\approx 5\text{--}10\%$  level of intrinsic  $P_{\text{rot}}$  scatter at old ages. While this accounts for the statistical impact of differential rotation on the inferred age distribution, the accuracy issue remains; any one individual star’s observed rotation period might be faster or slower than average stars of the same mass and age. This effect can yield a  $\approx 20\%$  age error for individual Sun-like stars with ages of a few billion years; the statistical uncertainties in Table 2 should be treated with appropriate caution.)

(Deleted: Metallicity may also be relevant (Amard et al. 2020; See et al. 2024). An additional caveat is that our rotation-based ages used photometric effective temperatures (Section 3), even though spectroscopic temperatures are available for the planet-hosting subset. This decision was driven by a desire for homogeneity irrespective of planet-hosting status. However it implies that the estimated planet ages might shift if one were to account for the spectroscopic information.)

(Deleted: Other age indicators may help in verifying the ages of the  $\lesssim 3$  Gyr stars that are the main focus of this work. Specifically, chromospheric emission in the X-ray, Ca II HK line, and the UV can serve as an age tracer (Mamajek & Hillenbrand 2008; Vidotto et al. 2014; Engle 2024). However, these age indicators are all in a sense “rotation-powered”. The dynamo converts kinetic energy into magnetic energy, which is emitted through these chromospheric pathways. We did not attempt to incorporate these indicators into this study due to concerns regarding sensitivity, homogeneity, and the question of whether they in fact provide age information that is truly independent from rotation.)

### 8.3. Future Directions

(Added: *Ages of stars older than the Sun*: ) (Deleted: A broader question, beyond the scope of this work, concerns the ages of stars  $\gtrsim 4$  Gyr old. ) (Added: This work was limited to stars younger than 4 Gyr). For age-dating methods based only on rotation rates, an important challenge is that the detectability of the rotation signals for 3–10 Gyr stars, especially Sun-like stars, is at the limits of the Kepler data (Masuda 2022a). In the “old star” regime, methods that leverage either asteroseismology (van Saders et al. 2016; Saunderson et al. 2024), kinematics (Lu et al. 2021; See et al. 2024), or else that combine both the evolution of stellar luminosity and stellar rotation (Angus et al. 2019; Claytor et al. 2020; Mathur et al. 2023) seem the most capable of providing useful age constraints for single stars over the full span of their main-sequence lifetimes.

(Added: *Planet occurrence rates*: The age-dependent trends predicted for exoplanet populations, such as the Kelvin-Helmholtz cooling of mini-Neptunes (Gupta & Schlichting 2019), time-dependent carving of the photoevaporation desert (Owen & Lai 2018), and the time evolution of the radius valley (Rogers & Owen 2021) can be explored using our data. We defer this analysis to a separate publication. Some care is required since beyond age, exoplanet demographics also depend on stellar metallicity and mass (e.g. Petigura et al. 2018; Miyazaki & Masuda 2023).

*Field star ages from other surveys*: While this study focused on Kepler, other photometric surveys (e.g. K2, TESS, HAT, WASP, NGTS, ZTF, ATLAS) open opportunities for age-dating a far broader set of stars and planets. Future prospects also include PLATO (Rauer et al. 2014), Earth 2.0 (Ge et al. 2022), and the Roman Galactic Bulge Time Domain Survey (Wilson et al. 2023). Rotational ages from these surveys could yield new insight into (Replaced: whether the star formation history that we see in the Kepler field is universal across the thin disk. replaced with: the relative importance of star formation and kinematic heating across different components of the Galaxy.)

*Searches for new associations*: Some of the youngest stars in Tables 1 and 2 have been linked to their birth clusters. Others have not. A systematic search for the birth clusters of the youngest “field stars” could combine positions and velocities from Gaia with rotation measurements from TESS. Given the  $\sim 100$  Myr decoherence time of young clusters, (Replaced: the youngest “field stars” replaced with: these young stars) may have easily identifiable young neighbors.

## 9. CONCLUSIONS

We began (Deleted: this work) with two questions: how wrong is the assumption of a uniform age distribution for stars in the galactic thin disk? And why are only  $\approx 50$  sub-gigayear transiting planets known, rather than the  $\approx 500$  that would be expected under the assumption of a uniform star formation (Replaced: history replaced with: rate) for the  $\approx 5,000$  known planets?

Our approach to answering these questions was to curate a sample of stellar rotation periods, lithium equivalent widths, and temperatures using archival and new data from the Kepler field. We derived new ages using empirical interpolation-based methods, and assessed the reliability of these ages by comparing them against benchmark open clusters (Figure 5).

Tables 1 and 2 summarize the results for planets and stars, respectively. Our recovered ages are accurate for all 14 known Kepler planets in clusters. While lithium provided minimal added information for most of the sample (Added: due to its astrophysical depletion rate),  $t_{\text{Li}}$  and  $t_{\text{gyro}}$  agreed in over 90% of cases for which comparison was possible. Our (Replaced: resulting ages replaced with: results) included two-sided  $t_{\text{gyro}}$  and  $t_{\text{Li}}$  for 18 sub-gigayear confirmed planets, and two-sided  $t_{\text{gyro}}$  with one-sided  $t_{\text{Li}}$  for 37 (Replaced: sub-gigayear confirmed replaced with: such) planets. Allowing for “candidate” planets, grazing transit geometries, and stars with RUWE far from unity expands the counts by a factor of two.

The sizes and periods for the most secure set of planets are shown in Figure 7. While the new young planets are mostly mini-Neptunes, some are near the lower boundary of the “sub-Jovian desert” (Owen & Lai 2018), which could have an evolutionary connection to their youth. Other discoveries include Earth-sized planets (Added: (e.g. Kepler-1312 c and Kepler-1561 b) with (Deleted: new-) ages of only a few hundred million years.

(Added: Significant care is needed not only when assessing the reliability of individual rotational ages, but also when interpreting distributions of rotational ages inferred from many stars. Despite a slew of stellar quality flags, including on metallicity, evolutionary state, and binarity, up to a third of our “apparently clean” Kepler stars with  $P_{\text{rot}} < 10$  days could potentially be influenced by unresolved binaries (Figure 9). An additional nuance is that even after correcting for such effects, the star formation history does not directly trace the observed age distribution: particularly for pencil-beam surveys such as Kepler, kinematic heating of stars in the thin disk significantly sculpts the observed stellar (and therefore planetary) age distribution.)

Our (Deleted: main-) conclusions with respect to our original (Added: two) questions (Added: – what is the age distribution in the thin disk, and where are the missing young transiting exoplanets –) are as follows.

1. Rather than being uniform, the age distributions of both the Kepler target stars and the known Kepler planet hosts show a demographic cliff. There are (Added: at least) twice as many “old” (2-3 Gyr) stars in the Kepler field as “young” (0-1 Gyr) stars. (Deleted: The star formation rate today is  $2.81 \pm 0.12$  times lower than it was three billion years ago. )

2. (Added: The youngest Kepler stars (0-0.3 Gyr) are at least  $2.81 \pm 0.12$  times rarer than stars 2.7-3 Gyr old. The statistical uncertainty on this value is dwarfed by the systematic uncertainty in the contamination rate of unresolved photometric binaries, which could push it as high as  $4.81 \pm 0.26$  (Section 8.2)). (Replaced: This result replaced with: This range of values) (Replaced: from rotation-based ages broadly agrees replaced with: is broadly consistent) with (Replaced: recent reports of a declining star-formation rate replaced with: modeling expectations (Section 8.1), and with age distributions derived from) CMD fitting and white-dwarf chronology (Replaced: , though with the replaced with: . The ) advantage of (Added: rotation-based ages however is their ) (Replaced: sensitivity replaced with: accuracy) for individual FGK stars at  $t < 1$  Gyr (e.g. Figure 5).
3. Rather than expecting  $\approx 500$  cumulative exoplanets younger than one billion years, the age distribution of the Kepler field (Replaced: implies replaced with: suggests) that we (Replaced: should replaced with: might) instead expect  $\approx 250$ .
4. We have (Replaced: derived replaced with: reported) rotational ages for 63 Kepler planets younger than 1 Gyr at  $2\sigma$ , and 109 planets with median ages below 1 Gyr. Concatenating these planets against the existing literature yields  $\approx 170$  known sub-gigayear planets. This lessens the original factor of ten discrepancy to a factor of at most two.

## ACKNOWLEDGMENTS

We thank Kevin Schlaufman and the reviewer for suggestions which significantly improved the manuscript. This work was supported by the Heising-Simons 51 Pegasi b Fellowship (LGB, EKP) and the Arthur R. Adams SURF Fellowship (EKP). The HIRES data were obtained at the Keck Observatory. We recognize the importance that the summit of Maunakea has always had within the indigenous Hawaiian community, and we are deeply grateful for the opportunity to conduct observations from this mountain.

*Contributions:* Per <https://credit.niso.org/>: Conceptualization: LGB. Data curation: LGB, AWH, HI. Formal analysis: LGB, KM. Funding acquisition: LGB. Investigation: LGB, EKP. Methodology: LGB, EKP, KM. Project administration: LGB, LAH, AWH. Resources: LGB, AWH. Software: LGB. Supervision: LGB, LAH. Validation: LGB, KM. Visualization: LGB. Writing – original draft: LGB. Writing – review & editing: all authors.

*Facilities:* Gaia (Gaia Collaboration et al. 2023), Kepler (Borucki et al. 2010), Keck:I (HIRES) (Vogt et al. 1994), 2MASS (Skrutskie et al. 2006), SDSS (York et al. 2000).

*Software:* astropy (Astropy Collaboration et al. 2013, 2018, 2022), claude (Anthropic 2024), eagles (Jeffries et al. 2023), gyro-interp (Bouma et al. 2023), matplotlib (Hunter



1329 et al. 2007), numpy (Van Der Walt et al. 2011), scipy (Virta-  
1330 nen et al. 2020).

**Table 1.** Ages of Kepler planets and planet candidates. This version of the table is truncated to include the youngest systems, sorted by the minimum of either the rotation or lithium-based age. The full machine-readable table contains ages and age limits for 2,461 non-false positive KOIs with  $MES > 10$ . A bash script to decode the  $Q_{\text{star}}$  quality flag is [available online](#). A python script to select stars with specific bit flags is [also available](#). All quoted age uncertainties are statistical.

KOI	Kepler	$T_{\text{eff}}$	$P_{\text{rot}}$	$EW_{\text{Li}}^*$	$t_{\text{gyro}}$	$t_{\text{Li}}$	Consistent?	$R_p$	$P_{\text{orb}}$	$Q_{\text{planet}}$	$Q_{\text{star}}$	Spec?	Comment
–	–	K	days	mÅ	Myr	Myr	str	Earths	days	int	int	bool	–
K05245.01	Kepler-1627 b	5357	2.62	$225 \pm 7$	$81^{+158}_{-55}$	$51^{+38}_{-27}$	Yes	3.79	7.2	0	2176	1	Cep-Her
K07368.01	Kepler-1974 b	5068	2.56	$248 \pm 4$	$88^{+183}_{-60}$	$54^{+47}_{-25}$	Yes	2.22	6.84	0	2560	1	Cep-Her
K06228.01	Kepler-1644 b	5521	1.43	$-2 \pm 13$	$77^{+144}_{-53}$	$> 767$	No	1.88	21.09	4	2690	1	Unres. Binary
K06186.01	Kepler-1643 b	4918	5.05	$120 \pm 6$	$79^{+182}_{-54}$	$191^{+92}_{-76}$	Yes	2.11	5.34	0	0	1	Cep-Her
K03933.01	Kepler-1699 b	5496	4.16	$-11 \pm 7$	$85^{+106}_{-58}$	$> 889$	No	1.32	3.49	0	2688	1	Unres. Binary
K03916.01	Kepler-1529 b	4974	6.43	$200 \pm 6$	$109^{+117}_{-71}$	$90^{+53}_{-39}$	Yes	2.01	5.34	0	0	1	✓✓
K01804.01	Kepler-957 b	4947	4.52	$24 \pm 9$	$96^{+196}_{-66}$	$> 241$	Yes	6.9	5.91	0	0	1	✓
K07913.01	Kepler-1975 b	4450	3.36	$56 \pm 9$	$96^{+223}_{-66}$	$> 57$	Yes	2.03	24.28	0	2816	1	Cep-Her
K03936.02	Kepler-1930 b	4906	7.1	$170 \pm 4$	$174^{+106}_{-65}$	$115^{+55}_{-49}$	Yes	1.52	13.03	4	0	1	
K03876.01	Kepler-1928 b	5577	4.64	$137 \pm 4$	$148^{+102}_{-87}$	$189^{+150}_{-94}$	Yes	1.86	19.58	0	2048	1	MELANGE-3
K04069.01	Kepler-1938 b	4617	7.82	$6 \pm 16$	$152^{+112}_{-42}$	$> 208$	Yes	1.47	13.06	0	2048	1	✓
K02678.01	Kepler-1313 b	5236	6.13	$142 \pm 3$	$197^{+112}_{-91}$	$174^{+96}_{-72}$	Yes	1.71	3.83	4	2048	1	
K04194.01	Kepler-1565 b	4958	7.4	$133 \pm 7$	$230^{+111}_{-85}$	$174^{+81}_{-72}$	Yes	1.27	1.54	0	0	1	✓✓
K03835.01	Kepler-1521 b	4806	7.82	$117 \pm 5$	$207^{+98}_{-68}$	$176^{+79}_{-69}$	Yes	2.3	47.15	0	2048	1	✓✓
K01838.01	Kepler-970 b	4314	9.23	$36 \pm 14$	$177^{+124}_{-39}$	$> 92$	Yes	2.15	16.74	4	0	1	MELANGE-3
K00063.01	Kepler-63 b	5486	5.49	$89 \pm 4$	$223^{+98}_{-92}$	$542^{+475}_{-256}$	Yes	5.64	9.43	0	4096	1	✓✓
K01199.01	Kepler-786 b	4680	33.06	$83 \pm 6$	$4364^{+359}_{-374}$	$228^{+168}_{-87}$	No	2.31	53.53	0	0	1	Mystery
K03316.01	Kepler-1467 b	5252	6.31	$122 \pm 6$	$230^{+112}_{-98}$	$236^{+151}_{-95}$	Yes	3.11	47.06	0	0	1	✓✓
K01074.01	Kepler-762 b	5921	4.01	$-27 \pm 25$	$245^{+114}_{-100}$	$> 548$	Maybe	15.19	3.77	0	512	1	
K01839.01	Kepler-971 b	5447	6.22	$105 \pm 6$	$306^{+95}_{-115}$	$366^{+290}_{-164}$	Yes	3.93	9.59	0	128	1	
K01833.01	Kepler-968 b	4413	10.46	$10 \pm 18$	$328^{+108}_{-87}$	$> 159$	Yes	1.85	3.69	0	0	1	Theia-520
K01833.03	Kepler-968 c	4413	10.46	$10 \pm 18$	$328^{+108}_{-87}$	$> 159$	Yes	1.63	5.71	0	0	1	Theia-520
K01833.02	Kepler-968 d	4413	10.46	$10 \pm 18$	$328^{+108}_{-87}$	$> 159$	Yes	2.28	7.68	4	0	1	Theia-520
K02675.01	Kepler-1312 b	5584	6.13	$86 \pm 4$	$357^{+75}_{-109}$	$642^{+617}_{-318}$	Yes	2.07	5.45	0	4096	1	✓✓
K02675.02	Kepler-1312 c	5584	6.13	$86 \pm 4$	$357^{+75}_{-109}$	$642^{+617}_{-318}$	Yes	0.94	1.12	0	4096	1	✓✓
K00775.02	Kepler-52 b	4164	11.85	$22 \pm 18$	$360^{+208}_{-97}$	$> 100$	Yes	2.19	7.88	0	0	1	Theia-520
K00775.01	Kepler-52 c	4164	11.85	$22 \pm 18$	$360^{+208}_{-97}$	$> 100$	Yes	2.04	16.38	0	0	1	Theia-520
K00775.03	Kepler-52 d	4164	11.85	$22 \pm 18$	$360^{+208}_{-97}$	$> 100$	Yes	2.03	36.45	0	0	1	Theia-520
K04004.01	Kepler-1933 b	5576	6.21	$85 \pm 3$	$366^{+74}_{-109}$	$642^{+603}_{-318}$	Yes	1.01	4.94	4	0	1	
K02174.03	Kepler-1802 b	4245	11.45	–	$379^{+201}_{-109}$	–	–	1.71	7.73	4	820	0	
K02174.02	Kepler-1802 c	4245	11.45	–	$379^{+201}_{-109}$	–	–	2.05	33.14	0	820	0	
K03935.01	Kepler-1532 b	5554	6.48	$90 \pm 6$	$397^{+71}_{-102}$	$567^{+545}_{-278}$	Yes	1.26	1.09	0	0	1	✓✓
K01801.01	Kepler-955 b	5221	7.5	$79 \pm 4$	$397^{+99}_{-131}$	$536^{+481}_{-237}$	Yes	2.69	14.53	0	0	1	✓✓
K01800.01	Kepler-447 b	5648	6.4	$103 \pm 3$	$420^{+64}_{-78}$	$405^{+355}_{-203}$	Yes	18.49	7.79	4	2048	1	
K03370.02	Kepler-1481 b	4832	9.11	$22 \pm 7$	$407^{+126}_{-118}$	$> 210$	Yes	1.09	5.94	0	0	1	✓
K04156.01	Kepler-1943 b	6002	–	$99 \pm 5$	–	$409^{+520}_{-254}$	No	1.29	4.85	4	518	1	
K00448.01	Kepler-159 b	4511	10.5	$19 \pm 10$	$415^{+160}_{-110}$	$> 161$	Yes	2.3	10.14	0	4096	1	✓
K00448.02	Kepler-159 c	4511	10.5	$19 \pm 10$	$415^{+160}_{-110}$	$> 161$	Yes	2.75	43.59	0	4096	1	✓
K00046.01	Kepler-101 b	5498	–	$100 \pm 5$	–	$419^{+349}_{-196}$	No	5.9	3.49	0	1542	1	
K04169.01	Kepler-1561 b	5742	6.18	$66 \pm 3$	$426^{+74}_{-78}$	$1409^{+1718}_{-788}$	Maybe	0.94	1.01	0	2048	1	✓✓
K02708.01	Kepler-1320 b	4536	10.46	$25 \pm 32$	$430^{+173}_{-113}$	$> 140$	Yes	1.39	0.87	0	0	1	✓
K00119.01	Kepler-108 b	5626	–	$100 \pm 5$	–	$438^{+402}_{-220}$	No	8.2	49.18	0	418	1	
K00119.02	Kepler-108 c	5626	–	$100 \pm 5$	–	$438^{+402}_{-220}$	No	7.78	190.32	4	418	1	
K00323.01	Kepler-523 b	5267	7.6	$49 \pm 4$	$444^{+87}_{-114}$	$1824^{+2706}_{-1047}$	Maybe	1.9	5.84	0	0	1	✓✓
K02115.01	Kepler-67 b	5126	10.39	$83 \pm 10$	$878^{+108}_{-121}$	$458^{+451}_{-206}$	Yes	2.96	15.73	0	0	1	✓✓
K00002.01	Kepler-2 b	6436	–	$83 \pm 4$	–	$485^{+924}_{-353}$	–	16.42	2.2	0	7	1	
K03371.02	Kepler-1482 b	5330	7.7	$52 \pm 3$	$491^{+76}_{-91}$	$1630^{+2285}_{-904}$	Maybe	1.0	12.25	0	640	1	

**Table 1 continued**

Table 1 (continued)

K03497.01	Kepler-1512 b	4894	9.33	14 ± 8	505 <sup>+135</sup> <sub>-115</sub>	> 295	Yes	0.8	20.36	4	4738	1	
K03864.01	Kepler-1698 b	4866	9.49	2 ± 8	518 <sup>+151</sup> <sub>-123</sub>	> 358	Yes	0.9	1.21	0	1024	1	
K03010.01	Kepler-1410 b	3808	14.19	-21 ± 24	523 <sup>+612</sup> <sub>-96</sub>	> 80	Yes	1.39	60.87	0	512	1	
K05447.02	Kepler-1629 b	5585	7.45	–	529 <sup>+62</sup> <sub>-62</sub>	–	–	0.67	3.88	0	0	0	
K04246.01	Kepler-1576 b	5794	7.09	17 ± 8	559 <sup>+103</sup> <sub>-71</sub>	> 613	Yes	0.9	6.98	0	0	1	✓
K03324.01	Kepler-1469 b	5356	8.15	-5 ± 25	562 <sup>+75</sup> <sub>-82</sub>	> 535	Yes	2.53	21.86	0	0	1	✓
K01779.01	Kepler-318 b	5799	7.09	65 ± 3	562 <sup>+106</sup> <sub>-72</sub>	1507 <sup>+1876</sup> <sub>-858</sub>	Yes	3.97	4.66	0	0	1	✓✓
K01779.02	Kepler-318 c	5799	7.09	65 ± 3	562 <sup>+106</sup> <sub>-72</sub>	1507 <sup>+1876</sup> <sub>-858</sub>	Yes	3.1	11.82	4	0	1	
K02084.01	Kepler-1792 b	4942	9.49	13 ± 11	587 <sup>+155</sup> <sub>-144</sub>	> 312	Yes	2.15	4.2	0	0	1	✓
K02035.01	Kepler-1066 b	5847	7.0	60 ± 4	588 <sup>+158</sup> <sub>-85</sub>	2018 <sup>+2616</sup> <sub>-1187</sub>	Maybe	1.96	1.93	4	1024	1	
K03274.01	Kepler-1451 b	5675	7.82	42 ± 4	597 <sup>+79</sup> <sub>-65</sub>	> 316	Yes	2.33	35.62	0	0	1	✓
K01615.01	Kepler-908 b	5670	7.88	67 ± 4	602 <sup>+77</sup> <sub>-65</sub>	1317 <sup>+1607</sup> <sub>-724</sub>	Yes	1.36	1.34	0	4608	1	
K02022.01	Kepler-349 b	5756	7.71	64 ± 6	617 <sup>+103</sup> <sub>-72</sub>	1686 <sup>+2185</sup> <sub>-976</sub>	Yes	1.99	5.93	0	0	1	✓✓
K02022.02	Kepler-349 c	5756	7.71	64 ± 6	617 <sup>+103</sup> <sub>-72</sub>	1686 <sup>+2185</sup> <sub>-976</sub>	Yes	1.97	12.25	0	0	1	✓✓
K00620.01	Kepler-51 b	5635	8.14	48 ± 8	623 <sup>+75</sup> <sub>-65</sub>	> 258	Yes	6.62	45.16	0	0	1	✓
K00620.03	Kepler-51 c	5635	8.14	48 ± 8	623 <sup>+75</sup> <sub>-65</sub>	> 258	Yes	5.49	85.32	4	0	1	
K00620.02	Kepler-51 d	5635	8.14	48 ± 8	623 <sup>+75</sup> <sub>-65</sub>	> 258	Yes	9.04	130.18	0	0	1	✓
K02803.01	Kepler-1877 b	5506	8.37	3 ± 12	624 <sup>+69</sup> <sub>-65</sub>	> 726	Maybe	0.55	2.38	0	1024	1	
K00720.04	Kepler-221 b	5070	9.3	21 ± 5	636 <sup>+120</sup> <sub>-115</sub>	> 341	Yes	1.51	2.8	0	0	1	✓
K00720.01	Kepler-221 c	5070	9.3	21 ± 5	636 <sup>+120</sup> <sub>-115</sub>	> 341	Yes	2.86	5.69	0	0	1	✓
K00720.02	Kepler-221 d	5070	9.3	21 ± 5	636 <sup>+120</sup> <sub>-115</sub>	> 341	Yes	2.57	10.04	0	0	1	✓
K00720.03	Kepler-221 e	5070	9.3	21 ± 5	636 <sup>+120</sup> <sub>-115</sub>	> 341	Yes	2.58	18.37	4	0	1	
K03097.02	Kepler-431 b	6259	16.16	80 ± 3	–	671 <sup>+1092</sup> <sub>-458</sub>	–	0.93	6.8	4	4103	1	
K03097.03	Kepler-431 c	6259	16.16	80 ± 3	–	671 <sup>+1092</sup> <sub>-458</sub>	–	0.93	8.7	4	4103	1	
K03097.01	Kepler-431 d	6259	16.16	80 ± 3	–	671 <sup>+1092</sup> <sub>-458</sub>	–	1.08	11.92	4	4103	1	
K01982.01	Kepler-1781 b	5363	9.17	–	708 <sup>+80</sup> <sub>-78</sub>	–	–	1.95	4.89	0	0	0	
K03375.01	Kepler-1918 b	5522	9.11	–	722 <sup>+75</sup> <sub>-71</sub>	–	–	2.19	47.06	0	0	0	
K01835.02	Kepler-326 b	5142	9.56	8 ± 8	724 <sup>+108</sup> <sub>-103</sub>	> 506	Yes	1.25	2.25	0	1664	1	
K01835.01	Kepler-326 c	5142	9.56	8 ± 8	724 <sup>+108</sup> <sub>-103</sub>	> 506	Yes	1.38	4.58	0	1664	1	
K01835.03	Kepler-326 d	5142	9.56	8 ± 8	724 <sup>+108</sup> <sub>-103</sub>	> 506	Yes	1.31	6.77	0	1664	1	
K01797.01	Kepler-954 b	4736	10.68	4 ± 11	727 <sup>+222</sup> <sub>-184</sub>	> 270	Yes	2.16	16.78	0	1024	1	
K01821.01	Kepler-963 b	5383	9.42	–	748 <sup>+80</sup> <sub>-78</sub>	–	–	2.64	9.98	0	0	0	
K03681.01	Kepler-1514 b	5852	7.87	69 ± 2	758 <sup>+416</sup> <sub>-142</sub>	1302 <sup>+1622</sup> <sub>-754</sub>	Yes	11.94	217.83	0	1540	1	
K03681.02	Kepler-1514 c	5852	7.87	69 ± 2	758 <sup>+416</sup> <sub>-142</sub>	1302 <sup>+1622</sup> <sub>-754</sub>	Yes	1.17	10.51	4	1540	1	
K00647.01	Kepler-634 b	6272	–	77 ± 3	–	768 <sup>+1250</sup> <sub>-527</sub>	–	2.13	5.17	0	7	1	
K02037.01	Kepler-1995 b	4746	10.8	9 ± 21	772 <sup>+213</sup> <sub>-194</sub>	> 223	Yes	3.46	73.76	0	4096	1	✓
K01781.02	Kepler-411 b	4920	10.32	4 ± 3	775 <sup>+164</sup> <sub>-166</sub>	> 396	Yes	2.2	3.01	4	0	1	
K01781.01	Kepler-411 c	4920	10.32	4 ± 3	775 <sup>+164</sup> <sub>-166</sub>	> 396	Yes	3.47	7.83	0	0	1	✓
K07375.01	–	4212	3.88	117 ± 9	104 <sup>+235</sup> <sub>-72</sub>	70 <sup>+37</sup> <sub>-21</sub>	Yes	1.73	4.85	0	2560	1	
K03991.01	–	5226	5.22	98 ± 3	71 <sup>+91</sup> <sub>-48</sub>	354 <sup>+253</sup> <sub>-145</sub>	Maybe	1.37	1.57	0	2176	1	
K01546.01	–	5639	0.9	33 ± 24	75 <sup>+134</sup> <sub>-51</sub>	> 316	Maybe	11.92	0.92	0	2176	1	P <sub>rot</sub> -P <sub>orb</sub>   < 0.1 days
K05482.01	–	5519	0.81	–	77 <sup>+144</sup> <sub>-53</sub>	–	–	2.96	31.71	4	2690	0	
K00064.01	–	5306	2.23	3 ± 4	81 <sup>+129</sup> <sub>-55</sub>	> 567	No	10.49	1.95	4	4614	1	
K06188.01	–	5209	1.62	6 ± 11	84 <sup>+171</sup> <sub>-58</sub>	> 554	Maybe	2.75	1.65	0	2048	1	P <sub>rot</sub> -P <sub>orb</sub>   < 0.1 days
K02695.01	–	5174	2.88	–	86 <sup>+176</sup> <sub>-59</sub>	–	–	20.23	2.5	4	640	0	
K07449.01	–	4928	1.31	–	92 <sup>+194</sup> <sub>-63</sub>	–	–	20.43	1.32	4	2048	0	P <sub>rot</sub> -P <sub>orb</sub>   < 0.1 days
K06130.01	–	4560	3.02	-1 ± 25	94 <sup>+217</sup> <sub>-65</sub>	> 186	Yes	1.45	1.54	0	2976	1	
K06195.01	–	4677	1.42	-8 ± 30	94 <sup>+210</sup> <sub>-65</sub>	> 208	Yes	2.28	1.44	0	2048	1	P <sub>rot</sub> -P <sub>orb</sub>   < 0.1 days

NOTE—EW<sub>Li</sub><sup>\*</sup> is the lithium equivalent width *after* subtracting a constant 7.5 mÅ to account for the Fe I 6707.44 Å blend (see Section 2.3). Two checkmarks (✓✓) denote confirmed planets with two-sided  $t_{\text{gyro}}$  and  $t_{\text{Li}}$  for which both the age and the planet are expected to be reliable. One checkmark (✓) denotes confirmed planets with two-sided  $t_{\text{gyro}}$  only. “Confirmed” planets appear in the machine-readable version before “candidate” planets. Planetary sizes are mostly drawn from (in order of precedence) Petigura et al. (2022), Berger et al. (2020a), and Thompson et al. (2018), and might be unphysical for grazing planets. The bit quality flags for the rotation-based ages,  $Q_{\text{star}}$ , are described in Section 4.1. Concisely summarized, they are: Bit 0:  $T_{\text{eff}}/K \in [3800 - 6200]$ ? Bit 1:  $\log g < 4.2$ ? Bit 2:  $M_G < 3.9$  or  $M_G > 8.5$ ? Bit 3: In KEBC? Bit 4: Large  $d_{\text{xm, Kep-Gaia}}$ ? Bit 5: Confused Kep-Gaia crossmatch? Bit 6: Gaia DR3 non-single star? Bit 7: RUWE > 1.4? Bit 8: Crowded? Bit 9: Far from main sequence? Bit 10: Spectroscopic  $|[\text{Fe}/\text{H}]| > 0.3$ ? Bit 11: S21 CP/CB? Bit 12:  $P_{\text{rot}}$  not in the homogeneous S19 S21 sample? As an example, Kepler-1627 has  $Q_{\text{star}}$  flagged with bit 11 and bit 7. The analogous planet quality bitmask,  $Q_{\text{planet}}$ , has the following meaning. Bit 0: Candidate disposition not reliable? Bit 1: MES < 10? Bit 2: Grazing? For a star to have a high likelihood of being “reliable for gyrochronology” we suggest a  $Q_{\text{star}}$  bitmask with bits 0–10 not raised, and for a planet to be “reliable”, we suggest  $Q_{\text{planet}}$  to be zero.



**Table 2.** Ages of Kepler target stars derived from rotation periods. The full machine-readable table includes 37,892 stars with finite reported gyrochrone ages. The quality bitmask is as in Table 1; requiring bits zero to ten to be null (e.g. using the suggested [python script](#)) yields 23,813 stars for which gyrochronology is likely to be valid.

KIC	Gaia DR3	$T_{\text{eff}}$	$P_{\text{rot}}$	$t_{\text{gyro}}$	$Q_{\text{star}}$
–	–	K	days	Myr	int
8367679	2126373303927600000	5980	7.44	$1062^{+815}_{-369}$	0
11954016	2133663925009149952	5423	24.88	$3230^{+330}_{-301}$	0
1865152	2051034045638265984	5878	7.67	$879^{+891}_{-274}$	0
4447913	2100652256617096832	4654	35.67	> 4000	0
11085139	2129657064120722688	5507	20.49	$2596^{+322}_{-267}$	0
11072100	2131603612018238976	4814	23.87	$2726^{+253}_{-216}$	0
2860187	2052239591421076352	5546	26.29	$3639^{+354}_{-345}$	518
8737920	2106733414911973504	4844	25.94	$3035^{+286}_{-255}$	640
11962034	2130137688140649984	4925	27.61	$3327^{+326}_{-287}$	0
7958926	2126140894655659264	4323	17.0	$1990^{+214}_{-161}$	0

## REFERENCES

- 1331 Aigrain, S., Llama, J., Ceillier, T., et al. 2015, *MNRAS*, 450, 3211
- 1332 Akeson, R. L., Chen, X., Ciardi, D., et al. 2013, *PASP*, 125, 989
- 1333 Alzate, J. A., Bruzual, G., & Díaz-González, D. J. 2021, *MNRAS*, 501, 302
- 1334 Amard, L., & Matt, S. P. 2020, *ApJ*, 889, 108
- 1335 Amard, L., Palacios, A., Charbonnel, C., et al. 2019, *A&A*, 631, A77
- 1336 Amard, L., Roquette, J., & Matt, S. P. 2020, *MNRAS*, 499, 3481
- 1337 Angus, R., Aigrain, S., Foreman-Mackey, D., & McQuillan, A. 2015, *MNRAS*, 450, 1787
- 1338 Angus, R., Morton, T., Aigrain, S., Foreman-Mackey, D., & Rajpaul, V. 2018, *MNRAS*, 474, 2094
- 1339 Angus, R., Morton, T. D., Foreman-Mackey, D., et al. 2019, *AJ*, 158, 173
- 1340 Anthropic. 2024, Claude, v3, Anthropic, Conversational AI model used for editing manuscript and generating testable code; the authors wrote all of the original manuscript text. <https://claude.ai/>
- 1341 Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- 1342 Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, 156, 123
- 1343 Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, *ApJ*, 935, 167
- 1344 Barber, M. G., Mann, A. W., Bush, J. L., et al. 2022, *AJ*, 164, 88
- 1345 Barnes, S. A. 2003, *ApJ*, 586, 464
- 1357 Barnes, S. A., Weingrill, J., Fritzewski, D., Strassmeier, K. G., & Platais, I. 2016, *ApJ*, 823, 16
- 1358 Berger, T. A., Howard, A. W., & Boesgaard, A. M. 2018, *ApJ*, 855, 115
- 1359 Berger, T. A., Huber, D., Gaidos, E., van Saders, J. L., & Weiss, L. M. 2020a, *AJ*, 160, 108
- 1360 Berger, T. A., Huber, D., van Saders, J. L., et al. 2020b, *AJ*, 159, 280
- 1361 Binney, J., Dehnen, W., & Bertelli, G. 2000, *MNRAS*, 318, 658
- 1362 Bland-Hawthorn, J., & Gerhard, O. 2016, *ARA&A*, 54, 529
- 1363 Bonomo, A. S., Sozzetti, A., Lovis, C., et al. 2014, *A&A*, 572, A2
- 1364 Borucki, W. J., Koch, D., Basri, G., et al. 2010, *Science*, 327, 977
- 1365 Bouma, L. G., Curtis, J. L., Hartman, J. D., Winn, J. N., & Bakos, G. Á. 2021, *AJ*, 162, 197
- 1366 Bouma, L. G., Palumbo, E. K., & Hillenbrand, L. A. 2023, *ApJL*, 947, L3
- 1367 Bouma, L. G., Hartman, J. D., Brahm, R., et al. 2020, *AJ*, 160, 239
- 1368 Bouma, L. G., Curtis, J. L., Masuda, K., et al. 2022a, *AJ*, 163, 121
- 1369 Bouma, L. G., Kerr, R., Curtis, J. L., et al. 2022b, *AJ*, 164, 215
- 1370 Boyle, A. W., & Bouma, L. G. 2023, *AJ*, 166, 14
- 1371 Cantat-Gaudin, T., Jordi, C., Vallenari, A., et al. 2018, *A&A*, 618, A93
- 1372 Cantat-Gaudin, T., Anders, F., Castro-Ginard, A., et al. 2020, *A&A*, 640, A1
- 1373 Capistrant, B. K., Soares-Furtado, M., Vanderburg, A., et al. 2024, *AJ*, 167, 54

- Carlos, M., Meléndez, J., Spina, L., et al. 2019, *MNRAS*, 485, 4052
- Castro-Ginard, A., Jordi, C., Luri, X., et al. 2018, *A&A*, 618, A59
- Chaboyer, B., Demarque, P., & Pinsonneault, M. H. 1995, *ApJ*, 441, 865
- Chiti, F., van Saders, J. L., Heintz, T. M., et al. 2024, arXiv e-prints, arXiv:2403.12129
- Choi, J., Dotter, A., Conroy, C., et al. 2016, *ApJ*, 823, 102
- Christiansen, J. L., Zink, J. K., Hardegree-Ullman, K. K., et al. 2023, *AJ*, 166, 248
- Claytor, Z. R., van Saders, J. L., Santos, Â. R. G., et al. 2020, *ApJ*, 888, 43
- Curtis, J. 2024, in American Astronomical Society Meeting Abstracts, Vol. 243, American Astronomical Society Meeting Abstracts, 458.08
- Curtis, J. L., Agüeros, M. A., Douglas, S. T., & Meibom, S. 2019, *ApJ*, 879, 49
- Curtis, J. L., Vanderburg, A., Torres, G., et al. 2018, *AJ*, 155, 173
- Curtis, J. L., Agüeros, M. A., Matt, S. P., et al. 2020, *ApJ*, 904, 140
- David, T. J., Angus, R., Curtis, J. L., et al. 2022, *ApJ*, 933, 114
- David, T. J., Petigura, E. A., Luger, R., et al. 2019, *ApJL*, 885, L12
- David, T. J., Contardo, G., Sandoval, A., et al. 2021, *AJ*, 161, 265
- Denissenkov, P. A., Pinsonneault, M., Terndrup, D. M., & Newsham, G. 2010, *ApJ*, 716, 1269
- do Nascimento, J. D., J., de Almeida, L., Velloso, E. N., et al. 2020, *ApJ*, 898, 173
- Dotter, A. 2016, *ApJS*, 222, 8
- Dunee, R., van Saders, J., Gaidos, E., et al. 2022, *ApJ*, 938, 118
- Engle, S. G. 2024, *ApJ*, 960, 62
- Engle, S. G., & Guinan, E. F. 2023, *ApJL*, 954, L50
- Epstein, C. R., & Pinsonneault, M. H. 2014, *ApJ*, 780, 159
- Fortney, J. J., Marley, M. S., & Barnes, J. W. 2007, *ApJ*, 659, 1661
- Fritzewski, D. J., Barnes, S. A., James, D. J., & Strassmeier, K. G. 2021, *A&A*, 652, A60
- Fritzewski, D. J., Van Reeth, T., Aerts, C., et al. 2024, *A&A*, 681, A13
- Fulton, B. J., & Petigura, E. A. 2018, *AJ*, 156, 264
- Fulton, B. J., Petigura, E. A., Howard, A. W., et al. 2017, *AJ*, 154, 109
- Gaia Collaboration, Babusiaux, C., van Leeuwen, F., et al. 2018, *A&A*, 616, A10
- Gaia Collaboration, Arenou, F., Babusiaux, C., et al. 2023, *A&A*, 674, A34
- Gallet, F., & Bouvier, J. 2015, *A&A*, 577, A98
- Ge, J., Zhang, H., Zang, W., et al. 2022, arXiv e-prints, arXiv:2206.06693
- Gilbert, G. J. 2022, *AJ*, 163, 111
- Gillen, E., Briegal, J. T., Hodgkin, S. T., et al. 2020, *MNRAS*, 492, 1008
- Girardi, L., Groenewegen, M. A. T., Hatziminaoglou, E., & da Costa, L. 2005, *A&A*, 436, 895
- Green, G. M., Schlafly, E., Zucker, C., Speagle, J. S., & Finkbeiner, D. 2019, *ApJ*, 887, 93
- Greiss, S., Steeghs, D., Gänsicke, B. T., et al. 2012, *AJ*, 144, 24
- Gruner, D., Barnes, S. A., & Weingrill, J. 2023, *A&A*, 672, A159
- Gupta, A., & Schlichting, H. E. 2019, *MNRAS*, 487, 24
- Hall, O. J., Davies, G. R., van Saders, J., et al. 2021, *Nature Astronomy*, 5, 707
- Hedges, C., Hughes, A., Zhou, G., et al. 2021, *AJ*, 162, 54
- Herschel, J. F. W. 1864, *Philosophical Transactions of the Royal Society of London Series I*, 154, 1
- Huber, D., Chaplin, W. J., Christensen-Dalsgaard, J., et al. 2013, *ApJ*, 767, 127
- Huber, D., Zinn, J., Bojsen-Hansen, M., et al. 2017, *ApJ*, 844, 102
- Hunter, J. D., et al. 2007, *Computing in science and engineering*, 9, 90
- Isern, J. 2019, *ApJL*, 878, L11
- Izidoro, A., Ogihara, M., Raymond, S. N., et al. 2017, *MNRAS*, 470, 1750
- Jeffries, R. D., Jackson, R. J., Wright, N. J., et al. 2023, *MNRAS*, 523, 802
- Jenkins, J. M., Caldwell, D. A., & Borucki, W. J. 2002, *ApJ*, 564, 495
- Johnson, J. A., Petigura, E. A., Fulton, B. J., et al. 2017, *AJ*, 154, 108
- Kawaler, S. D. 1989, *ApJL*, 343, L65
- Kepler, M., Benisty, M., Müller, A., et al. 2018, *A&A*, 617, A44
- Kerr, R., Kraus, A. L., Krolikowski, D., Bouma, L. G., & Farias, J. P. 2024, arXiv e-prints, arXiv:2406.19530
- Kerr, R. M. P., Rizzuto, A. C., Kraus, A. L., & Offner, S. S. R. 2021, *ApJ*, 917, 23
- Kipping, D., & Bakos, G. 2011, *ApJ*, 733, 36
- Kirk, B., Conroy, K., Prša, A., et al. 2016, *AJ*, 151, 68
- Klein, B., Zicher, N., Kavanagh, R. D., et al. 2022, *MNRAS*, 512, 5067
- Kolbl, R., Marcy, G. W., Isaacson, H., & Howard, A. W. 2015, *AJ*, 149, 18
- Kounkel, M., & Covey, K. 2019, *AJ*, 158, 122
- Kounkel, M., Covey, K., & Stassun, K. G. 2020, *AJ*, 160, 279
- Livingston, J. H., Dai, F., Hirano, T., et al. 2018, *AJ*, 155, 115
- Lorenzo-Oliveira, D., Meléndez, J., Ponte, G., & Galarza, J. Y. 2020, *MNRAS*, 495, L61
- Lorenzo-Oliveira, D., Meléndez, J., Yana Galarza, J., et al. 2019, *MNRAS*, 485, L68
- Lu, Y., Angus, R., Foreman-Mackey, D., & Hattori, S. 2024, *AJ*, 167, 159
- Lu, Y. L., Angus, R., Curtis, J. L., David, T. J., & Kiman, R. 2021, *AJ*, 161, 189

- Lu, Y. L., Curtis, J. L., Angus, R., David, T. J., & Hattori, S. 2022, *AJ*, 164, 251
- Mamajek, E. E., & Hillenbrand, L. A. 2008, *ApJ*, 687, 1264
- Mann, A. W., Gaidos, E., Mace, G. N., et al. 2016, *ApJ*, 818
- Marois, C., Macintosh, B., Barman, T., et al. 2008, *Science*, 322, 1348
- Masuda, K. 2014, *ApJ*, 783, 53
- . 2022a, *ApJ*, 937, 94
- . 2022b, *ApJ*, 933, 195
- Masuda, K., Petigura, E. A., & Hall, O. J. 2022, *MNRAS*, 510, 5623
- Mathur, S., Huber, D., Batalha, N. M., et al. 2017, *ApJS*, 229, 30
- Mathur, S., Claytor, Z. R., Santos, Â. R. G., et al. 2023, *ApJ*, 952, 131
- Matt, S. P., Brun, A. S., Baraffe, I., Bouvier, J., & Chabrier, G. 2015, *ApJL*, 799, L23
- Mazeh, T., Perets, H. B., McQuillan, A., & Goldstein, E. S. 2015, *ApJ*, 801, 3
- Mazzi, A., Girardi, L., Trabucchi, M., et al. 2024, *MNRAS*, 527, 583
- McQuillan, A., Mazeh, T., & Aigrain, S. 2013, *ApJL*, 775, L11
- . 2014, *ApJS*, 211, 24
- Meibom, S., Barnes, S. A., Platais, I., et al. 2015, *Nature*, 517, 589
- Meibom, S., Mathieu, R. D., & Stassun, K. G. 2007, *ApJL*, 665, L155
- Meibom, S., Torres, G., Fressin, F., et al. 2013, *Nature*, 499, 55
- Mills, S. M., & Fabrycky, D. C. 2017, *AJ*, 153, 45
- Miyazaki, S., & Masuda, K. 2023, *AJ*, 166, 209
- Moe, M., & Kratter, K. M. 2021, *MNRAS*, 507, 3593
- Moe, M., Kratter, K. M., & Badenes, C. 2019, *ApJ*, 875, 61
- Mor, R., Robin, A. C., Figueras, F., Roca-Fàbrega, S., & Luri, X. 2019, *A&A*, 624, L1
- Morton, T. D., Bryson, S. T., Coughlin, J. L., et al. 2016, *ApJ*, 822, 86
- Nardiello, D., Malavolta, L., Desidera, S., et al. 2022, *A&A*, 664, A163
- Newton, E. R., Irwin, J., Charbonneau, D., et al. 2016, *ApJ*, 821, 93
- Newton, E. R., Mann, A. W., Kraus, A. L., et al. 2021, *AJ*, 161, 65
- Nordström, B., Mayor, M., Andersen, J., et al. 2004, *A&A*, 418, 989
- Noyes, R. W., Hartmann, L. W., Baliunas, S. L., Duncan, D. K., & Vaughan, A. H. 1984, *ApJ*, 279, 763
- O'Donovan, F. T., Charbonneau, D., Mandushev, G., et al. 2006, *ApJL*, 651, L61
- Owen, J. E. 2019, *Annual Review of Earth and Planetary Sciences*, 47, 67
- Owen, J. E., & Lai, D. 2018, *MNRAS*, 479, 5012
- Pecaut, M. J., & Mamajek, E. E. 2013, *ApJS*, 208, 9
- Petigura, E. A., Howard, A. W., Marcy, G. W., et al. 2017, *AJ*, 154, 107
- Petigura, E. A., Marcy, G. W., Winn, J. N., et al. 2018, *AJ*, 155, 89
- Petigura, E. A., Rogers, J. G., Isaacson, H., et al. 2022, *AJ*, 163, 179
- Plavchan, P., Barclay, T., Gagné, J., et al. 2020, *Nature*, 582, 497
- Quinn, S. N., White, R. J., Latham, D. W., et al. 2012, *ApJL*, 756, L33
- Raghavan, D., McAlister, H. A., Henry, T. J., et al. 2010, *ApJS*, 190, 1
- Rampalli, R., Agüeros, M. A., Curtis, J. L., et al. 2021, *ApJ*, 921, 167
- Rana, N. C. 1991, *ARA&A*, 29, 129
- Rauer, H., Catala, C., Aerts, C., et al. 2014, *Experimental Astronomy*, 38, 249
- Raymond, S. N., Kokubo, E., Morbidelli, A., Morishima, R., & Walsh, K. J. 2014, in *Protostars and Planets VI*, ed. H. Beuther, R. S. Klessen, C. P. Dullemond, & T. Henning, 595–618
- Rebull, L. M., Stauffer, J. R., Hillenbrand, L. A., et al. 2022, *AJ*, 164, 80
- Rebull, L. M., Stauffer, J. R., Bouvier, J., et al. 2016, *AJ*, 152, 114
- Reinhold, T., Bell, K. J., Kuszlewicz, J., Hekker, S., & Shapiro, A. I. 2019, *A&A*, 621, A21
- Reinhold, T., & Gizon, L. 2015, *A&A*, 583, A65
- Reinhold, T., Shapiro, A. I., Solanki, S. K., & Basri, G. 2023, *A&A*, 678, A24
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, *Journal of Astronomical Telescopes, Instruments, and Systems*, 1, 014003
- Rizzuto, A. C., Newton, E. R., Mann, A. W., et al. 2020, *AJ*, 160, 33
- Rogers, J. G., & Owen, J. E. 2021, *MNRAS*, 503, 1526
- Ruiz-Lara, T., Gallart, C., Bernard, E. J., & Cassisi, S. 2020, *Nature Astronomy*, 4, 965
- Sandoval, A., Contardo, G., & David, T. J. 2021, *ApJ*, 911, 117
- Santos, A. R. G., Breton, S. N., Mathur, S., & García, R. A. 2021, *ApJS*, 255, 17
- Santos, A. R. G., García, R. A., Mathur, S., et al. 2019, *ApJS*, 244, 21
- Saunders, N., van Saders, J. L., Lyttle, A. J., et al. 2024, *ApJ*, 962, 138
- Schmidt, S. P., Schlaufman, K. C., & Hamer, J. H. 2024, *AJ*, 168, 109
- See, V., Lu, Y. L., Amard, L., & Roquette, J. 2024, *MNRAS*, 533, 1290
- Sestito, P., & Randich, S. 2005, *A&A*, 442, 615
- Silva Aguirre, V., Bojsen-Hansen, M., Slumstrup, D., et al. 2018, *MNRAS*, 475, 5487
- Simonian, G. V. A., Pinsonneault, M. H., & Terndrup, D. M. 2019, *ApJ*, 871, 174
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Skumanich, A. 1972, *ApJ*, 171, 565



- 1581 Snodgrass, H. B., & Ulrich, R. K. 1990, *ApJ*, 351, 309  
 1582 Soderblom, D. R. 2010, *ARA&A*, 48, 581  
 1583 Southworth, J. 2011, *MNRAS*, 417, 2166  
 1584 Sozzetti, A., Torres, G., Charbonneau, D., et al. 2007, *ApJ*, 664,  
 1585 1190  
 1586 Spada, F., & Lanzafame, A. C. 2020, *A&A*, 636, A76  
 1587 Thompson, S. E., Coughlin, J. L., Hoffman, K., et al. 2018, *ApJS*,  
 1588 235, 38  
 1589 Tofflemire, B. M., Rizzuto, A. C., Newton, E. R., et al. 2021, *AJ*,  
 1590 161, 171  
 1591 Tran, Q. H., Bowler, B. P., Cochran, W. D., et al. 2024, *AJ*, 167,  
 1592 193  
 1593 Vach, S., Zhou, G., Huang, C. X., et al. 2024, *AJ*, 167, 210  
 1594 Van Der Walt, S., Colbert, S. C., & Varoquaux, G. 2011,  
 1595 *Computing in Science & Engineering*, 13, 22  
 1596 Van Eylen, V., Agentoft, C., Lundkvist, M. S., et al. 2018,  
 1597 *MNRAS*, 479, 4786  
 1598 van Saders, J. L., Ceillier, T., Metcalfe, T. S., et al. 2016, *Nature*,  
 1599 529, 181  
 1600 Vidotto, A. A., Gregory, S. G., Jardine, M., et al. 2014, *MNRAS*,  
 1601 441, 2361  
 1602 Villumsen, J. V. 1983, *ApJ*, 274, 632  
 1603 Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *Nature*  
 1604 *Methods*, 17, 261  
 1605 Vogt, S. S., Allen, S. L., Bigelow, B. C., et al. 1994, *SPIE*  
 1606 *Conference Series*, ed. D. L. Crawford & E. R. Craine, Vol. 2198  
 1607 Walkowicz, L. M., & Basri, G. S. 2013, *MNRAS*, 436, 1883  
 1608 Wilson, R. F., Barclay, T., Powell, B. P., et al. 2023, *ApJS*, 269, 5  
 1609 Winn, J. N., Johnson, J. A., Narita, N., et al. 2008, *ApJ*, 682, 1283  
 1610 Wood, M. L., Mann, A. W., Barber, M. G., et al. 2023, *AJ*, 165, 85  
 1611 Xiang, M., & Rix, H.-W. 2022, *Nature*, 603, 599  
 1612 York, D. G., Adelman, J., Anderson, John E., J., et al. 2000, *AJ*,  
 1613 120, 1579  
 1614 Zakhozhay, O. V., Launhardt, R., Trifonov, T., et al. 2022, *A&A*,  
 1615 667, L14  
 1616 Zari, E., Hashemi, H., Brown, A. G. A., Jardine, K., & de Zeeuw,  
 1617 P. T. 2018, *A&A*, 620, A172  
 1618 Zhou, G., Wirth, C. P., Huang, C. X., et al. 2022, *AJ*, 163, 289  
 1619 Ziegler, C., Law, N. M., Morton, T., et al. 2017, *AJ*, 153, 66  
 1620 Zong, W., Fu, J.-N., De Cat, P., et al. 2018, *ApJS*, 238, 30

## APPENDIX

## A. ROTATION PERIOD CATALOG COMPARISON


While we opted for the S19 and S21 rotation period catalogs, the ~~(Deleted: recent)~~ analysis by Reinhold et al. (2023) bears comment. Reinhold et al. (2023) used a similar selection function as Santos et al. (all Kepler stars), and considered a novel period measurement approach based on the gradient of the power spectrum (GPS). Their method likely provides greater completeness, due to its greater sensitivity to signals that are only weakly periodic.

Figure 10 shows histograms of the differences between reported periods from these various studies. The left panel compares overlapping stars from McQuillan et al. (2014) and the Santos et al. studies. The periods agree at a precision of  $\lesssim 0.01 P_{\text{rot}}$  for  $P_{\text{rot}} \lesssim 15$  days, and at  $\lesssim 0.03 P_{\text{rot}}$  for  $P_{\text{rot}} \approx 30$  days. A small bias exists at  $P_{\text{rot}} \approx 10$  days, in the sense that the Santos periods tend to be  $\approx 1\%$  faster for such stars.

The right panel compares overlapping periods from the Reinhold et al. (2023) GPS method and the Santos et al. studies. The periods agree at a precision of  $\lesssim 0.20 P_{\text{rot}}$  for  $P_{\text{rot}} \lesssim 15$  days, and at  $\lesssim 0.30 P_{\text{rot}}$  for  $P_{\text{rot}} \approx 30$  days. A bias develops past  $P_{\text{rot}} \gtrsim 30$  days, in the sense that the Santos et al. periods tend to be 10-20% longer than the GPS periods. The origin of this larger scatter could be connected to the need for the GPS method to determine the “ $\alpha$  calibration factor”, which is known only at a statistical level for large stellar populations (see Reinhold et al. 2023). While we considered deriving independent rotation-based ages using the Reinhold et al. (2023) GPS periods with an inflated empirical uncertainty, we ultimately opted against this path. A rotation period difference of 20-30% away from the “true period” could sufficiently skew a star’s age that we prefer to restrict our attention to the stars for which precision is a possible outcome.

## B. DEFINING AN “ISOCRONALLY FAR FROM THE MAIN SEQUENCE” BOUNDARY

The (arbitrary) method used to construct the orange line in the lower panel of Figure 1 was as follows. The line is defined as the maximum of two numerically-determined functions,  $f_0(T_{\text{eff}})$  and  $f_1(T_{\text{eff}})$ . We defined  $f_0$  by fitting an  $N^{\text{th}}$ -order polynomial to the stars with rotation periods for which  $t_{\text{B20,iso}}/(\sigma_{t,\text{B20,iso}}) \approx 3$  and  $T_{\text{eff}} \in [3700, 5900]$  K, where  $t_{\text{B20,iso}}$  was the isochronal age reported by Berger et al. (2020b). We let the order of the fit vary from  $N=1$  to 10, minimized the Bayesian Information Criterion, and found  $N=6$ . By eye, this yielded a plausible locus for  $T_{\text{eff}} \lesssim 5300$  K. However, for F and G dwarfs, the resulting locus allowed for stars that were “too evolved”; the density of KOIs was anomalously low in this region of the  $\log g$  vs.  $T_{\text{eff}}$  plane. We therefore visually selected KOIs that appeared to be near the main sequence – i.e. most of the yellow points in Figure 1 – and used them to fit a separate polynomial  $f_1$  through a similar BIC-minimization procedure, which yielded  $N=3$ . This polynomial,  $f_1$ , is shown in very faint opacity in Figure 1. The portion of the orange locus from  $\approx 5300$ – $6200$  K is set as  $f_1 + c$ , for  $c$  a constant offset that we defined to be 0.1 dex above the “KOI main sequence”. The exact break-point in temperature is automatically set by



**Figure 10. Comparison of reported literature stellar rotation periods.** “Santos” refers to the concatenation of Santos et al. (2019) and Santos et al. (2021). “M14” refers to McQuillan et al. (2014). “R23GPS” refers to the Gradient of Power Spectrum periods from Reinhold et al. (2023). The background is a 2-D histogram with a logarithmic color stretch that counts overlapping stars between these studies. The black errorbars show the median and  $\pm 1\sigma$  range of period the differences in 1-day bins. The solid gray “empirical  $\sigma_P/P$ ” line in the left panel shows the expected  $\pm 1\sigma$  range that would be spanned if the independent  $P_{\text{rot}}$  measurements had gaussian uncertainties drawn following the empirical estimate described in Section 2.1. Note the different vertical scales.

max( $f_0, f_1 + c$ ). While we do report rotation-based ages for stars above this orange locus, they are flagged as not being near the main sequence in the  $\log g$  vs.  $T_{\text{eff}}$  plane.

### C. CASES WITH “INCONSISTENT” ROTATION AND LITHIUM-BASED AGES

In this section we discuss the systems with confirmed planets that have nominally discrepant rotation and lithium-based ages. We find that only one of them, Kepler-786, is interesting.

*Kepler-1* — TrES-2/Kepler-1 (O’Donovan et al. 2006) is a  $\approx 2.5$  day near-grazing hot Jupiter orbiting a G0 V primary which has been studied in detail by multiple investigators (e.g. Sozzetti et al. 2007; Winn et al. 2008; Kipping & Bakos 2011; Southworth 2011). Our lithium-based age of this system ( $785^{+845}_{-419}$  Myr), derived from  $\text{EW}_{\text{Li}}^* = 81 \pm 3$  mÅ, qualitatively agrees with the previously noted lithium abundance (Sozzetti et al. 2007). However, we detect no photometric rotation signal, and the spectroscopic  $v \sin i$  is low. The lithium age is strongly asymmetric because of the scatter in  $\text{EW}_{\text{Li}}$  vs.  $T_{\text{eff}}$  over ages  $t \gtrsim 1$  Gyr for early G dwarfs. We calculate a  $2\sigma$  upper limit on  $t_{\text{Li}}$  of 2.8 Gyr, and a  $3\sigma$  upper limit of 6.1 Gyr. The non-detection of rotation, while mildly surprising, is not shocking if the system’s true age is in this long tail.

*Kepler-101* — This  $\approx 5500$  K star has  $\text{EW}_{\text{Li}}^* = 100 \pm 5$  mÅ, nominally implying  $t_{\text{Li}} \approx 200$ –800 Myr, but no rotation detection. Our automated flags noted that this star has a low surface gravity, high luminosity, and is far from the main sequence. Bonomo et al. (2014) concur: this star is a sub-giant slightly more massive than the Sun; no rotation detection is expected; the lithium likely simply survived over the star’s main sequence lifetime.

*Kepler-108* — This system of two mutually inclined giant planets (Mills & Fabrycky 2017) has a  $T_{\text{eff}} \approx 5600$  K host star with a strong lithium detection, and no rotation detection. The host star is entering the red giant branch; this has been previously noted through asteroseismology (Huber et al. 2013), and our analysis of the Keck/HIRES spectrum with SpecMatch-Synth (Petigura et al. 2017) concurs, yielding  $T_{\text{eff}} = 5668 \pm 100$  K,  $\log g = 3.8 \pm 0.1$ ,  $[\text{Fe}/\text{H}] = 0.38 \pm 0.06$ , and  $v \sin i = 3.2 \pm 1.0$  km s $^{-1}$ . Given the star’s high mass, the high lithium content and lack of rotation detection is not surprising.

*Kepler-786* — This K3 V star has a surprisingly high lithium content. The spectrum is single-lined, with SpecMatch-Synth derived parameters of  $T_{\text{eff}} = 4769 \pm 100$  K,  $\log g = 4.5 \pm 0.1$ ,  $[\text{Fe}/\text{H}] = 0.11 \pm 0.06$ , and  $v \sin i = 2.4 \pm 0.6$  km s $^{-1}$ . Yet with  $\text{EW}_{\text{Li}}^* = 83 \pm 6$  mÅ, the lithium age of  $228^{+168}_{-87}$  Myr would predict **(Replaced: an obvious replaced with: a short-period)** rotation signal. **(Replaced: None is present in the Kepler light curve replaced with: The actual signal that is present has an amplitude of  $\approx 0.1$ –0.3% and a period of  $\approx 33$  days)**, consistent with the low  $v \sin i$ . Out of all “apparently discrepant” lithium and rotation measurements discussed in this appendix, this is the only one that seems to remain discrepant after scrutiny. The Ca II doublet is in emission, with  $R'_{\text{HK}} = -4.7 \pm 0.5$ , which suggests an age at least as old as the Hyades (Mamajek & Hillenbrand 2008). The Balmer lines are in absorption, and display no obvious signatures of youth.


*Kepler-1644* — The rotational age of this system ( $77^{+144}_{-53}$  Myr) is nominally much lower than the lithium limit ( $> 767$  Myr). The Kepler light curve shows a  $\approx 1\%$  amplitude 1.4 day rotation signal with many flares. However the automated quality flags note that the star has low (photometric) surface gravity, a high RUWE ( $\text{RUWE}_{\text{DR3}} = 7.0$ ), and that it is far from the main sequence. The Keck/HIRES spectrum also shows visually narrow lines, with a SpecMatch-Synth  $v \sin i \leq 2$  km s $^{-1}$ . However, we performed a cross-correlation between the HIRES spectrum and the nearest matches in the Keck/HIRES template library (Kolbl et al. 2015), and found that on 31 July 2022 (UT) the system displayed a broad CCF with a blended second component at  $\approx +26$  km s $^{-1}$  relative to the primary, with a best-fit flux-ratio of  $\approx 3.4\%$ , and a preferred  $T_{\text{eff,B}} \approx 4400$  K. The spectrum and astrometric excess noise therefore point to this system being an unresolved binary, which calls the reliability of the rotational age into question. The non-detection of the companion from Robo-AO imaging at Palomar (Ziegler et al. 2017) suggests that the companion(s) are likely within  $\rho \lesssim 0.3''$ .

*Kepler-1699* — This system is in a similar qualitative regime as Kepler-1644, with an apparently young  $t_{\text{gyro}} = 85^{+106}_{-58}$  Myr derived from a  $\approx 2\%$  amplitude 4.2 day rotation signal, and no evidence for lithium. The system has  $\text{RUWE}_{\text{DR3}} = 19.1$ . From the same style of CCF analysis from the Keck/HIRES spectrum acquired on 31 Aug 2022 (UT), we also find a double-peaked CCF, in this case with a secondary component at  $\approx -16$  km s $^{-1}$  relative to the primary with  $T_{\text{eff,B}} \approx 4900$  K. This putative companion is similarly not detected in high-resolution imaging (Ziegler et al. 2017). The rotational age is questionable, given that we do not know which source the rotation signal is from, or whether the stars have interacted.

*Kepler-1943* — This system nominally has a  $409^{+520}_{-254}$  Myr lithium age, and no reported rotation detection. However, the star is flagged as being over-luminous, low-surface gravity, and far from the main sequence. In other words, it is a subgiant.

*Kepler-639, Kepler-320, Kepler-1719, Kepler-1876, Kepler-1072, Kepler-1743, Kepler-1929, Kepler-1488* — These eight systems all have nominally two-sided lithium age posteriors between 1 and 3 Gyr, and yet lack rotation period detections. All are subgiants, flagged with  $Q_{\text{star}}$  under various combinations of bits 1, 2, and 9.





**Figure 11. Rotation-based age vs. effective temperature.** The entire Kepler sample is shown on the left. KOI hosts are shown on the right. Each bar shows the  $\pm 1\sigma$  uncertainty for a star’s age as derived from its rotation period. Notable features are discussed in Appendix D.

#### D. AGE DIAGNOSTICS

1700


Figure 11 is a visualization of  $t_{\text{gyro}}$  vs.  $T_{\text{eff}}$  for our rotation-based age catalog. Each bar denotes the  $\pm 1\sigma$  uncertainty for a star’s (Deleted: rotation-based )age: this plot can be viewed as the transformation of the left panel of Figure 2. The overdensity of  $P_{\text{rot}} \approx 20$  day G dwarfs corresponds to the large overdensity at  $t_{\text{gyro}} \approx 3$  Gyr. The (non-physical) deficit of  $\sim 22$  day rotators appears as a deficit between 5500–6000 K. A second deficit is also visible from 3800–5000 K; it is associated with the “intermediate period gap”, which may be associated with a transition from spot-dominated to faculae-dominated light curves (e.g. Reinhold et al. 2019)(Added: , or with rapid spin-down at Rossby numbers near 0.5 (Lu et al. 2022)). At  $\approx 1$  Gyr, stalled spin-down yields larger uncertainties for K dwarfs. The pile-up near  $\approx 5$  Gyr is imposed by our choice of prior; given that gyro-interp returns non-calibrated ages in this regime, we truncated our analysis at 4 Gyr.

1709


#### E. WHAT IF WE ONLY CONSIDERED MCQUILLAN’S PERIODS?

We argued in Section 2.1 that adopting the periods from Santos et al. (2019, 2021) yielded the best possible balance between homogeneity and sensitivity for both KOIs and for the broader Kepler sample. Nonetheless, it is interesting to consider what would happen if we were to analyze only the rotation periods from McQuillan et al. (2014). Figures 12 and 13 repeat the analysis, but make this alternate choice. Figure 12 shows that while the overall contours of the  $P_{\text{rot}}$  vs.  $T_{\text{eff}}$  distributions are similar, the Santos et al. distribution includes more stars, particularly at longer rotation periods. This is connected to a stronger noise-dependent cutoff in the McQuillan et al. (2014) sample than in the Santos et al. samples (see Masuda 2022b Figures 10 and 15). Figure 13 shows the impact on the inferred rotation-based age histograms: while the Santos et al. distribution peaks at 2.8–3.0 Gyr, the McQuillan et al. distribution peaks at 2.4–2.6 Gyr. A number of metrics shift. For instance, the ratio of the star formation rate 3 Gyr age to today is  $2.81 \pm 0.12$  assuming the Santos et al. periods, and  $2.24 \pm 0.11$  assuming the McQuillan et al. periods. Similarly, the ratio of “old” (2–3 Gyr) to “young” (0–1 Gyr) stars shifts from 2.1 in the Santos et al. case to 1.9 in the McQuillan et al. case. The K dwarf age distributions (Figure 13 lower panel) however appear quite similar. Generally, these plots support the conclusion that the  $t_{\text{gyro}}$  age distribution in the Kepler field is non-uniform, with a peak in the (Replaced: 2.5 replaced with: 2.3)–3 Gyr range. Nonetheless, some of the details do shift depending on the choice of rotation detection pipeline.

1722



**Figure 12.** Variant of Figure 2, comparing the default rotation periods (*left*) from Santos et al. (2019, 2021) against the McQuillan et al. (2014) rotation periods (*right*). All stars are required to be amenable to rotation-based age dating (i.e. none of bit flags zero through **(Replaced: nine replaced with: ten)** raised).



**Figure 13.** Variant of Figure 6, comparing our default adopted age distribution (left) based on the Santos et al. (2019, 2021) rotation periods against the age distribution (middle) based only on rotation periods reported by McQuillan et al. (2014).

---

## List of Changes

---

Deleted: ~~accurately~~ on page 1.

Deleted: ~~We find that the scarcity of sub-gigayear systems can be attributed to the star formation rate in the Kepler field dropping by a factor of  $2.81 \pm 0.12$  over the past 3 Gyr. We observe this trend both for known planet hosts and for the parent stellar sample. This “demographic cliff” in the Galaxy’s star formation history has been previously reported, and its confirmation helps clarify the age distribution of the known transiting exoplanets.~~ on page 1.

Added: The age distribution that we observe, rather than being uniform, shows that the youngest stars in the Kepler field are 3-5 times rarer than stars 3 Gyr old. This trend holds for both known planet hosts and for the parent stellar sample. We attribute this “demographic cliff” to a combination of kinematic heating and a declining star formation rate in the Galaxy’s thin disk, and highlight its impact on the age distribution of known transiting exoplanets. , on page 1.

Deleted: ~~transiting~~ on page 2, line 81.

Added: **stellar** , on page 2, line 83.

Deleted: ~~such~~ on page 2, line 94.

Deleted: ~~a total of~~ on page 2, line 97.

Added: **than the 50 currently known**, on page 2, line 99.

Replaced: ~~binary stars~~ replaced with: **stars that are binaries and that have high metallicity**, on page 2, line 151.

Deleted: ~~than in the past~~ on page 2, line 153.

Replaced: ~~the available single-star~~ replaced with: **our**, on page 2, line 160.

Deleted: ~~points~~ on page 3.

Added: **apparent** , on page 3.

Replaced: ~~lower~~ replaced with: **upper**, on page 3.

Replaced: ~~lower~~ replaced with: **upper**, on page 3.

Replaced: ~~“mean-fits” to the rotation sequences of open clusters~~ replaced with: **polynomials fitted to the slow rotation sequences of open clusters**, on page 4.

Deleted: ~~mean~~ on page 4.

Replaced: ~~set of all~~ replaced with: **sample of**, on page 3, line 205.

Deleted: ~~and precise~~ on page 3, line 212.

Added: The gray lines in this figure are the seventh-order polynomials described by Bouma et al. (2023) that fit the slow rotation sequences of the Pleiades (Rebull et al. 2016), Praesepe (Rampalli et al. 2021), NGC-6811 (Curtis et al. 2019), NGC-6819 (Meibom et al. 2015), Ruprecht-147 (Curtis et al. 2020), and M67 (Barnes et al. 2016; Dungee et al. 2022; Gruner et al. 2023),, on page 4, line 260.

Replaced: ~~these~~ replaced with: **our adopted**, on page 4, line 268.

Deleted: ~~adopted~~ on page 4, line 270.

Deleted: ~~on achievable precision~~ on page 7.

Added: **v0.6**;, on page 6, line 395.

Added: **in v0.6**, on page 6, line 414.

Deleted: ~~Kepler~~ on page 6, line 416.

Added: **therefore**, on page 6, line 418.

Added: Finally, we opted to modify **gyro-interp** in **v0.6** to incorporate an age-dependent population-level Gaussian scatter, as detailed in the change log at [gyro-interp.readthedocs.io](https://gyro-interp.readthedocs.io). While age precisions at  $\leq 1$  Gyr are unaffected, this update accounts for the observed intrinsic scatter in Ruprecht-147 and M67 being larger than in Praesepe or NGC-6811, potentially due to differential rotation (e.g., Epstein & Pinsonneault 2014). For  $\approx 2$ -4 Gyr old stars with rotation periods measured at  $\approx 5\%$  precision, this change yields a  $\lesssim 50\%$  increase in age uncertainties., on page 6, line 419.

Replaced: ~~nine~~ replaced with: **ten**, on page 6, line 446.

Replaced: ~~11~~ replaced with: **12**, on page 6, line 449.



Added: **We also considered a deeper cut ( $\Delta G < 5$ ), and found it had negligible impact on our conclusions.**, on page 7, line 518.

Replaced: **HR** replaced with: **HR or Kiel**, on page 7, line 531.

Added: **s**, on page 7, line 532.

Added: *Metallicity (bit 10)*—**Models suggest that stars with non-solar metallicities spin down at different rates than solar-metallicity stars (Amard & Matt 2020). This expectation has been hard to verify due to a dearth of nearby non-solar metallicity open clusters. However, if true, this could cause systematic biases for gyrochrone ages (e.g. Figure 9 of Clayton et al. 2020). We therefore flagged stars with spectroscopic  $|\text{[Fe/H]}| > 0.3$  from either LAMOST DR5 (Zong et al. 2018) or CKS (Petigura et al. 2022) as being outside the range of our gyrochronology calibration. We discuss limitations of this approach in Section 8.**, on page 8, line 541.

Replaced: **10** replaced with: **11**, on page 8, line 553.

Replaced: **11** replaced with: **12**, on page 8, line 563.

Added: **, and for this plot, no attempt was made to exclude known binaries**, on page 9.

Replaced: **all stars with temperatures of 3800–6200 K for which we calculated rotational ages** replaced with: **stars with temperatures of 3800–6200 K**, on page 10.

Added: **metallicity**, on page 10.

Added: **(see Section 4.1)**, on page 10.

Replaced: **have more precise ages due to their fast spin-down** replaced with: **have more precise ages**, on page 10.

Replaced: **rotation-derived** replaced with: **rotational**, on page 10.

Replaced: **using CMD fitting** replaced with: **from CMD fits to other regions of the Galaxy**, on page 10.

Added: **, and decreases rapidly at older ages**, on page 10.

Replaced: **assume Poisson statistics** replaced with: **are Poissonian**, on page 10, line 691.

Replaced: **yields** replaced with: **shows**, on page 10, line 694.

Replaced: **seems to** replaced with: **might**, on page 11, line 711.

Added: *Completeness*—**Figure 6 has a gray overlay beyond  $\gtrsim 3$  Gyr because Kepler's sensitivity to rotation signals diminishes at old ages.**, on page 11, line 736.

Deleted: **this issue** on page 11, line 739.

Added: **Kepler's completeness to rotation signals using MIST-derived isochrone ages**, on page 11, line 739.

Added: **that are**, on page 11, line 742.

Deleted: **by Kepler** on page 11, line 742.

Deleted: **This trend is opposite in functional form to our derived age distributions.** on page 11, line 744.

Added: **This estimate suggests that while aspects of Figure 6 might be interpretable in terms of the Galaxy's star formation history at  $\lesssim 3$  Gyr, at older ages incompleteness precludes any such interpretation.**, on page 11, line 745.

Added: *Temperature sensitivity*—**There are quantitative differences in Figure 6 between the K dwarfs (4400–5400 K) and the FGK stars (3800–6200 K). For instance, the distribution peaks at 2.3 Gyr for K dwarfs, and at 2.9 Gyr for the FGK sample. The peak is also sharper for the K dwarfs; its decrease from 2.3–3 Gyr is in an age range in which we believe the rotation period catalog completeness to be near unity. We show further details of the  $t_{\text{gyro}} - T_{\text{eff}}$  distributions in Appendix D, and propose possible explanations for these differences in Section 8.2.**, on page 11, line 750.

Deleted: **Sub-gigayear planets can be particularly informative for studies of planet evolution.** on page 11, line 762.

Added: **This analysis provides an important path toward expanding the population of young and evolving exoplanets.**, on page 11, line 763.

Deleted: **a total of** on page 11, line 776.

Replaced: **reported by** replaced with: **from**, on page 11, line 786.

Deleted: **rotation-based** on page 11, line 794.

Replaced: **Star symbols denote 46 planets that are characterized in this work to have  $t_{\text{gyro}} < 1$  Gyr at  $2\sigma$ ; circles are planets from the literature meeting the same age requirement, but with a heterogeneous set of age provenances. Gray points are transiting planets from the NASA Exoplanet Archive older than 1 Gyr. If one were to instead select for precise ages ( $t/\sigma_t > 3$ ), this would hide  $\approx 10$   $< 0.5$  Gyr planets, and add  $\approx 15$   $0.5$ – $1$  Gyr planets. For this plot, we require our Kepler stellar hosts to not have  $Q_{\text{star}}$  bits 0–9 raised, and for the planets to similarly not have quality flags raised.** replaced with:

Systems are selected to have  $t < 1$  Gyr at  $2\sigma$ . Star symbols denote 46 planets characterized in this work to have such ages at the highest confidence, meaning that none of  $Q_{\text{star}}$  bits 0–10 are raised, and the planets have no quality flags raised. Transparent crosses denote 17 planets for which either the planet is grazing, or the star has  $\text{RUWE} > 1.4$ . Gray points are from the NASA Exoplanet Archive. , on page 12.

Replaced: **nine** replaced with: **ten**, on page 12, line 852.

Replaced: **This** replaced with: **The stars in this**, on page 12, line 859.

Deleted: **s** on page 12, line 860.

Deleted: **of which are** on page 12, line 861.

Replaced: **rank among** replaced with: **could be**, on page 12, line 877.

Added: **and are** , on page 12, line 879.

Added: Two separate astrophysical effects might explain our observed age distribution (Figure 6): a declining star formation rate (SFR), and kinematic heating. Kinematic heating is associated with an increase in both stellar velocity dispersions and the disk’s vertical scale height over time (e.g. Villumsen 1983; Schmidt et al. 2024). The Kepler field points toward  $(l, b) = (76.3^\circ, 13.5^\circ)$ . A typical Kepler star at 1 kpc is therefore  $\approx 230$  pc above the galactic plane, a significant fraction of the thin disk’s mass-weighted  $\approx 300$  pc scale height (Bland-Hawthorn & Gerhard 2016). , on page 13, line 888.

Added: We assessed the relative importance of these two effects using the TRILEGAL Monte Carlo population synthesis tool , on page 13, line 899.

Added: TRILEGAL treats the Galaxy as a linear combination of a thin disk, thick disk, halo, and bulge, and allows for two possible SFRs: constant, and a “two-step” SFR in which the thin disk SFR is 50% larger from 1–4 Gyr than at other ages. This two-step SFR was introduced to improve the model’s ability to match star counts from Hipparcos and 2MASS , on page 13, line 901.

Added: For kinematic heating, TRILEGAL assumes that the disk scale height grows as  $h_z = z_0(1 + t/t_0)^\alpha$ , for  $\alpha = 5/3$ ,  $z_0 = 94.7$  pc, and  $t_0 = 5.55$  Gyr, similar to Rana (1991). We queried the web interface toward the Kepler field, and toward the galactic plane (at  $l = 76.3^\circ$ ) to negate the effects of kinematic heating. We limited the output to stars with apparent Kepler magnitudes brighter than 16, and set the field of view to  $10 \text{ deg}^2$ , the maximum available through the web interface. , on page 13, line 909.

Added: Figure 8 compares the results against our observed age distribution. The middle panel shows that assuming a constant SFR, kinematic heating alone can yield a two-fold difference in the abundance of the youngest stars relative to those  $\approx 3$  Gyr old. The right-most panel shows that additionally including the time-variable SFR yields the best agreement with the slope of our observed age distribution. , on page 13, line 918.

Deleted: ~~Most stars observed by Kepler are in the thin disk. This can be verified by following Gaia Collaboration et al. (2018), and labeling stars with 2D tangential velocities  $v_T < 40 \text{ km s}^{-1}$  as thin disk members, and those with  $60 < v_T < 150 \text{ km s}^{-1}$  as thick disk members. These criteria yield 95,694 thin and 54,039 thick disk stars observed by Kepler. The thin disk stars have a threefold larger detected rotation period fraction: 35,675 rotators are in the thin disk, while 7,312 are in the thick disk. Further imposing the quality flags discussed in Section 4.1 yields 17,755 thin disk stars for which gyrochronology is applicable, and 2,462 thick disk stars.~~ on page 13, line 926.

Deleted: ~~Classifying Kepler stars as thin vs. thick disk members helps connect them to previous work on the Galaxy’s star formation history. On extragalactic scales, the star formation rate in spiral galaxies with mass similar to the Milky Way peaked  $\approx 10$  Gyr ago, and has since decreased by an order of magnitude.~~ on page 13, line 939.

Added: Despite the expected differences between pencil-beam surveys and volume-complete samples, our observed age distribution qualitatively agrees with a number of previous studies., on page 13, line 944.

Replaced: ~~In our own galaxy, previous studies have~~ replaced with: **Previous work has**, on page 13, line 948.

Replaced: ~~and Isern (2019) for instance focused on stars (and white dwarfs)~~ replaced with: **and Mazzi et al. (2024) for instance focused on stars**, on page 13, line 954.

Added: **weak** , on page 14, line 974.

Added:  $\approx 1$  Gyr and , on page 14, line 975.

Added: **s**, on page 14, line 975.

Deleted: ~~The  $\approx 1$  Gyr spike however is not recovered.~~ on page 14, line 976.

Replaced: ~~those studies is their improved accuracy at  $<$~~  replaced with: **these previous studies is their accuracy below**, on page 14, line 982.

Added: (see Figure 5), on page 14, line 984.

Deleted: ~~stellar~~ on page 14, line 987.

Deleted: ~~stellar~~ on page 14, line 988.

Added: Astrophysical factors include metallicity, latitudinal differential rotation, mergers, and tidal spin-up. Observational limitations include the difficulty of measuring small-amplitude, long-period signals that are only quasi-coherent due to the starspot lifetime approaching the stellar rotation period. There is also the issue that our age-dating approach is not calibrated beyond 4 Gyr. In the following, we attempt to quantify the importance of these effects in our analysis, ordered by what we perceive to be their descending importance., on page 14, line 989.

Added: Simonian et al. (2019) for instance found that 59% of Kepler stars with rotation periods between 1.5 and 7 days lie 0.3 mag above the main sequence, compared with 28% of their full rotation sample. We attempted to mitigate the effects of binarity with five distinct quality flags (Section 4.1). This attempt was at least somewhat effective, based on the peak near zero in the raw distributions of Figure 6 being flattened in the cleaned distributions. However, the completeness of our quality flags is hard to assess. One specific concern is that these flags might have minimal sensitivity to binaries with separations of  $\approx 0.05$ -0.5 AU, for all but the highest mass ratios. This inner limit is set by the onset of detectable ellipsoidal variations in Kirk et al. (2016); the outer limit is set by the typical orbital periods of astrometric binaries in Gaia DR3 , on page 14.

Added: If we were not at all sensitive to binaries with separations of 0.05-0.5 AU, then based on the binary fraction and separation distribution from Raghavan et al. (2010), as much as  $\approx 5\%$  of our “gyro applicable” sample (1,190/23,813 stars) might be contaminated. In a worst case scenario, all of these unidentified binaries might induce  $P_{\text{rot}} < 10$  days, which would then be mixed in with the bona fide young, single stars. This would constitute a major source of contamination, given that our nominal sample of apparently clean stars had 2,706 stars with  $P_{\text{rot}} < 10$  days. , on page 14.

Added: Figure 9 shows the rotation-based age distribution calculated by randomly dropping 1,190 of the  $P_{\text{rot}} < 10$  day stars. Relative to the nominal age distribution, there are three important differences. There is a much larger drop in the star formation rate over the past three gigayears, by a factor of  $4.81 \pm 0.26$ , rather than  $2.81 \pm 0.12$ . The rate of decline in the age distribution is also constant, rather than having a “kink” at 1.5 Gyr. And finally, the corrected distributions show closer agreement with both the SFHs derived from CMD fitting, and also with the KOI host distribution, which is almost devoid of close binaries within 1 AU (e.g. Moe & Kratter 2021, and references therein). We caution however that this correction assumes that *none* of our quality flags were able to identify any of these unresolved binaries. Since we did in fact include flags based on e.g., location in the HR diagram, this correction is best interpreted as an upper limit on the possible impact of binarity., on page 14.

Added: *Rotational outliers:* , on page 14.

Replaced: ~~young stars known to host planets~~ replaced with: each star, on page 14.

Deleted: ~~open~~ on page 14.

Deleted: ~~open~~ on page 14.

Added: While most of these outliers have known bad quality flags, some do not., on page 14.

Added: A separate plausible origin for rotational outliers is that they come from erroneously reported rotation periods. While the large surveys do show good internal consistency (Appendix E), the referee correctly pointed out that this is only a comparative test of methods, rather than an ability to recover ground truth (e.g. Aigrain et al. 2015). One could imagine an adversarial scenario in which, say, 10-20% of stars with  $P_{\text{rot}} > 20$  days are erroneously reported, and in fact have much longer rotation periods. In this scenario, a similar analysis to that in Figure 9 showed that our analysis would overestimate the degree by which the SFR in the Kepler field has decreased, by 7-14%. , on page 15.

Added: *Weakened magnetic braking:* If stellar rotation rates and spot-induced photometric amplitudes remained *constant* after  $\approx 3$  Gyr, then older stars could contaminate the peaks in Figure 6. This possibility warrants consideration, given asteroseismic evidence suggesting slowed spin-down rates after  $\approx 4$  Gyr for Sun-like stars (van Saders et al. 2016; Hall et al. 2021; Saunders et al. 2024), and the “long-period edge” in the  $P_{\text{rot}}-T_{\text{eff}}$  distribution which may pile up over similar timescales (David et al. 2022). However, this stalling is thought to happen at  $> 5$  Gyr for late-G dwarfs and early K dwarfs (e.g. Figure 4 by Saunders et al. 2024), which means that it cannot account for the 2.3 Gyr peak in the 4400-5400 K age distribution shown in Figure 6., on page 15.

Added: Additionally, Masuda (2022a) found that the McQuillan et al. (2014) sample, discussed in Appendix E, includes few stars older than  $\approx 4$  Gyr, based on isochrone-fitting to a high-quality subset of Kepler planet hosts. Some solar twins have also been shown to have rotation periods of 30-50 days (Lorenzo-Oliveira et al. 2019, 2020; do Nascimento et al. 2020), indicating that a completely stalled spin-down may be unlikely. These arguments suggest that while the peak in the age distribution is not primarily caused by an accumulation of much older stars with slowed spin-down rates, contamination by such stars beyond  $\approx 3$  Gyr remains possible, particularly for stars hotter than the Sun., on page 15.

Added: *Metallicity*: Stars with higher metallicity are predicted to have longer convective turnover times, which may slow their rate of spin-down (Amard & Matt 2020). While recent observational work has empirically confirmed that such an effect is present in the Kepler data (See et al. 2024), the anti-correlation between close, unresolved binaries and stellar metallicity (Moe et al. 2019) could be a complicating factor. Nonetheless, based on these studies, we flagged stars with known spectroscopic  $[\text{Fe}/\text{H}] > 0.3$ , using either LAMOST or Keck/HIRES spectra (Zong et al. 2018; Petigura et al. 2022). The main limitation of this approach is that only 53.3% of the stars in Table 2 with  $t_{\text{gyro}} < 4$  Gyr had such spectra available. For the remainder of our sample, our quality flags were not sensitive to metallicity outliers. The metallicity distribution of stars for which we do have spectra suggests that at most  $\approx 5\%$  of stars in our apparently clean sample might have  $[\text{Fe}/\text{H}] > 0.3$  and not be flagged, with a slight bias toward lower metallicity stars over higher metallicity. Based on work by Claytor et al. (2020), this will be most important for stars near the Kraft break, which could have ages over-estimated by  $\approx 50\text{--}100\%$  if they are in fact metal-poor, and under-estimated in the converse case. , on page 15.

Added: *Differential rotation*: The Sun’s surface rotates once per  $\approx 25\text{--}30$  days at latitudes between the equator and  $\approx \pm 60^\circ$  (e.g. Snodgrass & Ulrich 1990). This latitudinal surface differential rotation yields two separate limitations on rotational ages. The first is on the accuracy with which any one star’s rotation period can be measured over a few-year mission like Kepler; the second is on the degree to which the intrinsic scatter in open cluster rotation sequences can be measured (e.g. Epstein & Pinsonneault 2014). In this study, we accounted for the latter intrinsic scatter based on available measurements from Ruprecht-147 (Curtis et al. 2020) and M67 data (Barnes et al. 2016; Dungee et al. 2022; Gruner et al. 2023). This means that irrespective of the assumed  $P_{\text{rot}}$  measurement precision, our model imposed a  $\approx 5\text{--}10\%$  level of intrinsic  $P_{\text{rot}}$  scatter at old ages. While this accounts for the statistical impact of differential rotation on the inferred age distribution, the accuracy issue remains; any one individual star’s observed rotation period might be faster or slower than average stars of the same mass and age. This effect can yield a  $\approx 20\%$  age error for individual Sun-like stars with ages of a few billion years; the statistical uncertainties in Table 2 should be treated with appropriate caution., on page 16.

Deleted: ~~Metallicity may also be relevant (Amard et al. 2020; See et al. 2024). — An additional caveat is that our rotation-based ages used photometric effective temperatures (Section 3), even though spectroscopic temperatures are available for the planet-hosting subset. This decision was driven by a desire for homogeneity irrespective of planet-hosting status. However it implies that the estimated planet ages might shift if one were to account for the spectroscopic information. on page 16.~~

Deleted: ~~Other age indicators may help in verifying the ages of the  $\lesssim 3$  Gyr stars that are the main focus of this work. — Specifically, chromospheric emission in the X-ray, Ca II HK line, and the UV can serve as an age tracer (Mamajek & Hillenbrand 2008; Vidotto et al. 2014; Engle 2024). However, these age indicators are all in a sense “rotation-powered”. The dynamo converts kinetic energy into magnetic energy, which is emitted through these chromospheric pathways. We did not attempt to incorporate these indicators into this study due to concerns regarding sensitivity, homogeneity, and the question of whether they in fact provide age information that is truly independent from rotation. on page 16.~~

Added: *Ages of stars older than the Sun*: , on page 16.

Deleted: ~~A broader question, beyond the scope of this work, concerns the ages of stars  $\gtrsim 4$  Gyr old. on page 16.~~

Added: *This work was limited to stars younger than 4 Gyr*, on page 16.

Added: *Planet*, on page 16.

Replaced: ~~whether the star formation history that we see in the Kepler field is universal across the thin disk.~~ replaced with: *the relative importance of star formation and kinematic heating across different components of the Galaxy.*, on page 16.

Replaced: ~~the youngest “field stars”~~ replaced with: *these young stars*, on page 16.

Deleted: ~~this work~~ on page 16.

Replaced: ~~history~~ replaced with: *rate*, on page 16.

Added: *due to its astrophysical depletion rate*, on page 17.

Replaced: ~~resulting ages~~ replaced with: *results*, on page 17.

Replaced: ~~sub-gigayear confirmed~~ replaced with: *such*, on page 17.

Added: (e.g. Kepler-1312 c and Kepler-1561 b) , on page 17.

Deleted: ~~new~~ on page 17.

Added: *Significant care is needed not only when assessing the reliability of individual rotational ages, but also when interpreting distributions of rotational ages inferred from many stars. Despite a slew of stellar quality flags, including on metallicity, evolutionary state, and binarity, up to a third of our “apparently clean” Kepler stars with  $P_{\text{rot}} < 10$  days*



could potentially be influenced by unresolved binaries (Figure 9). An additional nuance is that even after correcting for such effects, the star formation history does not directly trace the observed age distribution: particularly for pencil-beam surveys such as Kepler, kinematic heating of stars in the thin disk significantly sculpts the observed stellar (and therefore planetary) age distribution., on page 17.

Deleted: ~~main~~ on page 17.

Added: **two** , on page 17.

Added: **– what is the age distribution in the thin disk, and where are the missing young transiting exoplanets –** , on page 17.

Added: **at least** , on page 17.

Deleted: ~~The star formation rate today is  $2.81 \pm 0.12$  times lower than it was three billion years ago.~~ on page 17.

Added: **The youngest Kepler stars (0-0.3 Gyr) are at least  $2.81 \pm 0.12$  times rarer than stars 2.7-3 Gyr old. The statistical uncertainty on this value is dwarfed by the systematic uncertainty in the contamination rate of unresolved photometric binaries, which could push it as high as  $4.81 \pm 0.26$  (Section 8.2), on page 17.**

Replaced: ~~This result~~ replaced with: **This range of values**, on page 17.

Replaced: ~~from rotation-based ages broadly agrees~~ replaced with: **is broadly consistent**, on page 17.

Replaced: ~~recent reports of a declining star-formation rate~~ replaced with: **modeling expectations (Section 8.1), and with age distributions derived from**, on page 17.

Replaced: ~~, though with the~~ replaced with: **. The** , on page 17.

Added: **rotation-based ages however is their** , on page 17.

Replaced: ~~sensitivity~~ replaced with: **accuracy**, on page 17.

Replaced: ~~implies~~ replaced with: **suggests**, on page 17.

Replaced: ~~should~~ replaced with: **might**, on page 17.

Replaced: ~~derived~~ replaced with: **reported**, on page 17.

Deleted: ~~recent~~ on page 25.

Replaced: ~~an obvious~~ replaced with: **a short-period**, on page 26.

Replaced: ~~None is present in the Kepler light curve~~ replaced with: **The actual signal that is present has an amplitude of  $\approx 0.1$ - $0.3\%$  and a period of  $\approx 33$  days**, on page 26.

Deleted: ~~rotation-based~~ on page 27.

Added: **, or with rapid spin-down at Rossby numbers near 0.5 (Lu et al. 2022)**, on page 27.

Replaced: ~~2.5~~ replaced with: **2.3**, on page 27.

Replaced: ~~nine~~ replaced with: **ten**, on page 28.