10

13

## Erratum: "Kepler and the Behemoth: Three Mini-Neptunes in a 40 Million Year Old Association" (2022, AJ, 164, 215)

L. G. Bouma, <sup>1,\*</sup> R. Kerr, <sup>2</sup> J. L. Curtis, <sup>3</sup> H. Isaacson, <sup>4</sup> L. A. Hillenbrand, <sup>1</sup> A. W. Howard, <sup>1</sup> A. L. Kraus, <sup>2</sup> A. Bieryla, <sup>5</sup> D. W. Latham, <sup>5</sup> E. A. Petigura, <sup>6</sup> and D. Huber <sup>7</sup>

<sup>1</sup>Cahill Center for Astrophysics, California Institute of Technology, Pasadena, CA 91125, USA
<sup>2</sup>Department of Astronomy, The University of Texas at Austin, Austin, TX 78712, USA
<sup>3</sup>Department of Astronomy, Columbia University, 550 West 120th Street, New York, NY 10027, USA
<sup>4</sup>Astronomy Department, University of California, Berkeley, CA 94720, USA
<sup>5</sup>Center for Astrophysics | Harvard & Smithsonian, 60 Garden St, Cambridge, MA 02138, USA
<sup>6</sup>Department of Physics & Astronomy, University of California Los Angeles, Los Angeles, CA 90095, USA
<sup>7</sup>Institute for Astronomy, University of Hawai'i, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

## Submitted to Astronomical Journal

In the original manuscript, the top-right panel of Figure 2 erroneously omitted KOI-7913 B due to an error in the plotting script. Figure 1 in this erratum corrects the omission.

Corresponding author: L. G. Bouma

luke@astro.caltech.edu




Figure 1. Age-diagnostic diagrams from the stellar groups near Kepler-1643, KOI-7368, and KOI-7913. *Top row*: Color-absolute magnitude diagram of candidate Cep-Her members, plotted over candidate members of the  $\delta$  Lyr cluster ( $\approx$  38 Myr; Bouma et al. 2022) and the Gaia EDR3 Catalog of Nearby Stars (gray background). The left and right columns shows stars in RSG-5 and CH-2, respectively. The range of colors is truncated to emphasize the pre-main-sequence; approximate spectral types are shown on the upper axes. Stars that fall far below the cluster sequences are field interlopers. *Bottom row*: TESS and ZTF-derived stellar rotation periods, with the Pleiades ( $\approx$  112 Myr) and Praesepe ( $\approx$  650 Myr) shown for reference (Rebull et al. 2016; Douglas et al. 2017). The detection efficiency for reliable rotation periods falls off beyond ( $G_{\rm BP}-G_{\rm RP}$ )<sub>0</sub>  $\gtrsim$  2.6.

## REFERENCES

Bouma, L. G., Curtis, J. L., Masuda, K., et al. 2022, AJ, 163, 121 16 Douglas, S. T., Agüeros, M. A., Covey, K. R., & Kraus, A. 2017,

17 ApJ, 842, 83

18 Rebull, L. M., Stauffer, J. R., Bouvier, J., et al. 2016, AJ, 152, 113