

A 38 Million Year Old Neptune-Sized Planet in the Kepler Field

² L. G. BOUMA,^{1,2,*} J. L. CURTIS,^{3,4} K. MASUDA,⁵ L. A. HILLENBRAND,¹ G. STEFANSSON,^{2,†} H. ISAACSON,⁶
³ N. NARITA,^{7,8,9,10} A. FUKUI,^{7,10} M. IKOMA,¹¹ M. TAMURA,^{12,9,11} A. L. KRAUS,¹³ E. FURLAN,¹⁴
⁴ C. L. GNILKA,^{15,14} K. V. LESTER,¹⁵ AND S. B. HOWELL¹⁵

⁵ Cahill Center for Astrophysics, California Institute of Technology, Pasadena, CA 91125, USA

⁶ Department of Astrophysical Sciences, Princeton University, 4 Ivy Lane, Princeton, NJ 08540, USA

⁷ Department of Astronomy, Columbia University, 550 West 120th Street, New York, NY 10027, USA

⁸ Department of Astrophysics, American Museum of Natural History, New York, NY 10024, USA

⁹ Department of Earth and Space Science, Osaka University, Osaka 560-0043, Japan

¹⁰ Astronomy Department, University of California, Berkeley, CA 94720, USA

¹¹ Komaba Institute for Science, The University of Tokyo, Tokyo 153-8902, Japan

¹² Japan Science and Technology Agency, PRESTO, Tokyo 153-8902, Japan

¹³ Astrobiology Center, Tokyo 181-8588, Japan

¹⁴ Instituto de Astrofísica de Canarias (IAC), 38205 La Laguna, Tenerife, Spain

¹⁵ Division of Science, National Astronomical Observatory of Japan, Tokyo 181-8588, Japan

¹⁶ Department of Astronomy, The University of Tokyo, Tokyo 113-0033, Japan

¹⁷ Department of Astronomy, The University of Texas at Austin, Austin, TX 78712, USA

¹⁸ NASA Exoplanet Science Institute, Caltech/IPAC, Pasadena, CA 91125, USA

¹⁹ NASA Ames Research Center, Moffett Field, CA 94035, USA

(Received 2021 Oct 12; Revised —; Accepted —)

Submitted to AAS Journals

ABSTRACT

Kepler 1627A is a G8V star previously known to host a $3.8 R_{\oplus}$ (**Replaced:** mini-Neptune replaced with: planet) on a 7.2 day orbit. The star was observed by the Kepler space telescope because it is nearby ($d = 329$ pc) and it resembles the Sun. Here we show using Gaia kinematics, TESS stellar rotation periods, and spectroscopic lithium abundances that Kepler 1627 is a member of the 38^{+6}_{-5} Myr old δ Lyr cluster. To our knowledge, this makes Kepler 1627Ab the youngest planet with a precise age yet found by the (**Replaced:** main replaced with: prime) Kepler mission. The Kepler photometry shows two peculiarities: the average transit profile is asymmetric, and the individual transit times might be correlated with the local light curve slope. We discuss possible explanations for each anomaly. More importantly, the δ Lyr cluster is one of $\sim 10^3$ coeval groups whose properties have been clarified by Gaia. Many other exoplanet hosts are candidate members of these clusters; their ages can be verified (**Replaced:** through replaced with: with) the trifecta of Gaia, TESS, and ground-based spectroscopy.

Keywords: exoplanet evolution (491), open star clusters (1160), stellar ages (1581)

1. INTRODUCTION

While thousands of exoplanets have been discovered orbiting nearby stars, the vast majority of

them are several billion years old. This makes it difficult to test origin theories for the different families of planets, since many evolutionary processes are expected to operate on timescales of less than 100 million years.

For instance, the “mini-Neptunes”, thought to be made of metal cores, silicate mantles (Kite et al. 2020), and extended hydrogen-dominated atmospheres, are expected to shrink in size by fac-

Corresponding author: L. G. Bouma
luke@astro.caltech.edu

* 51 Pegasi b Fellow

† Henry Norris Russell Fellow

tors of several over their first 10^8 years. Specifically, in the models of Owen & Wu (2016) and Owen (2020), the $\approx 5 M_\oplus$ planets start with sizes of $4 - 12 R_\oplus$ shortly after the time of disk dispersal ($\lesssim 10^7$ years), and shrink to sizes of $2 - 4 R_\oplus$ by 10^8 years. While the majority of this change is expected to occur within the first few million years after the disk disperses (Ikoma & Hori 2012), stellar irradiation and internal heat can also power gradual outflows which, if strong enough, can deplete or entirely strip the atmosphere (Lopez et al. 2012; Owen & Wu 2013; Ginzburg et al. 2018). Discovering young planets, measuring their masses, and detecting their atmospheric outflows are key steps toward testing this paradigm, which is often invoked to explain the observed radius distribution of mature exoplanets (Fulton et al. 2017; Van Eylen et al. 2018).

The K2 and TESS missions have now enabled the detection of about ten close-in planets younger than 100 million years, all smaller than Jupiter (Mann et al. 2016; David et al. 2016, 2019; Newton et al. 2019; Bouma et al. 2020; Plavchan et al. 2020; Rizzuto et al. 2020; Martioli et al. 2021). The Kepler mission however has not yielded any planets with precise ages below one gigayear (Meibom et al. 2013). The reason is that during the (Replaced: main replaced with: prime) Kepler mission (2009–2013), only four open clusters were known in the Kepler field, with ages spanning 0.7 Gyr to 9 Gyr (Meibom et al. 2011). Though isochronal, gyrochronal, and lithium-based analyses suggest that younger Kepler planets do exist (Walkowicz & Basri 2013; Berger et al. 2018; David et al. 2021), accurate and precise age measurements typically require an ensemble of stars. Fortunately, recent analyses of the Gaia data have greatly expanded our knowledge of cluster memberships (e.g., Cantat-Gaudin et al. 2018; Zari et al. 2018; Kounkel & Covey 2019; Meingast et al. 2021; Kerr et al. 2021). As part of our Cluster Difference Imaging Photometric Survey (CDIPS, Bouma et al. 2019), we concatenated the available analyses from the literature, which yielded a list of candidate young and age-dated stars (see Appendix A).

Matching our young star list against stars observed by Kepler revealed that Kepler observed a portion of the δ Lyr cluster (Stephenson-1; Theia 73). More specifically, a clustering analysis of the Gaia data by Kounkel & Covey (2019) reported that Kepler 1627 (KIC 6184894; KOI 5245) is a δ Lyr cluster member. Given the previous statistical validation of the close-in (Replaced: mini-Neptune replaced with: Neptune-sized planet) Kepler 1627b (Tenenbaum et al. 2012; Morton et al. 2016; Thompson et al. 2018), we begin

by scrutinizing the properties of the cluster (Section 2). We find that the δ Lyr cluster is 38^{+6}_{-5} Myr old, and in Section 3 show that Kepler 1627 is both a binary and also a member of the cluster. Focusing on the planet (Section 4), we confirm that despite the existence of the previously unreported M2.5V companion, hereafter Kepler 1627B, the planet orbits the G-dwarf primary, Kepler 1627A. We also analyze an asymmetry in the average transit profile, and a possible correlation between the individual transit times and the local light curve slope. We conclude by discussing broader implications for our ability to age-date a larger sample of planets (Section 5).

2. THE CLUSTER

To measure the age of the δ Lyr cluster, we first selected a set of candidate cluster members (Section 2.1), and then analyzed these stars using a combination of the isochronal and gyrochronal techniques (Section 2.2).

2.1. Selecting Cluster Members

Kounkel & Covey (2019) applied an unsupervised clustering algorithm to Gaia DR2 on-sky positions, proper motions, and parallaxes for stars within the nearest kiloparsec. For the δ Lyr cluster (Theia 73), they reported 3,071 candidate members. We matched these stars against the latest Gaia EDR3 observations using the dr2_neighbourhood table from the ESA archive, taking the stars closest in proper motion and epoch-corrected angular distance as the presumed match (Gaia Collaboration et al. 2021a). In Figure 1, (Added: we)have shown galactic positions only for the stars with parallax signal-to-noise exceeding 20. (Added: We calculated the tangential velocities for each of these stars relative to Kepler 1627 ($\Delta\mu^*$) by subtracting the observed proper motion from what the proper motion at the star’s position would be if it were co-moving with Kepler 1627.)The reported cluster members (gray and black points) extend over a much larger volume (Added: in both physical and kinematic space)than the cluster previously identified by Stephenson (1959) and later corroborated by Eggen (1968). While the non-uniform “clumps” of stars might comprise a *bona fide* cluster of identically-aged stars, they could also be heavily contaminated by field stars. (Added: One reason to suspect this is that the spread in tangential velocities exceeds typical limits for kinematic coherence (Meingast et al. 2021)). We therefore considered stars only in the immediate kinematic and spatial vicinity of Kepler 1627 as candidate cluster members. We performed (Replaced: the replaced with: this) selection cut(Deleted: s) manually, by drawing lassos with




Figure 1. Galactic positions and tangential velocities of stars in the δ Lyr cluster. Points are reported cluster members from Kounkel & Covey (2019). The tangential velocities relative to Kepler 1627 (bottom right) are computed assuming that every star has the same three-dimensional spatial velocity as Kepler 1627. Our analysis considers stars (black points) in the spatial and kinematic vicinity of Kepler 1627 (yellow star). The question of whether the other candidate cluster members (gray points) are part of the cluster is outside our scope. The location of the Sun is (\odot) is shown (Added: to clarify the direction along which parallax uncertainties are expected to produce erroneous clusters).




Figure 2. Kepler's view of the δ Lyr cluster, shown in equatorial coordinates. Each black circle is a candidate cluster member selected based on its position and kinematics (Figure 1). Of the 1,201 candidate cluster members, 58 have at least one quarter of Kepler data. TESS has also observed most of the cluster, for one to two lunar months to date.

the interactive `glue` visualization tool (Beaumont et al. 2014) in the four projections shown in Figure 1. The overlap between the Kepler field and the resulting candidate cluster members is shown in Figure 2. While this method will include some field interlopers in the “cluster star” sample, and vice-versa, it should suffice for our aim of verifying the existence of the cluster in the vicinity of Kepler 1627.

2.2. The Cluster’s Age

2.2.1. Color-Absolute Magnitude Diagram

We measured the isochrone age using an empirical approach. The left panel of Figure 3 shows the color-absolute magnitude diagram (CAMD) of candidate δ Lyr cluster members, IC 2602, the Pleiades, and the field. The stars from the Pleiades and IC 2602 were adopted from Cantat-Gaudin et al. (2018), and the field stars are from the Gaia EDR3 Catalog of Nearby Stars (Gaia Collaboration et al. 2021b). We cleaned these following the data filtering criteria from Gaia Collaboration et al. (2018a, Appendix B), except that we weakened the parallax precision requirement to $\varpi/\sigma_\varpi > 5$. (Added: This also involved cuts on the photometric signal to noise ratio, the number of visibility periods used, the astrometric χ^2 of the single-source solution, and the $G_{\text{BP}} - G_{\text{RP}}$ color excess factor.) These filters were designed to include genuine binaries while omitting instrumental artifacts. We then corrected for extinction by querying the 3-dimensional maps of Capitanio et al. (2017) and Lallement et al. (2018)¹, and applied the extinction coefficients $k_X \equiv A_X/A_0$ computed by Gaia Collaboration et al. (2018a) assuming that $A_0 = 3.1E(B-V)$. For IC 2602, the Pleiades, and the δ Lyr cluster, this procedure yielded a respective mean and standard deviation for the reddening of $E(B-V) = \{0.020 \pm 0.003, 0.045 \pm 0.008, 0.032 \pm 0.006\}$. These values (Replaced: agree reasonably well with replaced with: are within a factor of two of) previously reported values (Replaced: from replaced with: in) the literature (e.g., Gaia Collaboration et al. 2018a; Kounkel & Covey 2019; Bossini et al.

¹ <https://stilism.obspm.fr/>, 2021/09/25




Figure 3. The δ Lyr cluster is 38^{+6}_{-5} Myr old. *Top:* Color-absolute magnitude diagram of candidate δ Lyr cluster members, in addition to stars in IC 2602 (≈ 38 Myr), the Pleiades (≈ 115 Myr), and the Gaia EDR3 Catalog of Nearby Stars (gray background). The zoomed right panel highlights the pre-main-sequence. The δ Lyr cluster and IC 2602 are approximately the same isochronal age. *Bottom:* TESS and Kepler stellar rotation period versus dereddened Gaia color, with the Pleiades and Praesepe (650 Myr) shown for reference (Rebull et al. 2016; Douglas et al. 2017). Most candidate δ Lyr cluster members are gyrochronologically younger than the Pleiades; outliers are probably field interlopers.

204 2019)(Added: , and are all small enough that
 205 the choice of whether to use them *vs.* other ex-
 206 tinction estimates does not affect our primary
 207 conclusions).

208 Figure 3 shows that the δ Lyr cluster and IC 2602
 209 overlap, and therefore are approximately the same
 210 age. In our exploration, we also compared against
 211 μ -Tau (62 ± 7 Myr; Gagné et al. 2020) and the
 212 Upper-Centaurus-Lupus (UCL) component of the
 213 Sco OB2 association (≈ 16 Myr; Pecaut & Mama-
 214 jek 2016). The pre-main-sequence M dwarfs of the
 215 δ Lyr cluster were intermediate between the lat-
 216 ter two clusters. To turn this heuristic interpola-
 217 tion into an age measurement, we used the empir-
 218 ical method developed by Gagné et al. (2020). In
 219 brief, we fitted the pre-main-sequence loci of a set
 220 of reference clusters, and the locus of the target δ
 221 Lyr cluster was then modeled as a piecewise linear
 222 combination of these reference clusters. For our
 223 reference clusters, we adopted members of UCL,
 224 IC 2602, and the Pleiades from Damiani et al.
 225 (2019) and Cantat-Gaudin et al. (2018) respec-
 226 tively. We removed binaries by requiring RUWE <
 227 1.3, radial_velocity_error below the 80th
 228 percentile of each cluster’s distribution, and ex-
 229 cluded stars that were obvious photometric bina-
 230 ries in the CAMD.² We then passed a moving box
 231 average and standard deviation across the CAMD
 232 in 0.10 mag bins, fitted a univariate spline to the
 233 binned values, and assembled a piecewise grid
 234 of (Deleted: hybrid–) isochrones spanning the
 235 ages between UCL (Replaced: to replaced with:
 236 and) the Pleiades using Equation(Replaced: s 6
 237 and 7 replaced with: 6) from Gagné et al. (2020).
 238 (Added: To derive a probability distribution
 239 function for the age of δ Lyr cluster, we then
 240 assumed a Gaussian likelihood that treated the
 241 interpolated isochrones as the “model” and the
 242 δ Lyr cluster’s isochrone as the “data” (Equation
 243 7 from Gagné et al. 2020). The cluster’s age
 244 and its statistical uncertainty are then quoted as
 245 the the mean and standard deviation of this age
 246 posterior.)

247 The ages returned by this procedure depend on
 248 the ages assumed for each reference cluster. We
 249 adopted a 115 Myr age for the Pleiades (Dahm
 250 2015), and a 16 Myr age for UCL (Pecaut & Ma-
 251 majek 2016). The age of IC 2602 however is the
 252 most important ingredient, since it receives the
 253 most weight in the interpolation. Plausible ages
 254 for IC 2602 span 30 Myr to 46 Myr, with older ages
 255 being preferred by the lithium-depletion-boundary
 256 (LDB) measurements (Dobbie et al. 2010; Randich

² For a description of the renormalized unit weight error (RUWE), see the GAIA DPAC technical note http://www.rssd.esa.int/doc_fetch.php?id=3757412.

257 et al. 2018) and younger ages by the main-
 258 sequence turn-off (Stauffer et al. 1997; David &
 259 Hillenbrand 2015; Bossini et al. 2019). If we were
 260 to adopt the 30 Myr age for IC 2602, then the δ
 261 Lyr cluster would be 31^{+5}_{-4} Myr old. For the con-
 262 verse extreme of 46 Myr, the δ Lyr cluster would
 263 be 44^{+8}_{-7} Myr old. We adopt an intermediate 38 Myr
 264 age for IC 2602, which yields an age for the δ Lyr
 265 cluster of 38^{+6}_{-5} Myr.³ Follow-up studies of the LDB
 266 or main-sequence turn-off in the δ Lyr cluster could
 267 help determine a more precise and accurate age for
 268 the cluster, and are left for future work.

269 2.2.2. Stellar Rotation Periods

270 Of the 3,071 candidate δ Lyr cluster members re-
 271 ported by Kounkel & Covey (2019), 924 stars were
 272 amenable to rotation period measurements ($G < 17$
 273 and $(G_{\text{BP}} - G_{\text{RP}})_0 > 0.5$) using the TESS full frame
 274 image data. (Added: As a matter of scope, we re-
 275 stricted our attention to the 391 stars discussed
 276 in Section 2.1 in the spatial and kinematic prox-
 277 imity of Kepler 1627.) We extracted light curves
 278 from the TESS images using the nearest pixel to
 279 each star, and regressed them against systematics
 280 with the causal pixel model implemented in the
 281 unpopular package (Hattori et al. 2021). We
 282 then measured candidate rotation periods using a
 283 Lomb-Scargle periodogram (Lomb 1976; Scargle
 284 1982; Astropy Collaboration et al. 2018). To en-
 285 able cuts on crowding, we queried the Gaia source
 286 catalog for stars within a 21.⁰ radius of the tar-
 287 get star (a radius of 1 TESS pixel). Within this ra-
 288 dius, we recorded the number of stars with greater
 289 brightness than the target star, and with brightness
 290 within 1.25 TESS magnitudes of the target star.

291 We then cleaned the candidate TESS rotation pe-
 292 riod measurements through a combination of auto-
 293 mated and manual steps. (Deleted: As a matter
 294 of scope, we restricted our attention to the 391
 295 stars discussed in Section 2.1 in the spatial
 296 and kinematic proximity of Kepler 1627.)
 297 (Deleted: Kepler rotation periods were derived
 298 by McQuillan et al. (2014) for 28 of our 391
 299 stars; for these cases, we simply adopted the
 300 Kepler rotation period.) (Added: First, to val-
 301 idate the TESS rotation periods, we compared
 302 against 28 stars from McQuillan et al. (2014)
 303 that were also observed by Kepler. Of the 23
 304 stars with Kepler periods below 10 days, 21 of
 305 the TESS periods agreed with the Kepler ro-
 306 tation periods; the other 2 were measured at

³ Our exploration of the PARSEC and MIST isochrone mod-
 els over a grid of ages, metallicities, and reddenings, yielded
 the best agreement for this ≈ 38 Myr age as well, given
 $[{\text{Fe}}/{\text{H}}] = +0.1$ and $A_V = 0.2$ (Bressan et al. 2012; Choi et al. 2016); this preferred CAMD reddening is higher than the Lallement et al. (2018) value by a factor of two.

the double-period harmonic. Of the remaining 5 stars with Kepler rotation periods above 10 days, none were correctly recovered by TESS (3 were near the half-period harmonic). We therefore adopted a general stance of caution for TESS rotation periods exceeding 10 days, and used the Kepler rotation periods for the stars for which they were available.) For the remaining stars with only TESS data, we focused only on the stars for which no companions were known with a brightness exceeding one-tenth of the target star in a $21.^{\circ}0$ radius. There were 192 stars that met these crowding requirements, and that had TESS data available. For plotting purposes we then imposed a selection based on the strength of the signal itself: we required the Lomb Scargle power to exceed 0.2, and the period to be below 15 days.

The lower panel of Figure 3 shows the resulting 145 stars. The majority of these stars fall below the “slow sequence” of the Pleiades, consistent with a gyrochronal age for the δ Lyr cluster below 100 Myr. In fact, the rotation-color distributions of other 30 Myr to 50 Myr clusters (e.g., IC 2602 and IC 2391) are indistinguishable(Added: from each other) (Douglas et al. 2021). Approximately 10 of the δ Lyr cluster stars appear as outliers above the “slow sequence”. Assuming that they are all false positives (*i.e.*, field interlopers), our rotation period detection fraction would be $135/192 \approx 70\%$. (Replaced: The other stars are likely to be field contaminants. replaced with: Although some of these outlier stars might be unresolved F+K binaries that are in the cluster (Stauffer et al. 2016), assuming that they are field contaminants provides a more secure lower bound of the rotation period detection fraction.) A final possible confounding factor – binarity – is known to affect the “fast sequence” of stars beneath the slow sequence (Meibom et al. 2007; Gillen et al. 2020; Bouma et al. 2021). We do not expect it to change the central conclusion regarding the cluster’s age.

3. THE STARS

3.1. Kepler 1627A

3.1.1. Age

Based on the spatial and kinematic association of Kepler 1627 with the δ Lyr cluster, and the assumption that the planet formed shortly after the star, it seems likely that Kepler 1627 is the same age as the cluster. There are two consistency checks on whether this is true: rotation and lithium. Based on the Kepler light curve, the rotation period is 2.642 ± 0.042 days, where the quoted uncertainty is based on the scatter in rotation periods measured from each individual Kepler quarter. This is consistent with comparable cluster members (Figure 3).

To infer the amount of Li I from the 6708 Å doublet (e.g., Soderblom et al. 2014), we acquired an iodine-free spectrum from Keck/HIRES on the night of 2021 March 26 using the standard setup and reduction techniques of the California Planet Survey (Howard et al. 2010). Following the equivalent width measurement procedure described by Bouma et al. (2021), we find $\text{EW}_{\text{Li}} = 233^{+5}_{-7}$ mÅ. This value does not correct for the Fe I blend at 6707.44Å. Nonetheless, given the stellar effective temperature (Table 1), this measurement is in agreement with expectations for a ≈ 40 Myr star (e.g., as measured in IC 2602 by Randich et al. 2018). It is also larger than any lithium equivalent widths measured by Berger et al. (2018) in their analysis of 1,301 Kepler-star spectra.

3.1.2. Stellar Properties

The adopted stellar parameters are listed in Table 1. The stellar mass, radius, and effective temperature are found by interpolating against a 38 Myr MIST isochrone (Choi et al. 2016). The statistical uncertainties are propagated from the absolute magnitude (mostly originating from the parallax uncertainty) and the color; the systematic uncertainties are taken to be the difference between the PARSEC (Bressan et al. 2012) and MIST isochrones. Reported uncertainties are a quadrature sum of the statistical and systematic components. As a consistency check, we analyzed the aforementioned Keck/HIRES spectrum from the night of 2021 March 26 using a combination of SpecMatch-Emp for stellar properties, and SpecMatch-Synth for $v\sin i$ (Yee et al. 2017). This procedure yielded $T_{\text{eff}} = 5498 \pm 100$ K, $\log g = 4.6 \pm 0.1$, $[\text{Fe}/\text{H}] = 0.15 \pm 0.10$ from SpecMatch-Emp, and $v\sin i = 18.9 \pm 1.0$ from SpecMatch-Synth. These values are within the $1-\sigma$ uncertainties of our adopted values from the isochrone interpolation.

3.2. Kepler 1627B

We first noted the presence of a close neighbor in the Kepler 1627 system on 2015 July 22 when we acquired adaptive optics imaging using the NIRC2 imager on Keck-II. We used the narrow camera ($\text{FOV} = 10.2''$) to obtain 8 images in the K' filter ($\lambda = 2.12 \mu\text{m}$) with a total exposure time of 160 s. We analyzed these data following Kraus et al. (2016), which entailed using PSF-fitting to measure the separation, position angle, and contrast of the candidate companion. The best-fitting empirical PSF template was identified from among the near-contemporaneous observations of single stars in the same filter. The mean values inferred from the 8 images are reported in Table 1. To estimate the detection limits, we analyzed the residuals after subtracting the empirical PSF template.




Figure 4. Kepler 1627 is a binary. *Left:* High-resolution imaging from Gemini-North/‘Alopeke and Keck/NIRC2 shows an \approx M2.5V companion at $\rho \approx 0.^{\prime\prime}16$, which corresponds to a projected separation of 53 ± 4 AU. The inset shows a cutout of the stacked NIRC2 image (North is up, East is left, scale is set by the separation of the binary). The lines show $5-\sigma$ contrast limits for the ‘Alopeke filters, and $6-\sigma$ contrast limits for NIRC2 outside of $0.^{\prime\prime}15$. *Right:* Gaia EDR3 renormalized unit weight error (RUWE) point estimates for candidate δ Lyr cluster members. Since other members of the cluster with similar brightnesses have comparable degrees of photometric variability, the high RUWE independently suggests that Kepler 1627 is a binary.

Within each residual image, the flux was measured through 40 mas apertures centered on every pixel, and then the noise as a function of radius was estimated from the RMS within concentric rings. Finally, the detection limits were estimated from the strehl-weighted sum of the detection significances in the image stack, and we adopted the $6-\sigma$ threshold as the detection limit for ruling out additional companions.

We also observed Kepler 1627 on Gemini-North using the ‘Alopeke speckle imager on 2021 June 24. ‘Alopeke is a dual-channel speckle interferometer that uses narrow-band filters centered at $0.83 \mu\text{m}$ and $0.56 \mu\text{m}$. We acquired three sets of 1000×60 msec exposures during good seeing ($0.45''$), and used the autocorrelation function of these images to reconstruct a single image and $5-\sigma$ detection limits (see Howell et al. 2011). This procedure yielded a detection of the companion in the $0.83 \mu\text{m}$ notch filter, but not the $0.56 \mu\text{m}$ filter. The measured projected separation and magnitude difference are given in Table 1.

Figure 4 summarizes the results of the high-resolution imaging. The Gaia EDR3 parallax for the primary implies a projected separation of 53 ± 4 AU, assuming the companion is bound. Although the companion is unresolved in the Gaia source catalog (there are no comoving, codistant candidate companions brighter than $G < 20.5$ mag within $\rho < 120''$), its existence was also suggested by the primary star’s large (Replaced: renormalized unit weight error (RUWE), replaced with: RUWE) relative to other members of the δ Lyr cluster (Added: (RUWE ≈ 2.9 ; roughly the 98th percentile of the cluster’s distribution)). Based

on the apparent separation, the binary orbital period is of order hundreds of years. The large RUWE is therefore more likely to be caused by a PSF-mismatch skewing the Gaia centroiding during successive scans, rather than true astrometric motion. Regardless, given the low geometric probability that a companion imaged at $\rho \approx 0.^{\prime\prime}16$ is a chance line-of-sight companion, we proceed under the assumption that the companion is bound, and that Kepler 1627 is a binary. Given the distance and age, the models of Baraffe et al. (2015) imply a companion mass of $M_B \approx 0.33 M_{\odot}$ and companion temperature of $T_{\text{eff},B} \approx 3450$ K. The corresponding spectral type is roughly M2.5V (Pecaut & Mamajek 2013). These models combined with the NIRC2 contrast limits imply physical limits on tertiary companions of $M_{\text{ter}} < 50 M_{\text{Jup}}$ at $\rho = 50$ AU, $M_{\text{ter}} < 20 M_{\text{Jup}}$ at $\rho = 100$ AU, and $M_{\text{ter}} < 10 M_{\text{Jup}}$ at $\rho = 330$ AU.

4. THE PLANET

4.1. Kepler Light Curve

The Kepler space telescope observed Kepler 1627 at a 30-minute cadence from 2009 May 2 until 2013 April 8. Data gaps during quarters 4, 9, and 13 led to an average duty cycle over the 3.9 year interval of 67%. Kepler 1627 was also observed at 1-minute cadence from 2012 Oct 5 until 2013 Jan 11. The top panel of Figure 5 shows a portion of the 30-minute cadence PDCSAP light curve. Nonastrophysical variability has been removed using the methods discussed by Smith et al. (2017); the default optimal aperture was assumed (Smith et al. 2016). Cadences with non-zero quality flags (9% of the data) have been omitted. The resulting ph-




Figure 5. The light curve of Kepler 1627. *Top:* The Kepler data span 1,437 days (3.9 years), sampled at 30 minute cadence; a 100 day segment is shown. The top panel shows the PDCSAP median-subtracted flux in units of parts-per-thousand ($\times 10^{-3}$). The dominant signal is induced by starspots. The stellar variability model (orange line) is subtracted below, revealing the transits of Kepler 1627Ab. The online Figure Set spans the entire 3.9 years of observations. *Bottom:* Phase-folded transit of Kepler 1627Ab with stellar variability removed. Windows over 20 hours (*left*) and the entire orbit (*right*) are shown, and the residual after subtracting the transit is in the bottom-most row. The 2- σ model uncertainties and the best-fit model are the light purple band and the dark purple line. Gray points are individual flux measurements; black points bin these to 15 minute intervals, and have a representative 1- σ error bar in the lower right of each panel. The asymmetric residual during transit is larger than the out-of-transit scatter.

tometry is dominated by a quasi-periodic starspot signal with a peak-to-peak amplitude that varies between 2% and 8%. (Added: Given that the secondary companion’s brightness in the Kepler band is $\approx 1.5\%$ that of the primary, source confusion for the rotation signal is not expected to be an issue.) Previous analyses have identified and characterized the smaller transit signal (Tenenbaum et al. 2012; Thompson et al. 2018), validated its planetary nature (Morton et al. 2016), and even searched the system for transit timing variations (Holczer et al. 2016). Nonetheless, since the cluster membership provides us with more precise stellar parameters than those previously available, we opted to reanalyze the light curve.

4.1.1. Transit and Stellar Variability Model

We fitted the Kepler long cadence time series with a model that simultaneously included the planetary transit and the stellar variability. The stellar variability was modeled with the `RotationTerm` Gaussian Process kernel in `exoplanet` (Foreman-Mackey et al. 2020). This kernel assumes that the variability is generated by a mixture of two damped simple harmonic oscillators with characteristic frequencies set by $1/P_{\text{rot}}$ and its first harmonic. We additionally included a jitter term to inflate the flux uncertainties in a manner that accounted for otherwise unmodeled excess white noise, and let the eccentricity float. For the limb-darkening, we assumed a quadratic law, and sampled using the uninformative prior suggested by Kipping (2013).

Our model therefore included 10 free parameters for the transit ($\{P, t_0, \delta, b, u_1, u_2, R_*, \log g, e, \omega\}$), 2 free parameters for the light curve normalization and a white noise jitter ($\{\langle f \rangle, \sigma_f\}$), and 5 hyperparameters for the GP ($\{\sigma_{\text{rot}}, P_{\text{rot}}, Q_0, dQ, f\}$). We also considered including an additive `SHOTerm` kernel to account for stochastic noise, but found that this did not affect the results, and so opted for the simpler GP kernel. We fitted the models using `PyMC3` (Salvatier et al. 2016; Theano Development Team 2016), and accounted for the finite integration time of each exposure in the numerical integration when evaluating the model light curve (see Kipping 2010). We assumed a Gaussian likelihood, and after initializing each model with the parameters of the maximum *a posteriori* model, we sampled using `PyMC3`’s gradient-based No-U-Turn Sampler (Hoffman & Gelman 2014) in the bases indicated in Table 2. We used \hat{R} as our convergence diagnostic (Gelman & Rubin 1992).

Figure 5 shows the resulting best-fit model in orange (top) and purple (bottom). The model parameters and their uncertainties, given in Table 2, are broadly consistent with a (Replaced:

543 **mini-Neptune** replaced with: **Neptune**)-sized
 544 planet ($3.82 \pm 0.16 R_{\oplus}$) on a close-in circular⁴
 545 orbit around a G8V host star ($0.88 \pm 0.02 R_{\odot}$). This
 546 best-fit planet size is consistent with those previ-
 547 ously reported by Morton et al. (2016) and Berger
 548 et al. (2018), and corrects for the small amount of
 549 flux dilution from Kepler 1627B.

4.1.2. Transit Asymmetry

The transit fit however is not perfect: the lower panels of Figure 5 show an asymmetric residual in the data relative to the model: the measured flux is high during the first half of transit, and low in the second half. The semi-amplitude of this deviation is ≈ 50 ppm, which represents a $\approx 3\%$ distortion of the transit depth ($\delta = 1759 \pm 62$ ppm). Note that although this asymmetry is within the $2-\sigma$ model uncertainties, the model has a jitter term that grows to account for excess white noise in the flux. The significance of the asymmetry is therefore best assessed in comparison against the intrinsic out-of-transit scatter in the data (≈ 16 ppm), not the model uncertainties. The lower right panel of Figure 5 demonstrates that the scatter during transit is higher than during all other phases of the planet’s orbit.

To determine whether the asymmetry could be a systematic caused by our stellar variability model, we explored an alternative approach in which we isolated each transit window, locally fitted out polynomial trends, and then binned all the observed transits; the asymmetry was still present at a comparable amplitude. Appendix B describes a more detailed analysis, which finds that the asymmetry also seems to be robust to different methods of data binning in time and by local light curve slope. Possible astrophysical explanations are discussed in Section 5.

4.1.3. Transit Timing and the Local Slope

The previous analysis by Holczer et al. (2016) did not find any significant long-term transit timing or duration variations (TTVs or TDVs) for Kepler 1627. Quantitatively, the mean and standard deviation of the TTVs and TDVs they measured were -1.1 ± 13.8 min and -3.3 ± 22.1 min. In an earlier analysis however, Holczer et al. (2015) studied correlations between TTVs and local light curve slopes, and for Kepler 1627 found a weak correlation of -29 ± 13 min day $^{-1}$ between the two quantities. Given the possible connection between such correlations and the unresolved starspot crossings that we expect to be present in the Kepler 1627

⁴ Our transit fitting yields $e < 0.48$ at $2-\sigma$; the constraints on the eccentricity are not particularly strong.




Figure 6. Possible evidence for a prograde orbit of Kepler 1627 Ab. The time of each Kepler transit was measured, along with the local slope of the light curve. The two quantities are weakly anti-correlated ($\approx 2\sigma$), which might be caused by starspot crossings during the first (second) half of transit inducing a positive (negative) TTV, provided that the orbit is prograde (Mazeh et al. 2015). The units along the abscissa can be understood by considering that the stellar flux changes by ~ 60 ppt per half rotation period (~ 1.3 days).

594 light curve (Mazeh et al. 2015), we opted to re-
595 examine the individual transit times.

596 We therefore isolated each of the 144 observed
597 transits to within ± 4.5 hr of each transit, and fitted
598 each window with both *i*) a local (Added: sec-
599 ond or fourth-order) polynomial baseline plus
600 the transit, and *ii*) a local linear trend(Added:
601 plus the transit). (Deleted: We considered
602 the results both for a second and fourth-order
603 time-dependence in the local baseline.) We let
604 the mid-time of each transit float, and then calcu-
605 lated the residual between the measured mid-time
606 and that of a periodic orbit. This residual, the
607 transit timing variation, is plotted in Figure 6 against
608 the local linear slope for the fourth-order polyno-
609 mial baseline. The slope of -21 ± 10 min day $^{-1}$ is
610 similar to that found by Holczer et al. (2015).

611 One concern we had in this analysis was whether
612 our transit fitting procedure might induce spuri-
613 ous correlations between the slope and transit time.
614 In particular, using the second-order polynomial
615 baseline yielded a larger anti-correlation between
616 the TTVs and local slopes, of -79 ± 14 min day $^{-1}$.
617 We therefore performed an injection-recovery pro-
618 cedure in which we injected transits at differ-
619 ent phases in the Kepler 1627 light curve and re-
620 peated the TTV analysis. This was done at ≈ 50
621 phases, each separated by $0.02 P_{\text{orb}}$ while omitting
622 the phases in transit. For the second-order poly-
623 nomial baseline, this procedure yielded a similar anti-
624 correlation in the injected transits as that present in
625 the real transit; using this baseline would therefore




Figure 7. Keck/HIRES radial velocities and MuS-CAT3 photometry from the transit of 2021 Aug 7. *Top:* The radial velocity jitter across the 15 minute exposures ($\sigma_{\text{RV}} \approx 30 \text{ m s}^{-1}$) prevented us from detecting the RM effect; a model including the RM anomaly and a quadratic trend in time to fit the spot-induced $\approx 250 \text{ m s}^{-1}$ trend is shown (see Appendix C). Shaded bands show 2σ model uncertainties. *Middle:* The RV variations are strongly correlated with varying emission in the Ca H and K lines. *Bottom:* The photometric transit depths are consistent across the *griz* bandpasses. The photometry is binned at 10 minute intervals.

626 bias the result. However for the fourth-order base-
627 line, the correlation present in the data was stronger
628 than in all but one of the injected transits. Possi-
629 ble interpretations are discussed below. Given the
630 (Added: lack of) statistical significance, this anal-
631 ysis should be interpreted as suggestive at best.

632 4.2. Planet Confirmation




Figure 8. Radii, orbital periods, and ages of transiting exoplanets. Planets younger than a gigayear with $\tau/\sigma_\tau > 3$ are emphasized, where τ is the age and σ_τ is its uncertainty. Kepler 1627Ab is shown with a star. The large sizes of the youngest transiting planets could be explained by their primordial atmospheres not yet having evaporated; direct measurements of the atmospheric outflows or planetary masses would help to confirm this expectation. Selection effects may also be important. Parameters are from the NASA Exoplanet Archive (2021 Sept 15).

If the Kepler 1627Ab transit signal is created by a genuine planet, then to our knowledge it would be the youngest planet yet found by the (Replaced: main replaced with: prime) Kepler mission.⁵ Could the transit be produced by anything other than a planet orbiting this near-solar analog? Morton et al. (2016) validated the planet based on the transit shape, arguing that the most probable false positive scenario was that of a background eclipsing binary, which had a model-dependent probability of $\approx 10^{-5}$. However, this calculation was performed without knowledge of the low-mass stellar companion ($M_B \approx 0.33 M_\odot$). Validated planets have also previously been refuted (e.g., Shporer et al. 2017). We therefore reassessed false positive scenarios in some detail.

As an initial plausibility check, Kepler 1627B contributes 1% to 2% of the total flux observed in the Kepler aperture. For the sake of argument, assume the former value. The observed transit has a depth of $\approx 0.18\%$. A 18% deep eclipse of Ke-

pler 1627B would therefore be needed to produce a signal with the appropriate depth. The shape of the transit signal however requires the impact parameter to be below 0.77 (2- σ); the tertiary transiting the secondary would therefore need to be non-grazing with $R_3/R_2 \approx 0.4$. This yields a contradiction: this scenario requires an ingress and egress phase that each span $\approx 40\%$ of the transit duration (≈ 68 min). The actual measured ingress and egress duration is ≈ 17 min, a factor of four times too short. The combination of Kepler 1627B's brightness, the transit depth, and the ingress duration therefore disfavor the scenario that Kepler 1627B might host the transit signal.

Beyond this simple test, a line of evidence that effectively confirms the planetary interpretation is that the stellar density implied by the transit duration and orbital period is inconsistent with an eclipsing body around the M-dwarf companion. We find $\rho_\star = 2.00 \pm 0.24 \text{ g cm}^{-3}$, while the theoretically expected density for Kepler 1627B is $\approx 4.6 \text{ g cm}^{-3}$ (Baraffe et al. 2015). The transit duration is therefore too long to be explained by a star eclipsing the M dwarf secondary at 10- σ . While the planet might hypothetically still orbit a hidden close and bright companion, this possibility is im-

⁵ The re-purposed K2 mission however has found two younger systems containing five planets: K2-33b (9 ± 1 Myr; Mann et al. 2016; David et al. 2016) and V1298 Tau (23 ± 4 Myr; David et al. 2019).

680 plausible given *i*) the lack of secondary lines in the
 681 HIRES spectra, *ii*) the lack of secondary rotation
 682 signals in the Kepler photometry, and *iii*) the prox-
 683 imity of Kepler 1627 to the δ Lyr cluster locus on
 684 the Gaia CAMD (Figure 3).

685 The correlation noted in Section 4.1.3 between
 686 the TTVs and the local light curve slope might
 687 be an additional line of evidence in support of the
 688 planetary interpretation. Unless it is a statistical
 689 fluke (a $\approx 5\%$ possibility), then the most likely
 690 cause of the correlation is unresolved starspot
 691 crossings (Mazeh et al. 2015). These would only
 692 be possible if the planet transits the primary star,
 693 which excludes a background eclipsing binary sce-
 694 nario. The correlation would also suggest that the
 695 planet’s orbit is prograde. The latter point as-
 696 sumes that the dominant photometric variability
 697 is induced by dark spots, and not bright faculae.
 698 Given the observed transition of Sun-like stellar
 699 variability from spot to faculae-dominated regimes
 700 between young and old ages, we expect this lat-
 701 ter assumption to be reasonably secure (Shapiro
 702 et al. 2016; Montet et al. 2017; Reinhold & Hekker
 703 2020).

704 A third supporting line of evidence for the plan-
 705 etary interpretation also exists. We observed a tran-
 706 sit of Kepler 1627Ab on the night of 2021 Aug
 707 7 (Replaced: simultaneously with Keck/HIRES
 708 and MuSCAT3 replaced with: spectroscopically
 709 with HIRES at the Keck-I telescope and pho-
 710 tomatically in *griz* bands with MuSCAT3 at
 711 Haleakalā Observatory). (Added: Details of
 712 the observation sequence are discussed in Ap-
 713 pendix C; Figure 7 shows the results.) (Deleted:
 714 . We scheduled the observations using the
 715 ephemeris of Holezer et al. (2016).) Al-
 716 though we did not detect the Rossiter-McLaughlin
 717 (RM) anomaly, the multi-band MuSCAT3 light
 718 curves show that the transit is achromatic (Deleted:
 719 (Figure 7)). Quantitatively, when we fitted the
 720 MuSCAT3 photometry with a model that lets the
 721 transit depths vary across each bandpass, we found
 722 *griz* depths consistent with the Kepler depth at
 723 0.6, 0.3, 0.3, and $1.1-\sigma$ respectively. (Added:
 724 The achromatic transits strongly favor Kepler
 725 1627A as the transit host, since Kepler 1627B
 726 is much redder.) Conditioned on the ephemeris
 727 and transit depth from the Kepler data, the MuS-
 728 CAT3 observations also suggested a transit dura-
 729 tion 17.3 ± 4.3 min shorter than the Kepler transits.
 730 However, given both the lack of TDVs in the Ke-
 731 pler data and the relatively low signal-to-noise of
 732 the MuSCAT3 transit, further photometric follow-
 733 up would be necessary to determine whether the
 734 transit duration is indeed changing. For our RM
 735 analysis, the details are discussed in Appendix C.
 736 While the velocities are marginally more consistent

737 with a prograde or polar orbit than a retrograde or-
 738 bit, the spot-corrected exposure-to-exposure scat-
 739 ter ($\sigma_{RV} \approx 30 \text{ m s}^{-1}$) is comparable to the expected
 740 RM anomaly assuming an aligned orbit ($\Delta v_{RM} \approx$
 741 20 ms^{-1}). We are therefore not in a position to
 742 claim a spectroscopic detection of the RM effect,
 743 nor to quantify the stellar obliquity.

744 5. DISCUSSION & CONCLUSIONS

745 Kepler 1627Ab provides a new extremum in the
 746 ages of the Kepler planets, and opens multiple av-
 747 enues for further study. Observations of spectro-
 748 scopic transits at greater (Replaced: precision re-
 749 placed with: signal-to-noise) should yield a mea-
 750 surement of the stellar obliquity, which would con-
 751 firm or refute the prograde orbital geometry sug-
 752 gested by the TTV-local slope correlation. Sepa-
 753 rately, transit spectroscopy aimed at detecting at-
 754 mospheric outflows could yield insight into the
 755 evolutionary state of the atmosphere (e.g., Ehren-
 756 reich et al. 2015; Spake et al. 2018; Vissapra-
 757 gada et al. 2020). Observations that quantify the
 758 amount of high-energy irradiation incident on the
 759 planet would complement these efforts, by helping
 760 to clarify the expected outflow rate (e.g., Poppen-
 761 haeger et al. 2021). Finally, a challenging but infor-
 762 mative quantity to measure would be the planet’s
 763 mass. Measured at sufficient precision, for instance
 764 through a multi-wavelength radial velocity cam-
 765 paign, the combination of the size, mass, and age
 766 would yield constraints on both the planet’s com-
 767 position and its initial entropy (Owen 2020).

768 More immediately, the Kepler data may yet con-
 769 tain additional information. For instance, one pos-
 770 sible explanation for the transit asymmetry shown
 771 in Figure 5 is that of a dusty asymmetric out-
 772 flow. Dusty outflows are theoretically expected
 773 for young mini-Neptunes, and the amplitude of the
 774 observed asymmetry is consistent with predictions
 775 (Wang & Dai 2019). A second possibility is that
 776 the planetary orbit is slightly misaligned from the
 777 stellar spin axis, and tends to transit starspot groups
 778 at favored stellar latitudes. This geometry would
 779 be necessary in order to explain how the starspot
 780 crossings could add up coherently (Added: , given
 781 that the planetary orbital period (7.203 days)
 782 and the stellar rotation period (2.642 days) are
 783 not a rational combination). Other possibilities
 784 including gravity darkening or TTVs causing the
 785 asymmetry are disfavored (see Appendix B).

786 Beyond the asymmetric transits, Appendix D
 787 highlights an additional abnormality in the short-
 788 cadence Kepler data, in the arrival time distribu-
 789 tion of stellar flares. We encourage its exploration
 790 by investigators more versed in the topic than our-
 791 selves.
 792 In the context of the transiting planet population,
 793 Kepler 1627Ab is among the youngest known (Fig-

ure 8). Comparable systems with precise ages include K2-33 (Mann et al. 2016; David et al. 2016), DS Tuc (Benatti et al. 2019; Newton et al. 2019), HIP 67522 (Rizzuto et al. 2020), TOI 837 (Bouma et al. 2020), the two-planet AU Mic (Plavchan et al. 2020; Martioli et al. 2021) and the four-planet V1298 Tau (David et al. 2019). Kepler 1627Ab is one of the smaller planets in this sample ($3.82 \pm 0.16 R_{\oplus}$), which could be linked to the selection effects imposed by spot-induced photometric variability at very young ages (e.g., Zhou et al. 2021). **(Replaced: However, replaced with: However, it seems that)** smaller planets could have been detected: **(Added: based on the per-target detection contours,**) the Kepler pipeline’s median completeness extended to $1.6 R_{\oplus}$ at 10 day orbital periods, and $3.3 R_{\oplus}$ at 100 days (Burke & Catanzarite 2021). **(Added: These limits account for the spot-induced variability in the system through a correction based on the Combined Differential Photometric Precision (CDPP) in the Kepler 1627 light curve over the relevant transit timescales (Burke & Catanzarite 2017).)** The large size of Kepler 1627Ab relative to most Kepler mini-Neptunes might therefore support a picture in which the typical $5 M_{\oplus}$ **(Deleted: to $10 M_{\oplus}$)** mini-Neptune (Wu 2019) loses a significant fraction of its primordial atmosphere over its first gigayear (Owen & Wu 2013; Ginzburg et al. 2018). **(Added: This of course would need to be confirmed through a measurement of the planetary mass; alternative explanations for its large size could be that it is abnormally massive, or that it has an abnormally large envelope to core mass ratio.)**

Ultimately, the main advance of this work is a precise measurement of the age of Kepler 1627Ab. This measurement was enabled by identifying the connection of the star to the δ Lyr cluster using Gaia kinematics, and by then using the Gaia color–absolute magnitude diagram and TESS stellar rotation periods to verify the cluster’s existence. Table 3 enables similar cross-matches for both known and forthcoming exoplanet systems (e.g., Guerrero et al. 2021). Confirming these candidate associations using independent age indicators is essential because their false positive rates are not known. A related path is to identify new kinematic associations around known exoplanet host stars using positions and tangential velocities from Gaia, and to verify these associations with stellar rotation periods and spectroscopy (e.g., Tofflemire et al. 2021). Each approach seems likely to expand the census of planets with precisely measured ages over the coming years, which will help in deciphering the early stages of exoplanet evolution.

ACKNOWLEDGMENTS

The authors are grateful to J. Winn, J. Spake, A. Howard, and T. David for illuminating discussions and suggestions, and to R. Kerr for providing us with the Kerr et al. (2021) membership list prior to its publication. The authors are also grateful to K. Collins for helping resolve the scheduling conflict that would have otherwise prevented the MuSCAT3 observations. L.G.B. acknowledges support from a Charlotte Elizabeth Procter Fellowship from Princeton University, as well as from the TESS GI Program (NASA grants 80NSSC19K0386 and 80NSSC19K1728) and the Heising-Simons Foundation (51 Pegasi b Fellowship). Keck/NIRC2 imaging was acquired by program 2015A/N301N2L (PI: A. Kraus). In addition, this paper is based in part on observations made with the MuSCAT3 instrument, developed by the Astrobiology Center and under financial support by JSPS KAKENHI (JP18H05439) and JST PRESTO (JPMJPR1775), at Faulkes Telescope North on Maui, HI, operated by the Las Cumbres Observatory. This work is partly supported by JSPS KAKENHI Grant Numbers 22000005, JP15H02063, JP17H04574, JP18H05439, JP18H05442, JST PRESTO Grant Number JPMJPR1775, the Astrobiology Center of National Institutes of Natural Sciences (NINS) (Grant Number AB031010). This paper also includes data collected by the TESS mission, which are publicly available from the Mikulski Archive for Space Telescopes (MAST). Funding for the TESS mission is provided by NASA’s Science Mission directorate. We thank the TESS Architects (G. Ricker, R. Vandervort, D. Latham, S. Seager, J. Jenkins) and the many TESS team members for their efforts to make the mission a continued success. Finally, this research has made use of the Keck Observatory Archive (KOA), which is operated by the W. M. Keck Observatory and the NASA Exoplanet Science Institute (NExScI), under contract with the National Aeronautics and Space Administration. We also thank the Keck Observatory staff for their support of HIRES and remote observing. We recognize the importance that the summit of Maunakea has within the indigenous Hawaiian community, and are deeply grateful to have the opportunity to conduct observations from this mountain.

Software: `altaipony` (Ilin et al. 2021), `astrobase` (Bhatti et al. 2018), `astropy` (Astropy Collaboration et al. 2018), `astroquery` (Ginsburg et al. 2018), `corner` (Foreman-Mackey 2016), `exoplanet` (Foreman-Mackey et al. 2020), and its dependencies (Agol et al. 2020; Kip-

903 ping 2013; Luger et al. 2019; Theano Develop-
904 ment Team 2016), PyMC3 (Salvatier et al. 2016),
905 scipy (Jones et al. 2001), TESS-point (Burke
906 et al. 2020), wotan (Hippke et al. 2019).

907 *Facilities:* Astrometry: Gaia (Gaia Col-
908 laboration et al. 2018b, 2021a). Imaging:
909 Second Generation Digitized Sky Survey.
910 Keck:II (NIRC2; [www2.keck.hawaii.edu/inst/
nirc2](http://www2.keck.hawaii.edu/inst/nirc2)). Gemini:North ('Alopeke; Scott et al. 2018,
911 2021). Spectroscopy: Keck:I (HIRES; Vogt et al.
912 1994). Photometry: Kepler (Borucki et al. 2010),
913 MuSCAT3 (Narita et al. 2020), TESS (Ricker et al.
914 2015).
915

Table 1. Literature and Measured Properties for Kepler 1627

Primary Star			
TIC 120105470 GAIADR2 [†] 2103737241426734336			
Parameter	Description	Value	Source
$\alpha_{J2015.5}$	Right Ascension (hh:mm:ss)	18:56:13.6	1
$\delta_{J2015.5}$	Declination (dd:mm:ss)	+41:34:36.22	1
V	Johnson V mag.	13.11 \pm 0.08	2
G	Gaia G mag.	13.02 \pm 0.02	1
G_{BP}	Gaia BP mag.	13.43 \pm 0.02	1
G_{RP}	Gaia RP mag.	12.44 \pm 0.02	1
T	TESS T mag.	12.53 \pm 0.02	2
J	2MASS J mag.	11.69 \pm 0.02	3
H	2MASS H mag.	11.30 \pm 0.02	3
K _S	2MASS K _S mag.	11.19 \pm 0.02	3
π	Gaia EDR3 parallax (mas)	3.009 \pm 0.032	1
d	Distance (pc)	329.5 \pm 3.5	1, 4
μ_α	Gaia EDR3 proper motion	1.716 \pm 0.034	1
	in RA (mas yr ⁻¹)		
μ_δ	Gaia EDR3 proper motion	-1.315 \pm 0.034	1
	in DEC (mas yr ⁻¹)		
RUWE	Gaia EDR3 renormalized	2.899	1
	unit weight error		
RV	Systemic radial velocity (km s ⁻¹)	-16.7 \pm 1.0	5
Spec. Type	Spectral Type	G8V	5
$v \sin i_*$	Rotational velocity* (km s ⁻¹)	18.9 \pm 1.0	5
Li EW	6708Å Equiv. Width (mÅ)	235 ⁺⁵ ₋₇	5
T_{eff}	Effective Temperature (K)	5505 \pm 60	6
$\log g_*$	Surface Gravity (cgs)	4.53 \pm 0.05	6
R_*	Stellar radius (R_\odot)	0.881 \pm 0.018	6
M_*	Stellar mass (R_\odot)	0.953 \pm 0.019	6
Av	Interstellar reddening (mag)	0.2 \pm 0.1	6
[Fe/H]	Metallicity	0.1 \pm 0.1	6
P_{rot}	Rotation period (d)	2.642 \pm 0.042	7
Age	Adopted stellar age (Myr)	38 ⁺⁶ ₋₅	8
Δm_{832}	Mag difference ('Alopeke 832 nm)	3.14 \pm 0.15	9
θ_B	Position angle (deg)	92 \pm 1	9
ρ_B	Apparent separation of	0.164 \pm 0.002	9
	primary and secondary (as)		
ρ_B	Apparent separation of	53 \pm 4	1, 4, 9
	primary and secondary (AU)		
$\Delta m_{K'}$	Mag difference (NIR2 K')	2.37 \pm 0.02	10
θ_B	Position angle (deg)	95.9 \pm 0.5	10
ρ_B	Apparent separation of	0.1739 \pm 0.0010	10
	primary and secondary (as)		

NOTE—[†] The GAIADR2 and GAIAEDR3 identifiers for Kepler 1627A are identical. The secondary is not resolved in the Gaia point source catalog. * Given only $v \sin i$ and $2\pi R_*/P_{\text{rot}}$, $\cos i = 0.11^{+0.11}_{-0.08}$. Provenances are: ¹Gaia Collaboration et al. (2021a), ²Stassun et al. (2019), ³Skrutskie et al. (2006), ⁴Lindgren et al. (2021), ⁵HIPER spectra and Yee et al. (2017), ⁶Cluster isochrone (MIST adopted; PARSEC compared for quoted uncertainty), ⁷Kepler light curve, ⁸Pre-main-sequence CAMD interpolation (Section 2.2.1), ⁹'Alopeke imaging 2021 June 24 (Scott et al. 2021), ¹⁰NIRC2 imaging 2015 July 22, using the Yelda et al. (2010) optical distortion solution to convert pixel-space relative positions to on-sky relative astrometry. The “discrepancy” between the two imaging epochs likely indicates orbital motion.

Table 2. Priors and Posteriors for Model Fitted to the Long Cadence Kepler 1627Ab Light Curve.

Param.	Unit	Prior	Median	Mean	Std. Dev.	3%	97%	ESS	$\hat{R} - 1$
<i>Sampled</i>									
P	d	$\mathcal{N}(7.20281; 0.01000)$	7.2028038	7.2028038	0.0000073	7.2027895	7.2028168	7464	3.9e-04
$t_0^{(1)}$	d	$\mathcal{N}(120.79053; 0.02000)$	120.7904317	120.7904254	0.0009570	120.7886377	120.7921911	3880	2.0e-03
$\log \delta$	–	$\mathcal{N}(-6.3200; 2.0000)$	-6.3430	-6.3434	0.0354	-6.4094	-6.2767	6457	3.0e-04
$b^{(2)}$	–	$\mathcal{U}(0.000; 1.000)$	0.4669	0.4442	0.2025	0.0662	0.8133	1154	1.6e-03
u_1	–	Kipping (2013)	0.271	0.294	0.190	0.000	0.628	3604	1.5e-03
u_2	–	Kipping (2013)	0.414	0.377	0.326	-0.240	0.902	3209	1.4e-03
R_*	R_{\odot}	$\mathcal{N}(0.881; 0.018)$	0.881	0.881	0.018	0.847	0.915	8977	3.1e-04
$\log g$	cgs	$\mathcal{N}(4.530; 0.050)$	4.532	4.533	0.051	4.435	4.627	6844	1.6e-03
$\langle f \rangle$	–	$\mathcal{N}(0.000; 0.100)$	-0.0003	-0.0003	0.0001	-0.0005	-0.0000	8328	1.1e-03
$e^{(3)}$	–	Van Eylen et al. (2019)	0.154	0.186	0.152	0.000	0.459	1867	2.0e-03
ω	rad	$\mathcal{U}(0.000; 6.283)$	0.055	0.029	1.845	-3.139	2.850	3557	8.6e-05
$\log \sigma_f$	–	$\mathcal{N}(\log \langle \sigma_f \rangle; 2.000)$	-8.035	-8.035	0.008	-8.049	-8.021	9590	3.9e-04
σ_{rot}	d^{-1}	InvGamma(1.000; 5.000)	0.070	0.070	0.001	0.068	0.072	9419	1.4e-03
$\log P_{\text{rot}}$	log(d)	$\mathcal{N}(0.958; 0.020)$	0.978	0.978	0.001	0.975	0.980	8320	2.2e-04
$\log Q_0$	–	$\mathcal{N}(0.000; 2.000)$	-0.327	-0.326	0.043	-0.407	-0.246	9659	2.7e-04
$\log dQ$	–	$\mathcal{N}(0.000; 2.000)$	7.697	7.698	0.103	7.511	7.899	5824	3.7e-04
f	–	$\mathcal{U}(0.010; 1.000)$	0.01006	0.01009	0.00009	0.01000	0.01025	4645	4.0e-04
<i>Derived</i>									
δ	–	–	0.001759	0.001759	0.000062	0.001641	0.001875	6457	3.0e-04
R_p/R_*	–	–	0.039	0.039	0.001	0.037	0.042	1811	1.1e-03
ρ_*	g cm^{-3}	–	1.990	2.004	0.240	1.570	2.461	6905	2.1e-03
$R_p^{(4)}$	R_{Jup}	–	0.337	0.338	0.014	0.314	0.367	2311	1.0e-03
$R_p^{(4)}$	R_{Earth}	–	3.777	3.789	0.157	3.52	4.114	2311	1.0e-03
a/R_*	–	–	17.606	17.619	0.702	16.277	18.906	6905	2.1e-03
$\cos i$	–	–	0.027	0.025	0.010	0.004	0.040	1312	1.2e-03
T_{14}	hr	–	2.841	2.843	0.060	2.734	2.958	3199	3.6e-04
T_{13}	hr	–	2.555	2.539	0.094	2.360	2.692	1960	1.4e-03

NOTE— ESS refers to the number of effective samples. \hat{R} is the Gelman-Rubin convergence diagnostic. Logarithms in this table are base-e. \mathcal{U} denotes a uniform distribution, and \mathcal{N} a normal distribution. (1) The ephemeris is in units of BJDTDB - 2454833. (2) Although $\mathcal{U}(0, 1 + R_p/R_*)$ is formally correct, for this model we assumed a non-grazing transit to enable sampling in $\log \delta$. (3) The eccentricity vectors are sampled in the $(e \cos \omega, e \sin \omega)$ plane. (4) The true planet size is a factor of $((F_1 + F_2)/F_1)^{1/2}$ larger than that from the fit because of dilution from Kepler 1627B, where F_1 is the flux from the primary, and F_2 is that from the secondary; the mean and standard deviation of $R_p = 3.817 \pm 0.158 R_{\oplus}$ quoted in the text includes this correction, assuming $(F_1 + F_2)/F_1 \approx 1.015$.

Table 3. Young, Age-dated, and Age-dateable Stars Within the Nearest Few Kiloparsecs.

Parameter	Example Value	Description
source_id	1709456705329541504	Gaia DR2 source identifier.
ra	247.826	Gaia DR2 right ascension [deg].
dec	79.789	Gaia DR2 declination [deg].
parallax	35.345	Gaia DR2 parallax [mas].
parallax_error	0.028	Gaia DR2 parallax uncertainty [mas].
pmra	94.884	Gaia DR2 proper motion $\mu_{\alpha} \cos \delta$ [mas yr $^{-1}$].
pmdec	-86.971	Gaia DR2 proper motion μ_{δ} [mas yr $^{-1}$].
phot_g_mean_mag	6.85	Gaia DR2 G magnitude.
phot_bp_mean_mag	6.409	Gaia DR2 G_{BP} magnitude.
phot_rp_mean_mag	7.189	Gaia DR2 G_{RP} magnitude.
cluster	NASAExoArchive_ps_20210506.Uma,IR_excess	Comma-separated cluster or group name.
age	9.48,nan,nan	Comma-separated logarithm (base-10) of reported ^a age in years.
mean_age	9.48	Mean (ignoring NaNs) of age column.
reference_id	NASAExoArchive_ps_20210506,Ujjwal2020,CottenSong2016	Comma-separated provenance of group membership.
reference_bibcode	2013PASP..125..989A,2020AJ....159..166U,2016ApJS..225..15C	ADS bibcode corresponding to reference_id.

NOTE— Table 3 is published in its entirety in a machine-readable format. This table is a concatenation of the studies listed in Table 4. One entry is shown for guidance regarding form and content. In this particular example, the star has a cold Jupiter on a 16 year orbit, HD 150706b (Boisse et al. 2012). An infrared excess has been reported (Cotten & Song 2016), and the star was identified by Ujjwal et al. (2020) as a candidate UMa moving group member (≈ 400 Myr; Mann et al. 2020). The star's RV activity and TESS rotation period corroborate its youth.

Table 4. Provenances of Young and Age-dateable Stars.

Reference	N_{Gaia}	N_{Age}	$N_{G_{\text{RP}} < 16}$
Kounkel et al. (2020)	987376	987376	775363
Cantat-Gaudin & Anders (2020)	433669	412671	269566
Cantat-Gaudin et al. (2018)	399654	381837	246067
Kounkel & Covey (2019)	288370	288370	229506
Cantat-Gaudin et al. (2020)	233369	227370	183974
Zari et al. (2018) UMS	86102	0	86102
Wenger et al. (2000) Y*?	61432	0	45076
Zari et al. (2018) PMS	43719	0	38435
Gaia Collaboration et al. (2018a) $d > 250 \text{ pc}$	35506	31182	18830
Castro-Ginard et al. (2020)	33635	24834	31662
Kerr et al. (2021)	30518	25324	27307
Wenger et al. (2000) Y*○	28406	0	16205
Villa Vélez et al. (2018)	14459	14459	13866
Cantat-Gaudin et al. (2019)	11843	11843	9246
Damiani et al. (2019) PMS	10839	10839	9901
Oh et al. (2017)	10379	0	10370
Meingast et al. (2021)	7925	7925	5878
Wenger et al. (2000) pMS*	5901	0	3006
Gaia Collaboration et al. (2018a) $d < 250 \text{ pc}$	5378	817	3968
Kounkel et al. (2018)	5207	3740	5207
Ratzenböck et al. (2020)	4269	4269	2662
Wenger et al. (2000) TT*	4022	0	3344
Damiani et al. (2019) UMS	3598	3598	3598
Rizzuto et al. (2017)	3294	3294	2757
Akeson et al. (2013)	3107	868	3098
Tian (2020)	1989	1989	1394
Goldman et al. (2018)	1844	1844	1783
Cotten & Song (2016)	1695	0	1693
Gagné et al. (2018b)	1429	0	1389
Röser & Schilbach (2020) Psc-Eri	1387	1387	1107
Röser & Schilbach (2020) Pleiades	1245	1245	1019
Wenger et al. (2000) TT?	1198	0	853
Gagné & Faherty (2018)	914	0	913
Pavlidou et al. (2021)	913	913	504
Gagné et al. (2018a)	692	0	692
Ujjwal et al. (2020)	563	0	563
Gagné et al. (2020)	566	566	351
Esplin & Luhman (2019)	377	443	296
Roccatagliata et al. (2020)	283	283	232
Meingast & Alves (2019)	238	238	238
Fürnkranz et al. (2019) Coma-Ber	214	214	213
Fürnkranz et al. (2019) Neighbor Group	177	177	167
Kraus et al. (2014)	145	145	145

NOTE— Table 4 describes the provenances for the young and age-dateable stars in Table 3. N_{Gaia} : number of Gaia stars we parsed from the literature source. N_{Age} : number of stars in the literature source with ages reported. $N_{G_{\text{RP}} < 16}$: number of Gaia stars we parsed from the literature source with either $G_{\text{RP}} < 16$, or a parallax S/N exceeding 5 and a distance closer than 100 pc. The latter criterion included a few hundred white dwarfs that would have otherwise been neglected. Some studies are listed multiple times if they contain multiple tables. Wenger et al. (2000) refers to the SIMBAD database.

REFERENCES

- 916 Agol, E., Luger, R., & Foreman-Mackey, D. 2020, *AJ*,
917 159, 123
- 918 Akeson, R. L., Chen, X., Ciardi, D., et al. 2013, *PASP*,
919 125, 989
- 920 Astropy Collaboration, Price-Whelan, A. M., Sipőcz,
921 B. M., et al. 2018, *AJ*, 156, 123
- 922 Baraffe, I., Homeier, D., Allard, F., & Chabrier, G.
923 2015, *A&A*, 577, A42
- 924 Beaumont, C., Robitaille, T., Borkin, M., & Goodman,
925 A. 2014, glueviz v0.4: multidimensional data
926 exploration
- 927 Benatti, S., Nardiello, D., Malavolta, L., et al. 2019,
928 *A&A*, 630, A81
- 929 Berger, T. A., Howard, A. W., & Boesgaard, A. M.
930 2018, *ApJ*, 855, 115
- 931 Berger, T. A., Huber, D., Gaidos, E., & van Saders, J. L.
932 2018, *ApJ*, 866, 99
- 933 Bhatti, W., Bouma, L. G., & Wallace, J. 2018,
934 astrobase,
935 <https://doi.org/10.5281/zenodo.1469822>
- 936 Boisse, I., Pepe, F., Perrier, C., et al. 2012, *A&A*, 545,
937 A55
- 938 Borucki, W. J., Koch, D., Basri, G., et al. 2010, *Science*,
939 327, 977
- 940 Bossini, D., Vallenari, A., Bragaglia, A., et al. 2019,
941 *A&A*, 623, A108
- 942 Bouma, L. G., Curtis, J. L., Hartman, J. D., Winn, J. N.,
943 & Bakos, G. A. 2021, arXiv:2107.08050 [astro-ph]
- 944 Bouma, L. G., Hartman, J. D., Bhatti, W., Winn, J. N.,
945 & Bakos, G. Á. 2019, *ApJS*, 245, 13
- 946 Bouma, L. G., Hartman, J. D., Brahm, R., et al. 2020,
947 *AJ*, 160, 239
- 948 Bressan, A., Marigo, P., Girardi, L., et al. 2012,
949 *MNRAS*, 427, 127
- 950 Burke, C., & Catanzarite, J. 2021, KeplerPORTS:
951 Kepler Planet Occurrence Rate Tools
- 952 Burke, C. J., & Catanzarite, J. 2017, Planet Detection
953 Metrics: Per-Target Detection Contours for Data
954 Release 25, Kepler Science Document
955 KSCI-19111-002
- 956 Burke, C. J., Levine, A., Fausnaugh, M., et al. 2020,
957 TESS-Point: High precision TESS pointing tool,
958 Astrophysics Source Code Library, ascl:2003.001
- 959 Cantat-Gaudin, T., & Anders, F. 2020, *A&A*, 633, A99
- 960 Cantat-Gaudin, T., Jordi, C., Vallenari, A., et al. 2018,
961 *A&A*, 618, A93
- 962 Cantat-Gaudin, T., Jordi, C., Wright, N. J., et al. 2019,
963 *A&A*, 626, A17
- 964 Cantat-Gaudin, T., Anders, F., Castro-Ginard, A., et al.
965 2020, *A&A*, 640, A1
- 966 Capitanio, L., Lallement, R., Vergely, J. L., Elyajouri,
967 M., & Monreal-Ibero, A. 2017, *A&A*, 606, A65
- 968 Castro-Ginard, A., Jordi, C., Luri, X., et al. 2020, *A&A*,
969 635, A45
- 970 Choi, J., Dotter, A., Conroy, C., et al. 2016, *ApJ*, 823,
971 102
- 972 Claret, A., & Bloemen, S. 2011, *A&A*, 529, A75
- 973 Cotten, T. H., & Song, I. 2016, *ApJS*, 225, 15
- 974 Dahm, S. E. 2015, *ApJ*, 813, 108
- 975 Dai, F., Winn, J. N., Berta-Thompson, Z.,
976 Sanchis-Ojeda, R., & Albrecht, S. 2018, *AJ*, 155, 177
- 977 Damiani, F., Prisinzano, L., Pillitteri, I., Micela, G., &
978 Sciotino, S. 2019, *A&A*, 623, A112
- 979 Davenport, J. R. A. 2016, *ApJ*, 829, 23
- 980 Davenport, J. R. A., Hawley, S. L., Hebb, L., et al.
981 2014, *ApJ*, 797, 122
- 982 David, T., Hillenbrand, L., & Petigura, E. 2016, *Nature*,
983 534, 658
- 984 David, T. J., & Hillenbrand, L. A. 2015, *ApJ*, 804, 146
- 985 David, T. J., Petigura, E. A., Luger, R., et al. 2019,
986 *ApJL*, 885, L12
- 987 David, T. J., Contardo, G., Sandoval, A., et al. 2021, *AJ*,
988 161, 265
- 989 Dias, W. S., Monteiro, H., Caetano, T. C., et al. 2014,
990 *A&A*, 564, A79
- 991 Dobbie, P. D., Lodieu, N., & Sharp, R. G. 2010,
992 *MNRAS*, 409, 1002
- 993 Douglas, S. T., Agüeros, M. A., Covey, K. R., & Kraus,
994 A. 2017, *ApJ*, 842, 83
- 995 Douglas, S. T., Pérez Chávez, J., Cargile, P. A., et al.
996 2021, [10.5281/zenodo.5131306](https://doi.org/10.5281/zenodo.5131306)
- 997 Eggen, O. J. 1968, *ApJ*, 152, 77
- 998 Ehrenreich, D., Bourrier, V., Wheatley, P. J., et al. 2015,
999 *Nature*, 522, 459
- 1000 Esplin, T. L., & Luhman, K. L. 2019, *AJ*, 158, 54
- 1001 Feinstein, A. D., Montet, B. T., Johnson, M. C., et al.
1002 2021, arXiv:2107.01213 [astro-ph]
- 1003 Foreman-Mackey, D. 2016, *Journal of Open Source
Software*, 1, 24
- 1005 Foreman-Mackey, D., Czekala, I., Luger, R., et al. 2020,
1006 exoplanet-dev/exoplanet v0.2.6
- 1007 Fulton, B. J., Petigura, E. A., Howard, A. W., et al.
1008 2017, *AJ*, 154, 109
- 1009 Fürnkranz, V., Meingast, S., & Alves, J. 2019, *A&A*,
1010 624, L11
- 1011 Gagné, J., David, T. J., Mamajek, E. E., et al. 2020,
1012 *ApJ*, 903, 96

- 1013 Gagné, J., & Faherty, J. K. 2018, *ApJ*, 862, 138
 1014 Gagné, J., Roy-Loubier, O., Faherty, J. K., Doyon, R.,
 1015 & Malo, L. 2018a, *ApJ*, 860, 43
 1016 Gagné, J., Mamajek, E. E., Malo, L., et al. 2018b, *ApJ*,
 1017 856, 23
 1018 Gagné, J., David, T. J., Mamajek, E. E., et al. 2020,
 1019 *ApJ*, 903, 96
 1020 Gaia Collaboration, Babusiaux, C., van Leeuwen, F.,
 1021 et al. 2018a, *A&A*, 616, A10
 1022 Gaia Collaboration, Brown, A. G. A., Vallenari, A.,
 1023 et al. 2018b, *A&A*, 616, A1
 1024 —. 2021a, *A&A*, 649, A1
 1025 Gaia Collaboration, Smart, R. L., Sarro, L. M., et al.
 1026 2021b, *A&A*, 649, A6
 1027 Gelman, A., & Rubin, D. B. 1992, *Statistical Science*, 7,
 1028 457, publisher: Institute of Mathematical Statistics
 1029 Gibson, S. R., Howard, A. W., Marcy, G. W., et al. 2016,
 1030 in SPIE Conference Series, Vol. 9908, Ground-based
 1031 and Airborne Instrumentation for Astronomy VI, ed.
 1032 C. J. Evans, L. Simard, & H. Takami, 990870
 1033 Gillen, E., Briegal, J. T., Hodgkin, S. T., et al. 2020,
 1034 *MNRAS*, 492, 1008
 1035 Ginsburg, A., Sipocz, B., Madhura Parikh, et al. 2018,
 1036 *Astropy/Astroquery: V0.3.7 Release*
 1037 Ginzburg, S., Schlichting, H. E., & Sari, R. 2018,
 1038 *MNRAS*, 476, 759
 1039 Goldman, B., Röser, S., Schilbach, E., Moór, A. C., &
 1040 Henning, T. 2018, *ApJ*, 868, 32
 1041 Guerrero, N. M., Seager, S., Huang, C. X., et al. 2021,
 1042 *ApJS*, 254, 39
 1043 Günther, M. N., Zhan, Z., Seager, S., et al. 2020, *AJ*,
 1044 159, 60
 1045 Hattori, S., Foreman-Mackey, D., Hogg, D. W., et al.
 1046 2021, arXiv e-prints, arXiv:2106.15063
 1047 Hippke, M., David, T. J., Mulders, G. D., & Heller, R.
 1048 2019, *AJ*, 158, 143
 1049 Hirano, T., Suto, Y., Taruya, A., et al. 2010, *ApJ*, 709,
 1050 458
 1051 Hirano, T., Suto, Y., Winn, J. N., et al. 2011, *ApJ*, 742,
 1052 69
 1053 Hoffman, M. D., & Gelman, A. 2014, *Journal of
 Machine Learning Research*, 15, 1593
 1055 Holczer, T., Shporer, A., Mazeh, T., et al. 2015, *ApJ*,
 1056 807, 170
 1057 Holczer, T., Mazeh, T., Nachmani, G., et al. 2016,
 1058 *ApJS*, 225, 9
 1059 Howard, A. W., Johnson, J. A., Marcy, G. W., et al.
 1060 2010, *ApJ*, 721, 1467
 1061 Howell, S. B., Everett, M. E., Sherry, W., Horch, E., &
 1062 Ciardi, D. R. 2011, *AJ*, 142, 19
 1063 Ikoma, M., & Hori, Y. 2012, *ApJ*, 753, 66
 1064 Ilin, E., Schmidt, S. J., Poppenhöger, K., et al. 2021,
 1065 *A&A*, 645, A42
 1066 Jones, E., Oliphant, T., Peterson, P., et al. 2001, Open
 1067 source scientific tools for Python
 1068 Kerr, R. M. P., Rizzuto, A. C., Kraus, A. L., & Offner,
 1069 S. S. R. 2021, *ApJ*, 917, 23
 1070 Kharchenko, N. V., Piskunov, A. E., Schilbach, E.,
 1071 Röser, S., & Scholz, R.-D. 2013, *A&A*, 558, A53
 1072 Kipping, D. M. 2010, *MNRAS*, 408, 1758
 1073 Kipping, D. M. 2013, *MNRAS*, 435, 2152
 1074 Kite, E. S., Fegley, Jr., B., Schaefer, L., & Ford, E. B.
 1075 2020, *ApJ*, 891, 111
 1076 Klein, B., & Donati, J.-F. 2020, *MNRAS*, 493, L92
 1077 Kounkel, M., & Covey, K. 2019, *AJ*, 158, 122
 1078 Kounkel, M., Covey, K., & Stassun, K. G. 2020, *AJ*,
 1079 160, 279
 1080 Kounkel, M., Covey, K., Suárez, G., et al. 2018, *AJ*,
 1081 156, 84
 1082 Kounkel, M., Covey, K., Suárez, G., et al. 2018, *AJ*,
 1083 156, 84
 1084 Kraus, A. L., Ireland, M. J., Huber, D., Mann, A. W., &
 1085 Dupuy, T. J. 2016, *AJ*, 152, 8
 1086 Kraus, A. L., Shkolnik, E. L., Allers, K. N., & Liu,
 1087 M. C. 2014, *AJ*, 147, 146
 1088 Lallement, R., Capitanio, L., Ruiz-Dern, L., et al. 2018,
 1089 *A&A*, 616, A132
 1090 Lindegren, L., Bastian, U., Biermann, M., et al. 2021,
 1091 *A&A*, 649, A4
 1092 Lomb, N. R. 1976, *Astrophysics and Space Science*, 39,
 1093 447
 1094 Lopez, E. D., Fortney, J. J., & Miller, N. 2012, *ApJ*,
 1095 761, 59
 1096 Luger, R., Agol, E., Foreman-Mackey, D., et al. 2019,
 1097 *AJ*, 157, 64
 1098 Mann, A. W., Newton, E. R., Rizzuto, A. C., et al. 2016,
 1099 *AJ*, 152, 61
 1100 Mann, A. W., Johnson, M. C., Vanderburg, A., et al.
 1101 2020, *AJ*, 160, 179
 1102 Martioli, E., Hébrard, G., Correia, A. C. M., Laskar, J.,
 1103 & Lecavelier des Etangs, A. 2021, *A&A*, 649, A177
 1104 Masuda, K. 2015, *ApJ*, 805, 28
 1105 Mazeh, T., Holczer, T., & Shporer, A. 2015, *ApJ*, 800,
 1106 142
 1107 McCann, J., Murray-Clay, R. A., Kratter, K., &
 1108 Krumholz, M. R. 2019, *ApJ*, 873, 89
 1109 McKinney, W. 2010, in Proceedings of the 9th Python
 1110 in Science Conference, ed. S. van der Walt &
 1111 J. Millman, 51
 1112 McQuillan, A., Mazeh, T., & Aigrain, S. 2014, *ApJS*,
 1113 211, 24

- 1114 Meibom, S., Mathieu, R. D., & Stassun, K. G. 2007,
 1115 *ApJL*, **665**, L155
- 1116 Meibom, S., Barnes, S. A., Latham, D. W., et al. 2011,
 1117 *ApJL*, **733**, L9
- 1118 Meibom, S., Torres, G., Fressin, F., et al. 2013, *Nature*,
 1119 **499**, 55
- 1120 Meingast, S., & Alves, J. 2019, *A&A*, **621**, L3
- 1121 Meingast, S., Alves, J., & Rottensteiner, A. 2021, *A&A*,
 1122 **645**, A84
- 1123 Montet, B. T., Tovar, G., & Foreman-Mackey, D. 2017,
 1124 *ApJ*, **851**, 116
- 1125 Montet, B. T., Feinstein, A. D., Luger, R., et al. 2020,
 1126 *AJ*, **159**, 112
- 1127 Morris, B. M. 2020, *ApJ*, **893**, 67
- 1128 Morton, T. D., Bryson, S. T., Coughlin, J. L., et al.
 1129 2016, *ApJ*, **822**, 86
- 1130 Narita, N., Fukui, A., Yamamoto, T., et al. 2020, in
 1131 *SPIE Conference Series*, 114475K
- 1132 Newton, E. R., Mann, A. W., Tofflemire, B. M., et al.
 1133 2019, *ApJ*, **880**, L17
- 1134 Oh, S., Price-Whelan, A. M., Hogg, D. W., Morton,
 1135 T. D., & Spergel, D. N. 2017, *AJ*, **153**, 257
- 1136 Owen, J. E. 2020, *MNRAS*, **498**, 5030
- 1137 Owen, J. E., & Wu, Y. 2013, *ApJ*, **775**, 105
 1138 —. 2016, *ApJ*, **817**, 107
- 1139 Palle, E., Oshagh, M., Casasayas-Barris, N., et al. 2020,
 1140 *A&A*, **643**, 25
- 1141 Pavlidou, T., Scholz, A., & Teixeira, P. S. 2021,
 1142 *MNRAS*, **503**, 3232
- 1143 Pecaut, M. J., & Mamajek, E. E. 2013, *ApJS*, **208**, 9
- 1144 Pecaut, M. J., & Mamajek, E. E. 2016, *MNRAS*, **461**,
 1145 794
- 1146 Plavchan, P., Barclay, T., Gagné, J., et al. 2020, *Nature*,
 1147 **582**, 497
- 1148 Poppenhaeger, K., Ketzer, L., & Mallonn, M. 2021,
 1149 *MNRAS*, **500**, 4560
- 1150 Randich, S., Tognelli, E., Jackson, R., et al. 2018, *A&A*,
 1151 **612**, A99
- 1152 Ratzenböck, S., Meingast, S., Alves, J., Möller, T., &
 1153 Bomze, I. 2020, *A&A*, **639**, A64
- 1154 Rebull, L. M., Stauffer, J. R., Bouvier, J., et al. 2016,
 1155 *AJ*, **152**, 113
- 1156 Reinhold, T., & Hekker, S. 2020, *A&A*, **635**, A43
- 1157 Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015,
 1158 *JATIS*, **1**, 014003
- 1159 Rizzuto, A. C., Mann, A. W., Vanderburg, A., Kraus,
 1160 A. L., & Covey, K. R. 2017, *AJ*, **154**, 224
- 1161 Rizzuto, A. C., Newton, E. R., Mann, A. W., et al. 2020,
 1162 *AJ*, **160**, 33
- 1163 Roccatagliata, V., Franciosini, E., Sacco, G. G.,
 1164 Randich, S., & Sicilia-Aguilar, A. 2020, *A&A*, **638**,
 1165 A85
- 1166 Roettenbacher, R. M., Monnier, J. D., Korhonen, H.,
 1167 et al. 2017, *ApJ*, **849**, 120
- 1168 Röser, S., & Schilbach, E. 2020, *A&A*, **638**, A9
- 1169 Salvatier, J., Wiecki, T. V., & Fonnesbeck, C. 2016,
 1170 PyMC3: Python probabilistic programming
 1171 framework
- 1172 Scargle, J. D. 1982, *ApJ*, **263**, 835
- 1173 Scott, N. J., Howell, S. B., Horch, E. P., & Everett,
 1174 M. E. 2018, *PASP*, **130**, 054502
- 1175 Scott, N. J., Howell, S. B., Gnilka, C. L., et al. 2021,
 1176 *Frontiers in Astronomy and Space Sciences*, **8**, 138
- 1177 Seifahrt, A., Stürmer, J., Bean, J. L., & Schwab, C.
 1178 2018, in *SPIE Conference Series*, Vol. 10702,
 1179 *Ground-based and Airborne Instrumentation for
 1180 Astronomy VII*, ed. C. J. Evans, L. Simard, &
 1181 H. Takami, 107026D
- 1182 Shapiro, A. I., Solanki, S. K., Krivova, N. A., Yeo,
 1183 K. L., & Schmutz, W. K. 2016, *A&A*, **589**, A46
- 1184 Shporer, A., Zhou, G., Vanderburg, A., et al. 2017,
 1185 *ApJL*, **847**, L18
- 1186 Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006,
 1187 *AJ*, **131**, 1163
- 1188 Smith, J. C., Morris, R. L., Jenkins, J. M., et al. 2016,
 1189 *PASP*, **128**, 124501
- 1190 Smith, J. C., Stumpe, M. C., Jenkins, J. M., et al. 2017,
 1191 *Kepler Science Document*, 8
- 1192 Soderblom, D. R., Hillenbrand, L. A., Jeffries, R. D.,
 1193 Mamajek, E. E., & Naylor, T. 2014, *Protostars and
 1194 Planets VI*, 219
- 1195 Spake, J. J., Sing, D. K., Evans, T. M., et al. 2018,
 1196 *Nature*, **557**, 68
- 1197 Stassun, K. G., Oelkers, R. J., Paegert, M., et al. 2019,
 1198 *AJ*, **158**, 138
- 1199 Stauffer, J., Rebull, L., Bouvier, J., et al. 2016, *AJ*, **152**,
 1200 115
- 1201 Stauffer, J. R., Hartmann, L. W., Prosser, C. F., et al.
 1202 1997, *ApJ*, **479**, 776
- 1203 Stefansson, G., Mahadevan, S., Maney, M., et al. 2020,
 1204 *AJ*, **160**, 192
- 1205 Stephenson, C. B. 1959, *PASP*, **71**, 145
- 1206 Strassmeier, K. G. 2009, *Astronomy and Astrophysics
 1207 Review*, **17**, 251
- 1208 Tenenbaum, P., Christiansen, J. L., Jenkins, J. M., et al.
 1209 2012, *ApJS*, **199**, 24
- 1210 Theano Development Team. 2016, *arXiv e-prints*,
 1211 *abs/1605.02688*
- 1212 Thompson, S. E., Coughlin, J. L., Hoffman, K., et al.
 1213 2018, *ApJS*, **235**, 38

- 1214 Tian, H.-J. 2020, [ApJ](#), 904, 196
- 1215 Tofflemire, B. M., Rizzato, A. C., Newton, E. R., et al.
1216 2021, [AJ](#), 161, 171
- 1217 Ujjwal, K., Kartha, S. S., Mathew, B., Manoj, P., &
1218 Narang, M. 2020, [AJ](#), 159, 166
- 1219 Van Eylen, V., Agentoft, C., Lundkvist, M. S., et al.
1220 2018, [MNRAS](#), 479, 4786
- 1221 Van Eylen, V., Albrecht, S., Huang, X., et al. 2019, [AJ](#),
1222 157, 61
- 1223 Villa Vélez, J. A., Brown, A. G. A., & Kenworthy,
1224 M. A. 2018, [RNAAS](#), 2, 58
- 1225 Vissapragada, S., Knutson, H. A., Jovanovic, N., et al.
1226 2020, [AJ](#), 159, 278
- 1227 Vogt, S. S., Allen, S. L., Bigelow, B. C., et al. 1994,
1228 SPIE Conference Series, ed. D. L. Crawford & E. R.
1229 Craine, Vol. 2198
- 1230 Walkowicz, L. M., & Basri, G. S. 2013, [MNRAS](#), 436,
1231 1883, arXiv: 1309.2159
- 1232 Wang, L., & Dai, F. 2019, [ApJL](#), 873, L1
- 1233 Wenger, M., Ochsenbein, F., Egret, D., et al. 2000,
1234 [A&AS](#), 143, 9
- 1235 Wirth, C. P., Zhou, G., Quinn, S. N., et al. 2021, [ApJL](#),
1236 917, L34
- 1237 Wu, Y. 2019, [ApJ](#), 874, 91
- 1238 Yee, S. W., Petigura, E. A., & von Braun, K. 2017, [ApJ](#),
1239 836, 77
- 1240 Yelda, S., Lu, J. R., Ghez, A. M., et al. 2010, [ApJ](#), 725,
1241 331
- 1242 Zari, E., Hashemi, H., Brown, A. G. A., Jardine, K., &
1243 de Zeeuw, P. T. 2018, [A&A](#), 620, A172
- 1244 Zhou, G., Winn, J. N., Newton, E. R., et al. 2020, [ApJ](#),
1245 892, L21
- 1246 Zhou, G., Quinn, S. N., Irwin, J., et al. 2021, [AJ](#), 161, 2

1247

APPENDIX

1248

A. YOUNG, AGE-DATED, AND AGE-DATEABLE STAR COMPILED

1249 The v0.6 CDIPS target catalog (Table 3) includes stars that are young, age-dated, and age-dateable. By
 1250 “age-dateable”, we mean that the stellar age should be measurable at greater precision than that of a typical
 1251 FGK field star, through either isochronal, gyrochronal, or spectroscopic techniques. As in Bouma et al.
 1252 (2019), we collected stars that met these criteria from across the literature. Table 4 gives a list of the studies
 1253 included, and brief summary statistics. The age measurement methodologies adopted by each study differ: in
 1254 many, spatial and kinematic clustering has been performed on the Gaia data, and ensemble isochrone fitting
 1255 of the resulting clusters has been performed (typically focusing on the turn-off). In other studies however,
 1256 the claim of youth is based on the location of a single star in the color-absolute magnitude diagram, or on
 1257 spectroscopic information.

1258 One major change in Table 3 relative to the earlier iteration from Bouma et al. (2019) is that the extent
 1259 of Gaia-based analyses has now matured to the point that we can neglect pre-Gaia cluster memberships,
 1260 except for a few cases with spectroscopically confirmed samples of age-dated stars. The membership lists for
 1261 instance of Kharchenko et al. (2013) and Dias et al. (2014) (MWSC and DAML) are no longer required. This
 1262 is helpful for various post-processing projects, since the field star contamination rates were typically much
 1263 higher in these catalogs than in the newer Gaia-based catalogs.

1264 The most crucial parameters of a given star for our purposes are the Gaia DR2 `source_id`, the cluster
 1265 or group name (`cluster`), and the `age`. Given the hierarchical nature of many stellar associations, we do
 1266 not attempt to resolve the cluster names to a single unique string. The Orion complex for instance can be
 1267 divided into almost one hundred kinematic subgroups (Kounkel et al. 2018). (**Added: Based on Figure 1,**
 1268 **the δ Lyr cluster may also be part of a similar hierarchical association.**) Similar complexity applies to the
 1269 problem of determining homogeneous ages, which we do not attempt to resolve. Instead, we simply merged
 1270 the cluster names and ages reported by various authors into a comma-separated string.

1271 This means that the `age` column can be null, for cases in which the original authors did not report an age,
 1272 or for which a reference literature age was not readily available. Nonetheless, since we do prefer stars with
 1273 known ages, we made a few additional efforts to populate this column. When available, the age provenance
 1274 is from the original analysis of the cluster. In a few cases however we adopted other ages when string-
 1275 based cross-matching on the cluster name was straightforward. In particular, we used the ages determined
 1276 by Cantat-Gaudin et al. (2020) to assign ages to the clusters from Gaia Collaboration et al. (2018a), Cantat-
 1277 Gaudin et al. (2018), Castro-Ginard et al. (2020), and Cantat-Gaudin & Anders (2020).

1278 The catalogs we included for which ages were not immediately available were those of Cotten & Song
 1279 (2016), Oh et al. (2017), Zari et al. (2018), Gagné et al. (2018a), Gagné et al. (2018b), Gagné & Faherty
 1280 (2018), and Ujjwal et al. (2020). While in principle the moving group members discussed by Gagné et al.
 1281 (2018a,b); Gagné & Faherty (2018) and Ujjwal et al. (2020) have easily associated ages, our SIMBAD cross-
 1282 match did not retain the moving group identifiers given by those studies, which should therefore be recovered
 1283 using tools such as BANYAN Σ (Gagné et al. 2018b). We also included the SIMBAD object identifiers `TT*`,
 1284 `Y*O,Y*?`, `TT?`, and `pMS*`. Finally, we included every star in the NASA Exoplanet Archive planetary system
 1285 (`ps`) table that had a Gaia identifier available (Akeson et al. 2013). If the age had finite uncertainties, we also
 1286 included it, since stellar ages determined through the combination of isochrone-fitting and transit-derived
 1287 stellar densities typically have higher precision than from isochrones alone.

1288 For any of the catalogs for which Gaia DR2 identifiers were not available, we either followed the spa-
 1289 tial (plus proper-motion) cross-matching procedures described in Bouma et al. (2019), or else we pulled the
 1290 Gaia DR2 source identifiers associated with the catalog from SIMBAD. We consequently opted to drop the
 1291 `ext_catalog_name` and `dist` columns maintained in Bouma et al. (2019), as these were only popu-
 1292 lated for a small number of stars. The technical manipulations for the merging, cleaning, and joining were
 1293 performed using pandas (McKinney 2010). The eventual cross-match (using the Gaia DR2 `source_id`)
 1294 against the Gaia DR2 archive was performed asynchronously on the Gaia archive website.

1295

B. THE TRANSIT ASYMMETRY

1296

B.1. How Robust is the Asymmetric Transit?

1297 As a means of exploring the robustness of the transit asymmetry, Figures 9, 10, and 11 show the Kepler data
 1298 binned in three ways: over Kepler quarters, Julian years, and quartiles of local slope. Over Kepler quarters
 1299 (Figure 9), Quarter 6 shows the strongest asymmetry: a deviation of about 3 ppt from expectation. Quarter




Figure 9. Transit model residuals through time (binned by Kepler quarter). *Left:* Phase-folded transit of Kepler 1627b, with stellar variability removed. Black points are binned to 20 minute intervals. The 2σ model uncertainties and the maximum *a posteriori* model are shown as the faint purple band, and the dark purple line. *Right:* As on the left, with the transit removed.




Figure 10. Transit model residuals through time (binned by year of observation). *Left:* Phase-folded transit of Kepler 1627b, with stellar variability removed. Points and models are as in Figure 9. *Right:* As on the left, with the transit removed.




Figure 11. Transit models and residuals, binned by quartiles in the local slope of the light curve. Representative uncertainties for the black points (binned at 20 minute intervals) are shown in the lower right of each panel. A similar transit asymmetry to that shown in Figure 5 seems to be present in three of the four bins.

1300 7 shows an anomaly at roughly the same transit phase. Year 2 correspondingly shows the strongest anomaly
 1301 out of any year in Figure 10; the asymmetry is visually apparent however in each of the four years.

1302 To bin by quartiles in local slope, we used our measurements of the local linear slopes in each of the
 1303 observed transit windows (144 transits total). Four outlier transits were removed, leaving 140 transits. These
 1304 were then divided into quartiles, so that each panel shows 35 transits binned together. The exact light curve
 1305 slope intervals are listed in the lower left panels of Figure 11. Binned by local slope quartiles (Figure 11),
 1306 the asymmetry is visually present in three of the four quartiles: the only bin in which it does not appear is
 1307 $df/dt \in [3.0, 32.5] \text{ ppt day}^{-1}$.

1308 Within the theory presented by Mazeh et al. (2015), unresolved starspot crossings cause the weak correlation
 1309 between TTVs and the local light curve slope (Figure 6). In this model, we would expect the light curves
 1310 with the most negative local slopes to have the largest positive TTVs, due to spot crossing events during the
 1311 latter half of transit. The upper-left panel of Figure 11 agrees with this expectation. However, we would also
 1312 expect the sign of the effect to reverse when considering the most positive local slopes (most negative TTVs).
 1313 The lower-right panel of Figure 11 contradicts this expectation: the residual in both cases maintains the same
 1314 parity! On the one hand, this shows that the residual is not dependent on the local light curve slope, which
 1315 lowers the likelihood that it might be an artifact of our detrending methods. On the other, it raises the question
 1316 of whether unresolved starspot crossings are indeed the root cause of the correlation shown in Figure 6. While
 1317 we do not have a solution to this contradiction, the injection-recovery tests discussed in Section 4.1.3 provide
 1318 some assurance that the TTV-slope correlation is not simply a systematic artifact.

1319

B.2. Interpretation

1320 The transit asymmetry seems robust against most methods of binning the data, though with some caveats
 1321 (e.g., the “middle quartile” in local flux, $df/dt \in [3.0, 32.5] \text{ ppt day}^{-1}$, where the asymmetry does not appear).
 1322 Nonetheless, if the asymmetric were systematic we might expect its parity to reverse as a function of the sign
 1323 of the local slope, and it does not. We therefore entertained four possible astrophysical explanations: gravity
 1324 darkening, transit timing variations, spot-crossing events, and a persistent asymmetric dusty outflow.

1325 Gravity darkening is based on the premise that the rapidly rotating star is oblate, and brighter near the
 1326 poles than the equator (e.g., Masuda 2015). The fractional transit shape change due to gravity darkening
 1327 is on the order of $(P_{\text{break}}/P_{\text{rot}})^2$, for P_{break} the break-up rotation period, and P_{rot} the rotation period. Using

the parameters from Table 2, this yields an expected 0.14% distortion of the \approx 1.8 ppt transit depth: *i.e.*, an absolute deviation of \approx 2.5 ppm. The observed residual has a semi-amplitude of \approx 50 ppm. Since the expected signal is smaller than the observed anomaly by over an order of magnitude, gravity darkening seems to be an unlikely explanation.

The scenario of transit timing variations (TTVs) producing the asymmetry seems unlikely because the transit timing variations we do observe are correlated with the local light curve slope, which increases roughly as much as it decreases. From our analysis, the mean TTV and its standard deviation are 0.66 ± 5.53 min; similarly the mean local slope and its standard deviation are 0.59 ± 45.50 ppt day $^{-1}$. There is therefore little expectation for TTVs to produce the asymmetry. A separate line of argument comes from Figure 11. If the local slope were essential to producing the transit asymmetry, we would expect that in the largest df/dt bin, $df/dt \in [3.0, 32.5]$ ppt day $^{-1}$, the sign of the asymmetry would reverse. We do not see evidence for this being the case.

The third and related possibility is that of starspot crossings. Young stars have higher spot-covering fractions than old stars (*e.g.*, Morris 2020). Young solar-type stars may also host dark starspots at high stellar latitudes (*e.g.*, EK Dra; Strassmeier 2009), though interferometric imaging of spotted giant stars has shown different starspot latitude distributions than those inferred from Doppler imaging (Roettenbacher et al. 2017). Regardless, for any spot-crossing anomalies to add coherently over the 140 Kepler transits, it seems likely that we would need either for spots to be persistent at a particular latitude (and for the planetary orbit to be somewhat misaligned), or for a “stroboscopic” longitudinal phasing (*e.g.*, Dai et al. 2018). For our system, $P_{\text{orb}}/P_{\text{rot}} \approx 2.76$, which means that every 4 transits and 11 stellar rotations, the planet crosses over roughly the same stellar longitude, which might enable the necessary phasing if the spot-groups are large and long-lived. Unfortunately, the S/N per Kepler transit is ≈ 8 , which renders individual spot-crossing events unresolved. This explanation seems marginally plausible, mainly because the expected spot-crossing anomaly amplitudes (\approx 100 ppm) resemble the observed amplitude of the asymmetry (\approx 50 ppm). One issue with this explanation however is that there is no reason to expect starspot crossing events to last exactly half the transit duration.

A persistent feature of the planet itself might therefore be needed to explain the transit asymmetry. An asymmetric outflow from the planet’s atmosphere could at least geometrically meet the requirements (*e.g.*, McCann et al. 2019). To explain the asymmetric transit, a small, dense component would lead the planet, and a long, more rarefied (and variable) component would trail it. This might also explain the slight flux decrement visible for \sim 1 hour pre-ingress (Figure 5). The amplitude of the asymmetry is roughly in line with theoretical expectations for dusty outflows (Wang & Dai 2019), and based on the planet’s size, its mass is likely in a regime where such outflows are possible. Out of the four explanations discussed, this one at least theoretically seems the most plausible. By composition, the expectation would be that the envelope is mostly hydrogen and helium gas, with a dust or haze component providing the broadband opacity in the Kepler bandpass. A natural path for testing this idea would be to observe additional transits of the planet in hydrogen absorption, metastable helium absorption, or across a broad wavelength range in the near-infrared.

1364 C. SPECTROSCOPY AND PHOTOMETRY DURING THE TRANSIT OF 2021 AUG 7

1365 (Added: We used the ephemeris of Holczer et al. (2016) to observe a transit of Kepler 1627Ab on
 1366 the night of 2021 Aug 7 both spectroscopically and photometrically. We used the HIRES echelle spec-
 1367 trograph at the Keck-I telescope and MuSCAT3 at Haleakalā Observatory on Maui, HI (Narita et al.
 1368 2020).) (Deleted: -We monitored Kepler 1627 with Keck/HIRES before, during, and after transit on
 1369 the night of 2021 Aug 7.) (Replaced: We used the iodine cell for wavelength calibration replaced with:
 1370 For the HIRES wavelength calibration, we used the iodine cell), and extracted the 1-D spectra using the
 1371 standard California Planet Survey pipeline (Howard et al. 2010). (Added: Given the faintness of the tar-
 1372 get ($V = 13.1$), we observed using the C2 decker, which yielded an instrument resolution of \approx 50,000.)
 1373 The airmass ranged between 1.1 and 2.2 from the start through the end of observations; the seeing ranged
 1374 from $1.^{\prime\prime}1$ at the beginning to $1.^{\prime\prime}5$ at the end. (Added: The HIRES exposure time was set at \approx 15 min-
 1375 utes in order to resolve the 2.8 hour transit event, which yielded a S/N per resolution element of \approx 75
 1376 (low for precision radial velocity standards).) (Replaced: We also simultaneously observed across *griz*
 1377 bands using MuSCAT3 at Haleakalā Observatory on Maui, HI. replaced with: For the MuSCAT3 ob-
 1378 servations, we observed simultaneously across the *griz* bands.) (Added: The exposure times in each
 1379 bandpass ranged between 23 and 46 seconds, and were chosen in order to yield a S/N in the peak pixel
 1380 that exceeded 130 while also preventing saturation.) Performing aperture photometry on the latter image
 1381 stack yielded the data given in Table 5.




Figure 12. Spectroscopic activity indicators during the transit of 2021 Aug 7. The *top panels* show the median line profiles Ca K, Ca H, and H α line profiles from the HIRES spectra. The *lower panels* show the differences of each individual spectrum relative to the median spectrum. The bump in the red wing of Ca H is He. The spectra in the lower panels are smoothed for visualization purposes.

We considered two approaches to measuring the velocities: in the first, hereafter “Method 1”, we cross-correlated against a template found via spectral classification with SpecMatch-Emp (Yee et al. 2017). In “Method 2”, we used a high S/N template of V1298 Tau. Although V1298 Tau is cooler than Kepler 1627A by ≈ 500 K, it has a comparable amount of line-broadening ($v \sin i = 23$ km s $^{-1}$), and a comparable level of stellar activity. The mean and standard deviation of the internal RV uncertainties averaged over all epochs were 16.2 ± 1.1 m s $^{-1}$ from Method 1, and 12.6 ± 0.6 m s $^{-1}$ from Method 2. The corresponding time-averaged reduced χ^2 from the template match was 1.57 ± 0.04 (Method 1) and 1.30 ± 0.02 (Method 2). Given these diagnostics, we adopted the velocities from the second approach, which are reported in Table 6.

Figure 7 shows the results. The MuSCAT3 photometry shows the expected starspot trend, along with the transit and what is likely a chromatic starspot crossing event at JD – 2459433 = 0.955. The radial velocities decrease by ≈ 250 m s $^{-1}$ over the six hour window. This decrease in RV is correlated with a decrease in the S-indices derived from the Ca HK lines. One outlying RV point is apparent shortly before egress; it is temporally coincident with an outlying value in the S-index time series.

Overall, we expect the dominant trends in both the photometry and radial velocities to be caused by starspots on the stellar photosphere rotating into and out of view. The plasma in the leading and receding limbs of the stellar disk has an apparent line-of-sight velocity of ± 20 km s $^{-1}$. Over 10% of a rotation cycle ($P_{\text{rot}} = 2.6$ days), spots near these limbs come into and out of view, modulate the stellar velocity profile, and can thereby produce the overall ≈ 250 m s $^{-1}$ trend. The Ca HK and H α emission profiles support this interpretation; Figure 12 shows that each line gradually decreases in intensity over the course of the six hour sequence.

The expectation however is for the starspot-induced signals to be smooth, at worst with contributions at $0.5P_{\text{rot}}$ or $0.25P_{\text{rot}}$ (Klein & Donati 2020). We therefore fitted the RVs using the Hirano et al. (2010, 2011) models for the Rossiter-McLaughlin (RM) effect, and allowed for an optional linear and quadratic trend in time to fit the ≈ 250 m s $^{-1}$ spot-induced trend. We followed the methodology developed by Stefansson et al. (2020). We allowed the sky-projected obliquity, the projected stellar equatorial velocity, and the Gaussian dispersion of the spectral lines to vary, and fixed the limb-darkening using the V-band tabulation from Claret & Bloemen (2011). We assumed a Gaussian prior on $v \sin i$ and a/R_{\star} from Table 1, and also allowed for a white-noise jitter term to be added in quadrature to the measurement uncertainties. We used a 15 minute exposure time to numerically evaluate the model.

1410 The quadratic model with the RM effect is shown in Figure 7; the jitter term is incorporated in the model
 1411 uncertainties, but not the plotted measurement uncertainties. The plotted measurement uncertainties are the
 1412 internal uncertainties on the RVs ($\approx 13 \text{ m s}^{-1}$), and are dominated by the $v \sin i$ broadening. However, between
 1413 exposures, the RVs show significant additional scatter that is not captured by the slow quadratic trend. The
 1414 white-noise jitter for this particular model is $\sigma_{\text{RV}} = 27^{+6}_{-5} \text{ m s}^{-1}$, which is comparable to the expected RM
 1415 anomaly of $\Delta v_{\text{RM}} \approx f_{\text{LD}} \cdot \delta \cdot v \sin i \cdot \sqrt{1 - b^2} \approx 20 \text{ m s}^{-1}$, assuming a perfectly aligned orbit.

The presence of this additional scatter prevents a convincing detection of the RM effect. The reason can be understood via model comparison. If we compare the model with a quadratic trend and the RM effect against a model with a linear trend and the RM effect, or even a model with no RM effect at all, then the respective Bayesian Information Criterion (BIC) values are as follows.

$$\begin{aligned} \text{BIC} &= 227.1 \quad (\text{Quadratic + RM}) \\ \text{BIC} &= 231.1 \quad (\text{Linear + RM}) \\ \text{BIC} &= 221.4 \quad (\text{Only Quadratic}). \end{aligned} \quad (\text{C1})$$

1416 There is therefore no evidence to prefer the model with the RM effect against a model that only accounts
 1417 for the stellar variability. The “only quadratic” model does particularly well because it can inflate the jitter
 1418 term to account for scatter during the transit (even if the scatter contains astrophysics!), and it has fewer free
 1419 parameters. However, we cannot justify a physical prior on the jitter term, because we do not understand
 1420 the origin of the exposure-to-exposure scatter. As noted above, the velocity deviations from starspots are
 1421 expected to have contributions at the stellar rotation frequency, or harmonics thereof. This jitter is present on
 1422 the exposure timescale (15 minutes), which is only 0.4% of the stellar rotation period; it is not obvious that
 1423 starspots would be the culprit.

1424 The amplitude of both the spot-induced trend and the jitter are somewhat larger than recent comparable
 1425 measurements in systems such as AU Mic (Palle et al. 2020), DS Tuc (Montet et al. 2020; Zhou et al. 2020)
 1426 and TOI 942 (Wirth et al. 2021). One possible explanation for the jitter is that it is astrophysical in origin,
 1427 and that it is caused by some novel process operating on the surface of Kepler 1627A. Another possibility
 1428 is that our RV analysis underestimates our measurement uncertainties; in order to achieve the requisite time-
 1429 sampling the S/N per resolution element in our spectra was 70 to 80, which is lower than desired for deriving
 1430 high-precision velocities. In addition, the rapid rotation of the star could affect accuracy of the uncertainties
 1431 from the velocity extraction. Observations at higher S/N are necessary to distinguish these two possibilities,
 1432 and remain worthwhile in order to clarify the orbital geometry of Kepler 1627Ab. Useful next steps would
 1433 include transit observations with a stabilized spectrograph in the optical (e.g., Gibson et al. 2016; Seifahrt
 1434 et al. 2018), or in the near-infrared (e.g., Feinstein et al. 2021).

Table 5. MuSCAT3 photometry of Kepler 1627.

Time [BJD _{TDB}]	Rel. Flux	Rel. Flux Err.	Bandpass
2459433.829202	0.99719	0.00091	g
2459433.829324	0.99849	0.00112	r
2459433.829117	0.99611	0.00116	i
2459433.829406	0.99941	0.00136	z

NOTE— Table 5 is published in its entirety in a machine-readable format. Example entries are shown for guidance regarding form and content.

Table 6. Kepler 1627 radial velocities.

Time [BJD _{TDB}]	RV [m s^{-1}]	σ_{RV} [m s^{-1}]	S-value
2459433.801306	152.97	12.29	0.7355
2459433.812255	100.5	13.23	0.7434

NOTE— Table 6 is published in its entirety in a machine-readable format. Example entries are shown for guidance regarding form and content.

1435 **D. FLARE ANALYSIS** 1438 acquired over 97.7 days during Quarter 15. The
 1436 In addition to the 3.9 years of long cadence data,
 1437 short cadence (1-minute) Kepler observations were




Figure 13. Flares in Kepler 1627. *Top:* The 1-minute cadence Kepler data (black points) is shown with a stellar variability model superposed (blue line). *Middle:* Residual after subtracting the stellar variability model. Flares appear as spikes. *Bottom:* Zooms of the brightest, and third-brightest flares. A timing coincidence – that both flares have “successors” approximately one (Added: planetary) orbital period after the initial event – is emphasized.




Figure 14. Statistics of inter-flare arrival times. 24 flares were recorded with amplitudes exceeding 0.5% over the 97.7 days of short cadence observations. The histogram of the time intervals between every possible pair of flares is shown in black. Some plausibly important timescales for star-planet interactions, namely the planetary orbital period and synodic period (the orbital period as seen from the rotating stellar frame) are shown along with their linear combinations. Monte Carlo draws from a Poisson distribution are shown with the gray bands. While peaks in the observed distribution do coincide with some of the “special periods”, the statistical evidence for a non-Poissonian process driving the flares does not clear the 5σ threshold.

short cadence light curve shows a higher rate of flaring than visible in the long cadence data (Figure 13). We analyzed the short cadence light curve and its flares according to the following procedure.

1. Fit the starspot-induced variability using a Gaussian Process with a SHOTerm kernel, a white-noise jitter term, and the mean flux.
2. Select points more than twice the median absolute deviation from the residual, and exclude them from the light curve (these points include the flares). Repeat Step 1.
3. Using the residual from Step 2, identify all flares, requiring them to be at least 20 cadences apart, at least 7 median absolute deviations above the median baseline, and lasting at least 2 cadences in duration. Build the mask spanning these times, from 5 minutes before each flare begins to 2.5 minutes after the final flare cadence. Repeat Step 1 a final time.

The final step of flare identification and fitting was performed using `altaipony` (Davenport 2016; Ilin et al. 2021). The analytic flare model is from Davenport et al. (2014) and it parametrizes the flare with a start time, an exponential lag time, and an amplitude.

There were $N_f = 24$ flares that exceeded 0.5% in relative flux during the short cadence observations. These 24 flares spanned a total of 6.5 hours (~ 15 minutes per flare). Inspecting the data, we noticed a coincidence in the flare arrival times. The coincidence is that despite the low flare duty cycle, one orbital period after the brightest flare, a second flare followed. This and a similar event are shown in Figure 13. The timing error is good to a $\approx 0.2\%$ difference from the orbital period, which given the duty cycle seems *a priori* unlikely. If we consider flares falling within 2% of the planet’s orbital period after a previous flare, then 4 of the 24 flare events have candidate “successors”.

As with any coincidence, if one does not have a firm prediction, it is difficult to assess whether a surprise is statistically significant. Since our surprise was specifically at the inter-arrival time of certain flares coinciding with special time intervals, we performed the following analysis. First, we considered all unordered pairs of flares. For N flares there are $\binom{n}{2}$ such pairs (for our case, 276 pairs). We then compared the distribution of the pair separations against that of a Poisson distribution. Specifically, we drew $N_f = 24$ samples from a Poisson distribution with $\lambda = \Delta t / N_f$, for $\Delta t = 97.7$ days the total duration of the observations, and repeated the draw 10^3 times with unique random seeds.

Figure 14 shows the results. The vertical lines in the figure show the planetary orbital period, the synodic period $P_{\text{syn}} = (P_{\text{rot}}^{-1} - P_{\text{orb}}^{-1})^{-1}$, and linear combinations thereof. The tidal period (half the synodic period) is not shown. The bins are logarithmically spaced to give 100 bins between the minimum and maximum ordinate values. The gray bands express the range of values observed from the Poissonian draws. While it does seem like an odd coincidence for peaks in the observed flare arrival time distribution to coincide with the locations of these “special intervals”, the statistical evidence for a non-Poissonian process driving the flares does not seem especially overwhelming. More quantitatively, the peaks observed at the orbital and synodic periods are within the $\pm 2\sigma$ range of a Poissonian process, and those at $P_{\text{orb}} + P_{\text{syn}}$ and $P_{\text{orb}} + 2P_{\text{syn}}$ are only slightly above this range. With that said, future analyses of these data by investigators with more knowledge of this topic could very well yield more quantitative insights. Such analyses should keep in mind an important caveat: the amplitude distribution of M-dwarf flares extends up to many times the quiescent flux (see Figure 7 of Günther et al. 2020). A flare on Kepler 1627B producing double its quiescent white-light flux would yield a $\approx 1\%$ apparent amplitude. Such flares could represent a significant fraction of those in the Kepler observations.

List of Changes

Replaced: **mini-Neptune** replaced with: **planet**, on page 1.

Replaced: **main** replaced with: **prime**, on page 1.

Replaced: **through** replaced with: **with**, on page 1.

Replaced: **main** replaced with: **prime**, on page 2, line 74.

Replaced: **mini-Neptune** replaced with: **Neptune-sized planet**, on page 2, line 100.

Added: **to clarify the direction along which parallax uncertainties are expected to produce erroneous clusters**, on page 3.

Added: **we**, on page 2, line 136.

Added: We calculated the tangential velocities for each of these stars relative to Kepler 1627 ($\Delta\mu^*$) by subtracting the observed proper motion from what the proper motion at the star's position would be if it were co-moving with Kepler 1627., on page 2, line 138.

Added: **in both physical and kinematic space**, on page 2, line 145.

Added: One reason to suspect this is that the spread in tangential velocities exceeds typical limits for kinematic coherence (Meingast et al. 2021), on page 2, line 151.

Replaced: **the** replaced with: **this**, on page 2, line 158.

Deleted: **s** on page 2, line 159.

Added: This also involved cuts on the photometric signal to noise ratio, the number of visibility periods used, the astrometric χ^2 of the single-source solution, and the $G_{\text{BP}} - G_{\text{RP}}$ color excess factor., on page 3, line 183.

Replaced: **agree reasonably well with** replaced with: **are within a factor of two of**, on page 3, line 199.

Replaced: **from** replaced with: **in**, on page 3, line 201.

Added: **, and are all small enough that the choice of whether to use them vs. other extinction estimates does not affect our primary conclusions**, on page 5, line 204.

Deleted: **hybrid** on page 5, line 234.

Replaced: **to** replaced with: **and**, on page 5, line 235.

Replaced: **s 6 and 7** replaced with: **6**, on page 5, line 236.

Added: To derive a probability distribution function for the age of δ Lyr cluster, we then assumed a Gaussian likelihood that treated the interpolated isochrones as the “model” and the δ Lyr cluster’s isochrone as the “data” (Equation 7 from Gagné et al. 2020). The cluster’s age and its statistical uncertainty are then quoted as the mean and standard deviation of this age posterior., on page 5, line 238.

Added: As a matter of scope, we restricted our attention to the 391 stars discussed in Section 2.1 in the spatial and kinematic proximity of Kepler 1627., on page 5, line 274.

Deleted: ~~As a matter of scope, we restricted our attention to the 391 stars discussed in Section 2.1 in the spatial and kinematic proximity of Kepler 1627.~~ on page 5, line 293.

Deleted: ~~Kepler rotation periods were derived by McQuillan et al. (2014) for 28 of our 391 stars; for these cases, we simply adopted the Kepler rotation period.~~ on page 5, line 297.

Added: First, to validate the TESS rotation periods, we compared against 28 stars from McQuillan et al. (2014) that were also observed by Kepler. Of the 23 stars with Kepler periods below 10 days, 21 of the TESS periods agreed with the Kepler rotation periods; the other 2 were measured at the double-period harmonic. Of the remaining 5 stars with Kepler rotation periods above 10 days, none were correctly recovered by TESS (3 were near the half-period harmonic). We therefore adopted a general stance of caution for TESS rotation periods exceeding 10 days, and used the Kepler rotation periods for the stars for which they were available., on page 5, line 300.

Added: **from each other**, on page 6, line 330.

Replaced: ~~The other stars are likely to be field contaminants.~~ replaced with: Although some of these outlier stars might be unresolved F+K binaries that are in the cluster (Stauffer et al. 2016), assuming that they are field contaminants provides a more secure lower bound of the rotation period detection fraction., on page 6, line 336.

Replaced: ~~renormalized unit weight error (RUWE)~~, replaced with: RUWE, on page 7, line 448.

Added: **(RUWE \approx 2.9; roughly the 98th percentile of the cluster’s distribution)**, on page 7, line 451.

Added: Given that the secondary companion’s brightness in the Kepler band is \approx 1.5% that of the primary, source confusion for the rotation signal is not expected to be an issue., on page 9, line 489.

Replaced: ~~mini-Neptune~~ replaced with: **Neptune**, on page 9, line 542.

Added: **second or fourth-order**, on page 10, line 598.

Added: **plus the transit**, on page 10, line 600.

Deleted: ~~We considered the results both for a second and fourth-order time-dependence in the local baseline~~, on page 10, line 601.

Added: **lack of**, on page 10, line 630.

Replaced: **main** replaced with: **prime**, on page 11, line 635.

Replaced: ~~simultaneously with Keek/HIRES and MuSCAT3~~ replaced with: spectroscopically with HIRES at the Keck-I telescope and photometrically in *griz* bands with MuSCAT3 at Haleakalā Observatory, on page 12, line 707.

Added: Details of the observation sequence are discussed in Appendix C; Figure 7 shows the results., on page 12, line 711.

Deleted: ~~. We scheduled the observations using the ephemeris of Holezer et al. (2016)~~, on page 12, line 713.

Deleted: ~~(Figure 7)~~ on page 12, line 718.

Added: The achromatic transits strongly favor Kepler 1627A as the transit host, since Kepler 1627B is much redder., on page 12, line 723.

Replaced: **precision** replaced with: **signal-to-noise**, on page 12, line 748.

Added: , given that the planetary orbital period (7.203 days) and the stellar rotation period (2.642 days) are not a rational combination, on page 12, line 780.

Replaced: **However**, replaced with: **However, it seems that**, on page 13, line 805.

Added: based on the per-target detection contours., on page 13, line 807.

Added: These limits account for the spot-induced variability in the system through a correction based on the Combined Differential Photometric Precision (CDPP) in the Kepler 1627 light curve over the relevant transit timescales (Burke & Catanzarite 2017), on page 13, line 811.

Deleted: ~~to $10 M_{\oplus}$~~ on page 13, line 819.

Added: This of course would need to be confirmed through a measurement of the planetary mass; alternative explanations for its large size could be that it is abnormally massive, or that it has an abnormally large envelope to core mass ratio., on page 13, line 823.

Added: Based on Figure 1, the δ Lyr cluster may also be part of a similar hierarchical association., on page 22.

Added: We used the ephemeris of Holczer et al. (2016) to observe a transit of Kepler 1627Ab on the night of 2021 Aug 7 both spectroscopically and photometrically. We used the HIRES echelle spectrograph at the Keck-I telescope and MuSCAT3 at Haleakalā Observatory on Maui, HI (Narita et al. 2020)., on page 25.

Deleted: ~~We monitored Kepler 1627 with Keek/HIRES before, during, and after transit on the night of 2021 Aug 7.~~ on page 25.

Replaced: ~~We used the iodine cell for wavelength calibration~~ replaced with: For the HIRES wavelength calibration, we used the iodine cell, on page 25.

Added: Given the faintness of the target ($V = 13.1$), we observed using the C2 decker, which yielded an instrument resolution of $\approx 50,000$., on page 25.

Added: The HIRES exposure time was set at ≈ 15 minutes in order to resolve the 2.8 hour transit event, which yielded a S/N per resolution element of ≈ 75 (low for precision radial velocity standards)., on page 25.

Replaced: ~~We also simultaneously observed across griz bands using MuSCAT3 at Haleakalā Observatory on Maui, HI.~~ replaced with: For the MuSCAT3 observations, we observed simultaneously across the *griz* bands., on page 25.

Added: The exposure times in each bandpass ranged between 23 and 46 seconds, and were chosen in order to yield a S/N in the peak pixel that exceeded 130 while also preventing saturation., on page 25.

Added: **planetary**, on page 28.