

## A 30 Million Year Old Mini-Neptune in the Kepler Field

L. G. BOUMA<sup>1</sup> AND J. L. CURTIS<sup>2,3</sup>

<sup>1</sup>*Department of Astrophysical Sciences, Princeton University, 4 Ivy Lane, Princeton, NJ 08540, USA*

<sup>2</sup>*Department of Astronomy, Columbia University, 550 West 120th Street, New York, NY 10027, USA*

<sup>3</sup>*Department of Astrophysics, American Museum of Natural History, Central Park West, New York, NY 10024, USA*

(Received —; Revised —; Accepted —)

Submitted to Nature

### ABSTRACT


Kepler 1627 is an active G8V star previously found to host a  $3.56 \pm 0.04 R_{\oplus}$  mini-Neptune on a 7.2 day orbit. The star was selected for observation by the Kepler satellite because it is nearby and it resembles the Sun. Combining data from the Gaia satellite with rotation periods from TESS, we show that Kepler 1627 is a member of the  $\delta$  Lyr cluster. Based on isochronal and gyrochronal evidence, the cluster is  $30 \pm XX$  Myr old. Based on archival NIRC2 imaging, we also note the existence of a low-mass stellar companion. Our analysis advances Kepler 1627Ab as the youngest planet with a precise age detected by the main Kepler mission. Newly derived ages from Gaia offer the opportunity to significantly expand the census of age-dated planets – we offer a literature concatenation of young stars to support this effort. The properties of Kepler 1627Ab may also help clarify how the orbits and atmospheres of the mini-Neptune planets evolve.

**Keywords:** planetary evolution (XXXX), stellar associations (1582), open star clusters (1160), stellar ages (1581),

### 1. MANUSCRIPT

While thousands of exoplanets have been discovered orbiting nearby stars, the vast majority of them are several billion years old. This results in difficulty testing theories for the origins of the different families of planets, since many expected evolutionary processes operate on timescales of less than 100 million years.


For instance, the “mini-Neptune” planets, thought to be made of rocky cores and extended atmospheric envelopes of hydrogen and helium, are expected to shrink in size by factors




**Figure 1. Kepler and TESS views of the  $\delta$  Lyr cluster.** Colored points are kinematically selected members of the  $\delta$  Lyr cluster (black points in Figure 5). Both Kepler (gray squares) and TESS (colored points) observed portions of the cluster. Naked-eye stars ( $m_V < 6.5$ ) are shown in gray; four of them (orange points) have their names annotated. Kepler 1627 (green star) was observed during the entirety of the Kepler mission, and has been observed by TESS for two lunar months to date.

of several over their first  $10^8$  years (Owen 2020). Specifically, they start with sizes of 4– $12 R_\oplus$  at the time of disk dispersal ( $\approx 10^7$  years), and shrink to sizes of 2– $4 R_\oplus$  by  $10^8$  years. This change is driven by a combination of stellar irradiation and internal heat that powers an outflow, which eventually depletes or entirely strips the envelope from the rocky core (Owen & Wu 2013; Gupta & Schlichting 2019). Discovering young planets and measuring their compositions and atmospheric outflows are key steps toward testing this paradigm, which is often invoked to explain the observed radius distribution of both mature exoplanets (Fulton et al. 2017), as well as younger planets (Bouma et al. 2020).

At the time of the main Kepler mission (2009–2013), only four open clusters were known in the Kepler field: NGC 6866, NGC 6811, NGC 6819, and NGC 6791, with ages spanning 0.7 Gyr to 9 Gyr (Meibom et al. 2011). Since that time, analyses of the kinematic, photometric, and astrometric Gaia data have expanded our knowledge of open cluster and moving group memberships (e.g., Cantat-Gaudin et al. 2018; Zari et al. 2018; Kounkel & Covey 2019; Meingast et al. 2021). As part of our Cluster Difference Imaging Photometric Survey (CDIPS, Bouma et al. 2019), we concatenated the available analyses from the literature, which yielded a list of candidate young and age-dated stars (see Section A).




**Figure 2.**  $\delta$  Lyr cluster is  $35 \pm 15$  Myr old. *Left:* TESS and Kepler rotations periods and dereddened Gaia colors, with the Pleiades (125 Myr; ?) and Praesepe (650 Myr; ?) shown for reference. Stars of a given color (mass) move up through time due to magnetic braking. The  $\delta$  Lyr cluster has a 40 Myr gyrochronal age, as calibrated against the LDB-age of IC 2602. *Right:* HR diagram of the same comparison. **todo:** caption improve.



**Figure 3.** Zoomed view of the Kepler 1627 light curve. The full Kepler dataset spans 1,437 days (3.9 years), sampled at 30 minute cadence. The *top panel* shows the PDCSAP median-subtracted flux in units of parts-per-thousand ( $\times 10^{-3}$ ). The dominant signal is induced by starspots and plages. The model for the stellar variability (orange line) is subtracted in the *bottom panel*, revealing the transits of Kepler 1627Ab, as well as other deviations from the stellar variability model. The Figure Set available online shows the entire 3.9 years of available data.

Comparing our young star list against the Kepler field yielded two discoveries. The first, to be discussed in an upcoming analysis by J. Curtis et al., is that Kepler observed the  $\approx 350$  Myr open cluster Theia 520 (UBC 1). At least five Kepler planets are known in the cluster, including the Kepler-52 and Kepler-968 systems. The second discovery, and the focus of this work, is that Kepler observed a portion of the  $\delta$  Lyr cluster, shown in Figure 1.



**Figure 4. Phase-folded transit of Kepler 1627b, with stellar variability removed.** The  $1-\sigma$  model uncertainties and the maximum *a posteriori* model are shown as the faint purple band, and the dark purple line. Gray points are individual PDCSAP flux measurements; black points are binned to 20 minutes, with a representative error bar shown. **LGB FIXME: seems like the uncertainty is bigger than the RMS in the out of transit...**

While we have yet to perform an in-depth planet search of the XXX light curves Kepler acquired for the  $\delta$  Lyr cluster, there is already a known planet in the cluster: Kepler 1627b. We considered the planet's previous statistical validation by Morton et al. (2016) to be sufficient cause to examine the age of the star and its cluster more closely. We collected the previously measured XXX Kepler rotation periods (CITEP MCQUILLAN2014), and measured YYY new stellar rotation periods for cluster members observed by TESS. We also performed an isochronal analysis. The results are shown in Figure 2. From the ensemble perspective, which provides the strongest constraints (SEE CITET SODERBLOM 2014), the age of the star is  $XX \pm YY$  Myr. Individual measurements of the star's rotation period and lithium abundance corroborate this conclusion.

If the Kepler 1627b transit signal truly corresponds to a planet, this age would make it the youngest planet yet found by the main Kepler mission. A portion of the light curve is shown in Figure 3, and the phase-folded transit is shown in Figure 4. The dominant signal in the PDCSAP photometry is a quasi-period starspot signal with a peak-to-peak amplitude that varies between roughly 2% and 8%, depending on the degree of asymmetry in spot coverage between the stellar hemispheres. The starspot signal repeats roughly every 2.606 days, though this value varies by a few percent depending on which portion of the

light curve is analyzed (**FIXME: do!**). The details of how we model this signal to derive the parameters of the planet are discussed in Section C. The results, given in Table 1, are consistent with a mini-Neptune sized planet ( $3.56 \pm 0.04 R_{\oplus}$ ) on a circular 7.2 day orbit, around a G8V star ( $0.91 \pm 0.05 R_{\odot}$ ) **fixme: age**.

Could the transit signal be produced by anything other than a planet orbiting this near-solar analog? Morton et al. (2016) considered the various false positive scenarios, including those of a foreground eclipsing binary, a hierarchical triple-star binary system, and a background eclipsing binary. Morton et al. (2016) reported the latter scenario to be the only possible case, with a (model-dependent) probability of  $\approx 10^{-5}$ . While this statistical validation is reassuring, “validated” planets have been demonstrated to be false positives (Shporer et al. 2017). Most failures of the statistical validation approach arise when attempting to separate the smallest stars from identically-sized gas giant planets (CITE). With a size of  $\approx 3.6 R_{\oplus}$ , Kepler 1627b does not meet this concern. The fact that we detect a low-mass stellar companion in archival Keck-NIRC2 Kp-band ( $2.12 \mu\text{m}$ ) imaging does not change the conclusion: the companion contributes 1% of the flux observed by Kepler, which combined with the observed transit shape cannot be invoked as an explanation for the observed data (Section D).

Kepler 1627Ab therefore provides an interesting extremum in the ages of the Kepler planets, and opens multiple avenues for detailed study. In order of ascending difficulty, observing its Doppler transit ( $\Delta v_{\text{RM}} \approx 20 \text{ m s}^{-1}$ ) with HIRES or the upcoming KPF spectrograph could yield a measurement of its orbital obliquity (CITEP,CITEP). (A simpler path to the same result might be to determine whether the  $\approx 80 \text{ ppm}$  residual apparent in our transit fit is caused by gravity darkening.) A transmission spectrum focused on either hydrogen or helium absorption could yield insight into its atmosphere, which is expected to be outflowing at a rate of  $\mathbf{X.X} M_{\oplus} \text{ yr}^{-1}$  (CITE). Finally, the most challenging measurement that may currently be in reach would be a measurement of its mass, through either detailed timing analyses of the Kepler data, or through a high-cadence multi-color radial velocity campaign. The mass would yield constraints on both the planet’s composition (do the mini-Neptunes form with gaseous envelopes?), as well as its initial entropy (“hot” or “cold”? CITE).

Ultimately, our age measurement for Kepler 6127Ab was enabled by identifying its connection to the  $\delta$  Lyr cluster through the Gaia data. The Supplementary Table 2 enables similar crossmatches to already known planets, as well as forthcoming planets from the TESS mission (CITE RickerJATIS2014). An alternative approach also likely to bear fruit is to identify kinematic associations around exoplanet host stars using the Gaia data, and to then verify these associations with ancillary rotation periods and spectroscopy (CITEP Tofflemire).

## ACKNOWLEDGMENTS

**Table 1.** Priors and posteriors for the transit and stellar variability model fitted to the long-cadence Kepler 1627b photometric timeseries.

Param.	Unit	Prior	Median	Mean	Std. Dev.	3%	97%	ESS	$\hat{R}-1$
<i>Sampled</i>									
$P$	d	$\mathcal{N}(7.20281; 0.01000)$	7.2028035	7.2028033	0.0000073	7.2027893	7.2028171	1910.5903732	0.0035905
$t_0^{(1)}$	d	$\mathcal{N}(120.79053; 0.02000)$	120.790504	120.790505	0.0009438	120.7886867	120.7922431	1564.1105056	0.0003213
$\log R_p/R_\star$	–	$\mathcal{U}(-4.605; 0.000)$	-3.33523	-3.33569	0.06618	-3.45772	-3.21617	1173.56574	0.00310
$b$	–	$\mathcal{U}(0; 1+R_p/R_\star)$	0.3971	0.3886	0.2070	0.0204	0.7289	378.9528	0.0177
$u_1$	–	Kipping (2013)	0.28	0.30	0.179	0.002	0.603	1161.95	0.005
$u_2$	–	Kipping (2013)	0.425	0.381	0.314	-0.197	0.912	900.884	0.002
$R_\star$	$R_\odot$	$\mathcal{T}(0.910; 0.052)$	0.911	0.910	0.051	0.814	1.004	2265.830	-0.001
$\log g$	cgs	$\mathcal{N}(4.600; 0.100)$	4.604	4.601	0.094	4.417	4.769	923.943	0.
$\langle f \rangle$	–	$\mathcal{N}(0.500; 0.100)$	0.4999	0.4999	0.0003	0.4993	0.5005	2964.6676	0.0013
$e^{(2)}$	–	Van Eylen et al. (2019)	0.127	0.168	0.147	0.	0.446	518.492	0.004
$\omega$	rad	$\mathcal{U}(0.000; 6.283)$	-0.235	-0.170	1.867	-2.879	3.132	1212.260	0.006
$\log \sigma_f$	–	$\mathcal{N}(\log \langle \sigma_f \rangle; 2.000)$	-8.016	-8.016	0.008	-8.031	-8.001	2224.427	-0.
$\rho$	d	$\mathcal{U}(1.000; 10.000)$	2.953	2.955	0.096	2.777	3.131	1936.492	0.
$\sigma$	$d^{-1}$	InvGamma(1.000; 5.000)	0.013	0.013	0.001	0.012	0.014	2155.687	-0.
$\sigma_{\text{rot}}$	$d^{-1}$	InvGamma(1.000; 5.000)	0.897	0.933	0.222	0.552	1.316	2147.324	0.003
$\log P_{\text{rot}}$	log(d)	$\mathcal{N}(0.958; 0.020)$	0.964	0.964	0.001	0.963	0.966	2486.029	0.
$\log Q_0$	–	$\mathcal{N}(0.000; 2.000)$	12.935	12.960	0.454	12.157	13.857	2126.581	0.004
$\log dQ$	–	$\mathcal{N}(0.000; 2.000)$	0.029	0.021	2.032	-3.785	3.780	1755.941	0.
$f$	–	$\mathcal{U}(0.100; 1.000)$	0.111	0.113	0.010	0.1	0.130	1253.358	-0.0
<i>Derived</i>									
$R_p/R_\star$	–	–	0.036	0.036	0.002	0.031	0.040	1173.566	0.003
$\rho_\star$	$\text{g cm}^{-3}$	–	2.27	2.312	0.516	1.408	3.272	910.746	0.001
$R_p$	$R_{\text{Jup}}$	–	0.316	0.317	0.037	0.251	0.386	1692.736	0.001
$a/R_\star$	–	–	18.396	18.407	1.375	15.690	20.780	910.717	0.001
$\cos i$	–	–	0.022	0.021	0.011	0.002	0.038	439.303	0.011
$T_{14}$	hr	–	2.822	2.823	0.057	2.710	2.917	1086.008	0.002
$T_{13}$	hr	–	2.578	2.566	0.083	2.430	2.714	590.630	0.011

NOTE— ESS refers to the number of effective samples.  $\hat{R}$  is the Gelman-Rubin convergence diagnostic. Logarithms through this table are in base- $e$ .  $\mathcal{U}$  denotes a uniform distribution,  $\mathcal{N}$  a normal distribution, and  $\mathcal{T}$  a truncated normal bounded between zero and an upper limit much larger than the mean. (1) The ephemeris is in units of BJDTDB - 2454833. (2) The eccentricity vectors are sampled in the  $(e \cos \omega, e \sin \omega)$  basis.

L.G.B. and J.L.C. are grateful to T. David for help with the transit fitting. L.G.B. acknowledges support by the TESS GI Program, programs G011103 and G022117, through NASA grants 80NSSC19K0386 and 80NSSC19K1728. L.G.B. was also supported by a Charlotte Elizabeth Procter Fellowship from Princeton University. This study was based in part on observations at Cerro Tololo Inter-American Observatory at NSF’s NOIRLab (NOIRLab Prop. ID 2020A-0146; 2020B-0029 PI: Bouma), which is managed by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation. This paper also includes data collected by the TESS mission, which are publicly available from the Mikulski Archive for Space Telescopes (MAST). Funding for the TESS mission is provided by NASA’s Science Mission Directorate. We thank the TESS Architects (G. Ricker, R. Vanderspek, D. Latham, S. Seager, J. Jenkins) and the many TESS team members for their efforts to make the mission a continued success.

*Software:* astrobase (Bhatti et al. 2018), astropy (Astropy Collaboration et al. 2018), astroquery (Ginsburg et al. 2018), corner (Foreman-Mackey 2016), exoplanet (Foreman-Mackey et al. 2020), and its dependencies (Agol et al. 2020; Kipping 2013; Luger et al. 2019; Theano Development Team 2016), IPython (Pérez & Granger 2007), matplotlib (Hunter 2007), numpy (Walt et al. 2011), pandas (McKinney 2010), PyMC3 (Salvatier et al. 2016), scipy (Jones et al. 2001), TESS-point (Burke et al. 2020), wotan (Hippke et al. 2019).

*Facilities:* *Astrometry:* Gaia (Gaia Collaboration et al. 2018b, 2020). *Imaging:* Second Generation Digitized Sky Survey. *Spectroscopy:* CTIO1.5m (CHIRON; Tokovinin et al. 2013), AAT (HERMES; Lewis et al. 2002; Sheinis et al. 2015), VLT:Kueyen (FLAMES; Pasquini et al. 2002). *Photometry:* TESS (Ricker et al. 2015).

## REFERENCES

- Agol, E., Luger, R., & Foreman-Mackey, D. 2020, *AJ*, **159**, 123
- Akeson, R. L., Chen, X., Ciardi, D., et al. 2013, *PASP*, **125**, 989
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, **156**, 123
- Bhatti, W., Bouma, L. G., & Wallace, J. 2018, astrobase, <https://doi.org/10.5281/zenodo.1469822>
- Boisse, I., Pepe, F., Perrier, C., et al. 2012, *A&A*, **545**, A55
- Bouma, L. G., Hartman, J. D., Bhatti, W., Winn, J. N., & Bakos, G. Á. 2019, *ApJS*, **245**, 13
- Bouma, L. G., Hartman, J. D., Brahm, R., et al. 2020, *AJ*, **160**, 239
- Burke, C. J., Levine, A., Fausnaugh, M., et al. 2020, TESS-Point: High precision TESS pointing tool, Astrophysics Source Code Library, [ascl:2003.001](https://ascl.net/2003.001)
- Cantat-Gaudin, T., & Anders, F. 2020, *A&A*, **633**, A99
- Cantat-Gaudin, T., Jordi, C., Vallenari, A., et al. 2018, *A&A*, **618**, A93
- Cantat-Gaudin, T., Jordi, C., Vallenari, A., et al. 2018, *A&A*, **618**, A93
- Cantat-Gaudin, T., Jordi, C., Wright, N. J., et al. 2019, *A&A*, **626**, A17
- Cantat-Gaudin, T., Anders, F., Castro-Ginard, A., et al. 2020, *A&A*, **640**, A1
- Castro-Ginard, A., Jordi, C., Luri, X., et al. 2020, *A&A*, **635**, A45
- Cotten, T. H., & Song, I. 2016, *ApJS*, **225**, 15
- Damiani, F., Prisinzano, L., Pillitteri, I., Micela, G., & Sciortino, S. 2019, *A&A*, **623**, A112
- Dias, W. S., Monteiro, H., Caetano, T. C., et al. 2014, *Astronomy and Astrophysics*, **564**, A79
- Esplin, T. L., & Luhman, K. L. 2019, *AJ*, **158**, 54
- Foreman-Mackey, D. 2016, *Journal of Open Source Software*, **1**, 24
- Foreman-Mackey, D., Czekala, I., Luger, R., et al. 2020, exoplanet-dev/exoplanet v0.2.6
- Fulton, B. J., Petigura, E. A., Howard, A. W., et al. 2017, *AJ*, **154**, 109
- Fürnkranz, V., Meingast, S., & Alves, J. 2019, *A&A*, **624**, L11
- Gagné, J., David, T. J., Mamajek, E. E., et al. 2020, *ApJ*, **903**, 96
- Gagné, J., & Faherty, J. K. 2018, *ApJ*, **862**, 138
- Gagné, J., Roy-Loubier, O., Faherty, J. K., Doyon, R., & Malo, L. 2018a, *ApJ*, **860**, 43
- Gagné, J., Mamajek, E. E., Malo, L., et al. 2018b, *ApJ*, **856**, 23
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2020, arXiv e-prints, arXiv:2012.01533
- Gaia Collaboration, Babusiaux, C., van Leeuwen, F., et al. 2018a, *A&A*, **616**, A10
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018b, *A&A*, **616**, A1
- Ginsburg, A., Sipocz, B., Madhura Parikh, et al. 2018, Astropy/Astroquery: V0.3.7 Release
- Goldman, B., Röser, S., Schilbach, E., Moór, A. C., & Henning, T. 2018, *ApJ*, **868**, 32
- Gupta, A., & Schlichting, H. E. 2019, *MNRAS*, **487**, 24
- Hippke, M., David, T. J., Mulders, G. D., & Heller, R. 2019, *AJ*, **158**, 143
- Hunter, J. D. 2007, Computing in Science & Engineering, **9**, 90
- Jones, E., Oliphant, T., Peterson, P., et al. 2001, Open source scientific tools for Python
- Kharchenko, N. V., Piskunov, A. E., Schilbach, E., Röser, S., & Scholz, R.-D. 2013, *A&A*, **558**, A53
- Kipping, D. M. 2013, *MNRAS*, **435**, 2152
- Kounkel, M., & Covey, K. 2019, *AJ*, **158**, 122
- Kounkel, M., & Covey, K. 2019, *AJ*, **158**, 122
- Kounkel, M., Covey, K., & Stassun, K. G. 2020, *AJ*, **160**, 279
- Kounkel, M., Covey, K., Suárez, G., et al. 2018, *AJ*, **156**, 84
- Kounkel, M., Covey, K., Suárez, G., et al. 2018, *AJ*, **156**, 84
- Kraus, A. L., Shkolnik, E. L., Allers, K. N., & Liu, M. C. 2014, *AJ*, **147**, 146
- Lewis, I. J., Cannon, R. D., Taylor, K., et al. 2002, *MNRAS*, **333**, 279
- Luger, R., Agol, E., Foreman-Mackey, D., et al. 2019, *AJ*, **157**, 64
- Mann, A. W., Johnson, M. C., Vanderburg, A., et al. 2020, *AJ*, **160**, 179

- McKinney, W. 2010, in Proceedings of the 9th Python in Science Conference, ed. S. van der Walt & J. Millman, 51
- Meibom, S., Barnes, S. A., Latham, D. W., et al. 2011, *The Astrophysical Journal Letters*, 733, L9
- Meingast, S., & Alves, J. 2019, *A&A*, 621, L3
- Meingast, S., Alves, J., & Rottensteiner, A. 2021, *A&A*, 645, A84
- Morton, T. D., Bryson, S. T., Coughlin, J. L., et al. 2016, *ApJ*, 822, 86
- Oh, S., Price-Whelan, A. M., Hogg, D. W., Morton, T. D., & Spergel, D. N. 2017, *AJ*, 153, 257
- Owen, J. E. 2020, arXiv:2009.03919 [astro-ph], arXiv: 2009.03919
- Owen, J. E., & Wu, Y. 2013, *ApJ*, 775, 105
- Pasquini, L., Avila, G., Blecha, A., et al. 2002, *The Messenger*, 110, 1
- Pavlidou, T., Scholz, A., & Teixeira, P. S. 2021, *MNRAS*, 503, 3232
- Pecaut, M. J., & Mamajek, E. E. 2013, *ApJS*, 208, 9
- Pérez, F., & Granger, B. E. 2007, *Computing in Science and Engineering*, 9, 21
- Ratzenböck, S., Meingast, S., Alves, J., Möller, T., & Bomze, I. 2020, *A&A*, 639, A64
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, *Journal of Astronomical Telescopes, Instruments, and Systems*, 1, 014003
- Rizzuto, A. C., Mann, A. W., Vanderburg, A., Kraus, A. L., & Covey, K. R. 2017, *AJ*, 154, 224
- Roccatagliata, V., Franciosini, E., Sacco, G. G., Randich, S., & Sicilia-Aguilar, A. 2020, *A&A*, 638, A85
- Röser, S., & Schilbach, E. 2020, *A&A*, 638, A9
- Salvatier, J., Wieckiâ, T. V., & Fonnesbeck, C. 2016, PyMC3: Python probabilistic programming framework
- Sheinis, A., Anguiano, B., Asplund, M., et al. 2015, *Journal of Astronomical Telescopes, Instruments, and Systems*, 1, 035002
- Shporer, A., Zhou, G., Vanderburg, A., et al. 2017, *ApJL*, 847, L18
- Theano Development Team. 2016, arXiv e-prints, abs/1605.02688
- Tian, H.-J. 2020, *ApJ*, 904, 196
- Tokovinin, A., Fischer, D. A., Bonati, M., et al. 2013, *PASP*, 125, 1336
- Ujjwal, K., Kartha, S. S., Mathew, B., Manoj, P., & Narang, M. 2020, *AJ*, 159, 166
- Van Eylen, V., Albrecht, S., Huang, X., et al. 2019, *AJ*, 157, 61
- Villa Vélez, J. A., Brown, A. G. A., & Kenworthy, M. A. 2018, *Research Notes of the American Astronomical Society*, 2, 58
- Walt, S. v. d., Colbert, S. C., & Varoquaux, G. 2011, Computing in Science & Engineering, 13, 22
- Wenger, M., Ochsenbein, F., Egret, D., et al. 2000, *A&AS*, 143, 9
- Zari, E., Hashemi, H., Brown, A. G. A., Jardine, K., & de Zeeuw, P. T. 2018, *A&A*, 620, A172
- Zari, E., Hashemi, H., Brown, A. G. A., Jardine, K., & de Zeeuw, P. T. 2018, *A&A*, 620, A172

**Table 2.** Young, Age-dated, and Age-dateable Stars Within the Nearest Few Kiloparsecs (v0.5 of the CDIPS Target List).

Parameter	Example Value	Description
source_id	1709456705329541504	Gaia DR2 source identifier.
ra	247.826	Gaia DR2 right ascension [deg].
dec	79.789	Gaia DR2 declination [deg].
parallax	35.345	Gaia DR2 parallax [mas].
parallax_error	0.028	Gaia DR2 parallax uncertainty [mas].
pmra	94.884	Gaia DR2 proper motion $\mu_\alpha \cos \delta$ [mas yr $^{-1}$ ].
pmdec	-86.971	Gaia DR2 proper motion $\mu_\delta$ [mas yr $^{-1}$ ].
phot_g_mean_mag	6.85	Gaia DR2 $G$ magnitude.
phot_bp_mean_mag	6.409	Gaia DR2 $G_{\text{BP}}$ magnitude.
phot_rp_mean_mag	7.189	Gaia DR2 $G_{\text{RP}}$ magnitude.
cluster	Uma,IR_excess,NASAExoArchive_ps_20210506	Comma-separated cluster or group name.
age	nan,nan,9.48	Comma-separated logarithm (base-10) of reported <sup>a</sup> age in years.
mean_age	9.48	Mean (ignoring NaNs) of age column.
reference_id	Ujjwal2020,CottenSong2016,NASAExoArchive_ps_20210506	Comma-separated provenance of group membership.
reference_bibcode	2020AJ....159..166U,2016ApJS..225...15C,2013PASP..125..989A	ADS bibcode corresponding to reference_id.

NOTE— Table 2 is published in its entirety in a machine-readable format. This table is a concatenation of the studies listed in Table 3. One entry is shown for guidance regarding form and content. In this particular example, the star has a cold Jupiter on a 16 year orbit, HD 150706b (Boisse et al. 2012). An infrared excess has been reported (Cotten & Song 2016), and the star was identified by Ujjwal et al. (2020) as a candidate UMa moving group member ( $\approx 400$  Myr; Mann et al. 2020). The star’s RV activity and TESS rotation period corroborate its youth.

## APPENDIX

### A. YOUNG, AGE-DATED, AND AGE-DATEABLE STAR COMPILATION

The v0.5 CDIPS target catalog (Table 2) includes some important updates from previous versions. As in Bouma et al. (2019), we collected membership information for young, age-dated, or age-dateable stars from across the literature. Table 3 gives a list of the sources included, and some brief summary statistics.

The first major important change is that the extent of analyses performed on the Gaia data at the time of our compilation was wide and deep enough that we opted to neglect pre-Gaia analyses, except in cases for which spectroscopically confirmed samples of stars had been collected. The membership lists for instance of Kharchenko et al. (2013) and Dias et al. (2014) (MWSC and DAML) were no longer required, especially given their relatively high field-star contamination rates compared to Gaia-derived membership catalogs.

For any of the catalogs for which Gaia DR2 identifiers were not immediately available, we either followed the spatial (plus proper-motion) crossmatching procedures described in Bouma et al. (2019), or else we pulled the Gaia DR2 source identifiers associated with the catalog from SIMBAD. We consequently opted to drop the ext\_catalog\_name and dist columns maintained in Bouma et al. (2019), as these were only populated for a small number of stars.

The most crucial parameters of a given star for our purposes are the Gaia DR2 source identifier (source\_id), the cluster name (cluster), and the (age). Given the hierarchical nature of many stellar associations, we do not attempt to resolve the cluster names to

**Table 3.** Provenances of Young and Age-Dateable Stars.

Reference	$N_{\text{Gaia}}$	$N_{\text{Age}}$	$N_{G_{\text{RP}} < 16}$
Kounkel et al. (2020)	987376	987376	775363
Cantat-Gaudin & Anders (2020)	433669	412671	269566
Cantat-Gaudin et al. (2018)	399654	381837	246067
Kounkel & Covey (2019)	288370	288370	229506
Cantat-Gaudin et al. (2020)	233369	227370	183974
Zari et al. (2018) UMS	86102	0	86102
Wenger et al. (2000) Y*?	61432	0	45076
Zari et al. (2018) PMS	43719	0	38435
Gaia Collaboration et al. (2018a) $d > 250 \text{ pc}$	35506	31182	18830
Castro-Ginard et al. (2020)	33635	24834	31662
Wenger et al. (2000) Y*O	28406	0	16205
Villa Vélez et al. (2018)	14459	14459	13866
Cantat-Gaudin et al. (2019)	11843	11843	9246
Damiani et al. (2019) PMS	10839	10839	9901
Oh et al. (2017)	10379	0	10370
Meingast et al. (2021)	7925	7925	5878
Wenger et al. (2000) pMS*	5901	0	3006
Gaia Collaboration et al. (2018a) $d > 250 \text{ pc}$	5378	817	3968
Kounkel et al. (2018)	5207	3740	5207
Ratzenböck et al. (2020)	4269	4269	2662
Wenger et al. (2000) TT*	4022	0	3344
Damiani et al. (2019) UMS	3598	3598	3598
Rizzuto et al. (2017)	3294	3294	2757
Akeson et al. (2013)	3107	868	3098
Tian (2020)	1989	1989	1394
Goldman et al. (2018)	1844	1844	1783
Cotten & Song (2016)	1695	0	1693
Gagné et al. (2018b)	1429	0	1389
Röser & Schilbach (2020) Psc-Eri	1387	1387	1107
Röser & Schilbach (2020) Pleiades	1245	1245	1019
Wenger et al. (2000) TT?	1198	0	853
Gagné & Faherty (2018)	914	0	913
Pavlidou et al. (2021)	913	913	504
Gagné et al. (2018a)	692	0	692
Ujjwal et al. (2020)	563	0	563
Gagné et al. (2020)	566	566	351
Esplin & Luhman (2019)	377	443	296
Roccatagliata et al. (2020)	283	283	232
Meingast & Alves (2019)	238	238	238
Fürnkranz et al. (2019) Coma-Ber	214	214	213
Fürnkranz et al. (2019) Neighbor Group	177	177	167
Kraus et al. (2014)	145	145	145

NOTE—Table 3 describes the provenances for the young and age-dateable stars in Table 2.  $N_{\text{Gaia}}$ : number of Gaia stars we parsed from the literature source.  $N_{\text{Age}}$ : number of stars in the literature source with ages reported.  $N_{G_{\text{RP}} < 16}$ : number of Gaia stars we parsed from the literature source with either  $G_{\text{RP}} < 16$ , or a parallax S/N exceeding 5 and a distance closer than 100 pc. The latter criterion included a few hundred white dwarfs that would have otherwise been neglected. Some studies appear multiple times when multiple tables from the analysis were included in the concatenation.

a single unique string. The Orion complex for instance, can be divided into almost one hundred kinematic subgroups (Kounkel et al. 2018). Similar complexity applies to the problem of determining homogeneous ages, which we do not attempt to resolve. Instead, we simply merged the cluster names and ages reported by various authors together.

This means that our “age” column can be null, for cases in which the original authors did not report an age, and a reference literature age was not readily available. Nonetheless, since we do generally prefer stars with known ages, we made a few additional efforts to populate this column. When available, the age provenance is from the original analysis of the cluster. However, in a few cases we adopted other ages when the string-based crossmatches on the “cluster” name was straightforward. In particular, we used the ages determined by Cantat-Gaudin et al. (2020) to assign ages to the catalogs from Gaia Collaboration et al. (2018a), Cantat-Gaudin et al. (2018), Castro-Ginard et al. (2020), and Cantat-Gaudin & Anders (2020).

The catalogs we included for which ages were not immediately available were those of Cotten & Song (2016), Oh et al. (2017), Zari et al. (2018), Gagné et al. (2018b), Gagné et al. (2018a), Gagné & Faherty (2018), and Ujjwal et al. (2020). While in principle the moving group members discussed by Gagné et al. (2018b,a); Gagné & Faherty (2018) and Ujjwal et al. (2020) have easily associated ages, our SIMBAD cross-matching lost the moving group association from those studies, which should therefore be recovered tools such as BANYAN  $\Sigma$ .<sup>1</sup>. We also included the SIMBAD object identifiers TT\*, Y\*O,Y\*?, TT?, and pMS\*. Finally, we also included every star in the NASA Exoplanet Archive ps table that had a Gaia identifier available (Akeson et al. 2013). If the age had finite uncertainties, we also included it, since stellar ages determined through the combination of isochrone-fitting and transit-derived stellar densities typically have higher precision than from isochrones alone.


The technical manipulations for the merging, cleaning, and joining were performed using pandas (McKinney 2010). The eventual crossmatch (using the Gaia DR2 source\_id) against the Gaia DR2 archive was performed asynchronously on the Gaia archive website<sup>2</sup>.

## B. KINEMATIC SELECTION OF $\delta$ LYR CLUSTER MEMBERS

Figure 5 shows members of the  $\delta$  Lyr cluster reported by Kounkel & Covey (2019) to be in the group. Galactic positions are determined and plotted only for stars with parallax signal-to-noise exceeding 20. The location of the Sun is shown on the plots. The non-uniform “clumps” might be an artifact of the data processing steps performed by Kounkel & Covey (2019). We therefore only consider stars in the immediate kinematic group around Kepler 1627. The tangential velocities relative to Kepler 1627 are shown in the bottom right panel. These are computed by assuming that every star has the same three-dimensional spatial velocity as Kepler 1627, where we assume a systemic radial velocity of  $-16.7 \pm 0.2$  km s<sup>-1</sup> based on the reconnaissance spectra obtained by A. Howard on HIRES

<sup>1</sup> <http://www.exoplanetes.umontreal.ca/banyan/banyansigma.php>

<sup>2</sup> <https://gea.esac.esa.int/archive/>



**Figure 5. Galactic position and tangential velocities of the  $\delta$  Lyr cluster (also known as Theia 73 and Stephenson 1).** Points are candidate cluster members with  $\bar{\omega}/\sigma_{\bar{\omega}} > 20$ , reported to be in the group by Kounkel & Covey (2019). We focus on stars in a small region (black points) in the kinematic vicinity of Kepler 1627 (yellow star). The other candidate cluster members (gray points) may or may not share the ages of the selected kinematic group. The location of the Sun is ( $\odot$ ) is shown.

and D. Latham on TRES. The relevant projection effects are then taken into account, as discussed by *e.g.*, Meingast et al. (2021) and L. Bouma et al (2021, submitted).


### C. TRANSIT AND STELLAR VARIABILITY MODEL

We assumed a quadratic limb-darkening law, with the uninformative prior advocated by CITET Kipping2013.

Figure 4 shows X, Y, Z. The residuals during the transit may hint at some small degree of unfitted signal – in the sense that the observations are systematically high in the first half of transit, and low in the second half. One possible explanation could be that we are seeing the effects of gravity darkening (*e.g.*, CITE Masuda 2015). The expected amplitude of the effect is XXX, based on CITE. However, there is also a dip of comparable amplitude shortly before the transit. Given the uncertainties in the stellar variability model, we expect that these patterns may correspond to more noise than signal.

### D. COMPANION STAR AND FALSE POSITIVE ASSESSMENT

Our analysis of archival Keck-NIRC2 K<sub>p</sub>-band (2.12  $\mu$ m) imaging revealed the existence of a previously unreported stellar neighbor, unresolved in the Gaia source catalog **FIXME: uncertainties**. The NIRC2 images yield a projected separation  $\rho = 0.^{\circ}17$ , with  $\Delta K_p = 2.5$ . Using the measured Gaia EDR3 parallax for the system, this implies a projected



**Figure 6.** Gaia EDR3 renormalized unit weight error (RUWE) point estimates for Kepler 1627A and other members of the  $\delta$  Lyr cluster. Since other members of the cluster with similar colors have comparable degrees of photometric variability, the high RUWE estimate suggests that Kepler 1627A is a binary.



**Figure 7.** Keck NIRC2 AO image of Kepler 1627A and Kepler 1627B. SCALEBAR DE-NOTES... North is up, East is left.

separation of 54 AU. The presence of this star is consistent with the excess noise in the Gaia astrometric time-series (see Section D). Given the low chance of a star imaged within **X.X arcseconds** to be a chance companion along the line of sight, we proceed under the assumption that it is bound, and that the Kepler 1627 system is binary.

Unforuntately, we do not have any reliable color information about Kepler 1627B. Based on the tabulation<sup>3</sup> by [Pecaut & Mamajek \(2013\)](#), the measured NIR-contrast for a main sequence G8V star corresponds to a spectral type for the companion of  $\approx$ M2V ( $M_* \approx 0.44M_\odot$ ). However, the companion should have a longer pre-main-sequence contraction phase than the primary, which would imply that this mass is overestimated by  $\sim 10$  to 20%.

Could the companion be creating a false positive signal? The companion star contributes  $\approx 1\%$  of the total flux observed in the Kepler aperture (0.7-2.0% V-band to G-band dmag difference from EEM table; todo is fix using model spectra). The observed transit has a depth of 0.18%. An 18% deep eclipse of the secondary star would therefore be needed to produce a deep enough signal. The shape of the observed signal requires allowed impact parameters to span 0.02 to 0.73 (Table 1); the body transiting the secondary would therefore need to be non-grazing with  $R_3/R_2 \approx 0.42$ . Assuming a  $\approx 0.42R_\odot$  radius of the imaged secondary, this would imply a tertiary stellar radius of  $\approx 0.2R_\odot$ . This scenario ultimately yields a contradiction, because it would require an ingress and egress phase that each span  $\approx 40\%$  of the transit duration ( $\approx 65$  minutes). The actual measured ingress and egress duration is  $\approx 15$  minutes), 4.4× shorter.

*Stellar density implied by transit duration*—The duration of the transit ( $2.823 \pm 0.057\text{hr}$ ) and the implied stellar density ( $2.31 \pm 0.52\text{ g cm}^{-3}$ ) could in theory help rule between whether the transiting body orbits the primary or secondary star. Ultimately, our stellar density measurement is not precise enough to render the blend scenario implausible. At 30 Myr, a  $0.40M_\odot$  solar-metallicity dwarf is  $\approx 26\%$  larger than when it is fully contracted on the main sequence ( $0.40R_\odot$  vs.  $0.50R_\odot$ ; CITEALT: Choi et al, MIST grids). The theoretically implied companion density of  $3.2\text{ g cm}^{-3}$  is indeed larger than the primary star's density of  $1.80\text{ g cm}^{-3}$  (measured through the HIRES reconnaissance spectroscopy). However the stellar density measured from the transit fitting ( $2.31 \pm 0.52\text{ g cm}^{-3}$ ) is only discrepant at the  $\approx 2\text{-}\sigma$  level from the M-dwarf blend scenario. It is instead the combination of the flux contrast, transit depth, and ingress duration that rule out this scenario.

<sup>3</sup> [http://www.pas.rochester.edu/~emamajek/EEM\\_dwarf\\_UBVIJHK\\_colors\\_Teff.txt](http://www.pas.rochester.edu/~emamajek/EEM_dwarf_UBVIJHK_colors_Teff.txt), version 2021/03/02.