

# Class 9: Halloween Mini Project

Linh Dang (PID: A16897764)

## Table of contents

Data import . . . . .	1
What is your favorite candy? . . . . .	2
Exploratory Analysis . . . . .	3
3. Overall Candy Rankings . . . . .	6
4. Winpercent vs Pricepercent . . . . .	13
5. Correlation Structure . . . . .	14
Principal Component Analysis (PCA) . . . . .	16

Today we will take a wee step back to some data we can taste and explore the correlation structure and principal components of some Halloween candy.

## Data import

```
candy_file <- "candy-data.csv"

candy = read.csv(candy_file, row.names=1)
head(candy)
```

	chocolate	fruity	caramel	peanutyalmondy	nougat	crispedricewafer
100 Grand	1	0	1	0	0	1
3 Musketeers	1	0	0	0	1	0
One dime	0	0	0	0	0	0
One quarter	0	0	0	0	0	0
Air Heads	0	1	0	0	0	0
Almond Joy	1	0	0	1	0	0

	hard	bar	pluribus	sugarpercent	pricepercent	winpercent
100 Grand	0	1	0	0.732	0.860	66.97173

3 Musketeers	0	1	0	0.604	0.511	67.60294
One dime	0	0	0	0.011	0.116	32.26109
One quarter	0	0	0	0.011	0.511	46.11650
Air Heads	0	0	0	0.906	0.511	52.34146
Almond Joy	0	1	0	0.465	0.767	50.34755

Q1. How many different candy types are in this dataset?

```
nrow(candy)
```

```
[1] 85
```

Q2. How many fruity candy types are in the dataset?

```
sum(candy$fruity)
```

```
[1] 38
```

**What is your favorite candy?**

```
candy["Twix",]$winpercent
```

```
[1] 81.64291
```

Q3. What is your favorite candy in the dataset and what is it's winpercent value?

My favorite candy is Twix with a winpercent of 81.6%.

Q4. What is the winpercent value for "Kit Kat"?

```
candy["Kit Kat",]$winpercent
```

```
[1] 76.7686
```

Q5. What is the winpercent value for "Tootsie Roll Snack Bars"?

```
candy["Tootsie Roll Snack Bars",]$winpercent
```

```
[1] 49.6535
```

## Exploratory Analysis

We can use the **skimr** package to get a quick overview of a given dataset. This can be useful for the first time you encounter a new dataset.

```
skimr::skim(candy)
```

Table 1: Data summary

Name	candy
Number of rows	85
Number of columns	12
<hr/>	
Column type frequency: numeric	12
<hr/>	
Group variables	None

### Variable type: numeric

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100	hist
chocolate	0	1	0.44	0.50	0.00	0.00	0.00	1.00	1.00	
fruity	0	1	0.45	0.50	0.00	0.00	0.00	1.00	1.00	
caramel	0	1	0.16	0.37	0.00	0.00	0.00	0.00	1.00	
peanutyalmondy	0	1	0.16	0.37	0.00	0.00	0.00	0.00	1.00	
nougat	0	1	0.08	0.28	0.00	0.00	0.00	0.00	1.00	
crispedricewafer	0	1	0.08	0.28	0.00	0.00	0.00	0.00	1.00	
hard	0	1	0.18	0.38	0.00	0.00	0.00	0.00	1.00	
bar	0	1	0.25	0.43	0.00	0.00	0.00	0.00	1.00	
pluribus	0	1	0.52	0.50	0.00	0.00	1.00	1.00	1.00	
sugarpercent	0	1	0.48	0.28	0.01	0.22	0.47	0.73	0.99	
pricepercent	0	1	0.47	0.29	0.01	0.26	0.47	0.65	0.98	
winpercent	0	1	50.32	14.71	22.45	39.14	47.83	59.86	84.18	

Q6. Is there any variable/column that looks to be on a different scale to the majority of the other columns in the dataset?


It looks like the last column `candy$winpercent` is on a different scale to all others.

Q7. What do you think a zero and one represent for the `candy$chocolate` column?

A zero means that candy has chocolate while one means that the candy has no chocolate in it.


Q8. Plot a histogram of winpercent values

```
hist(candy$winpercent, breaks = 30)
```



```
library(ggplot2)

ggplot(candy) +
  aes(winpercent) +
  geom_histogram(bins = 10, fill="lightblue")
```



Q9. Is the distribution of winpercent values symmetrical?

The distribution of winpercent values is not symmetrical.

Q10. Is the center of the distribution above or below 50%?

```
summary(candy$winpercent)
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
22.45	39.14	47.83	50.32	59.86	84.18

The center of the distribution is below 50% because median is 47.83.

Q11. On average is chocolate candy higher or lower ranked than fruit candy?

```
choc.inds <- candy$chocolate == 1
choc.candy <- candy[ choc.inds, ]
choc.win <- choc.candy$winpercent
mean(choc.win)
```

```
[1] 60.92153
```

```
fruit.win <- candy[ as.logical(candy$fruity), ]$winpercent
mean(fruit.win)
```

```
[1] 44.11974
```

On average, chocolate candy is higher than fruit candy.

Q12. Is this difference statistically significant?

```
ans <- t.test(choc.win, fruit.win)
ans
```

Welch Two Sample t-test

```
data:  choc.win and fruit.win
t = 6.2582, df = 68.882, p-value = 2.871e-08
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 11.44563 22.15795
sample estimates:
mean of x mean of y
 60.92153  44.11974
```

Yes, with a P-value of  $2.8713778 \times 10^{-8}$ .

### 3. Overall Candy Rankings

There are two related functions that can help here, one is the classic `sort()` and `order()`

```
x <- c(5, 10, 1, 4)
sort(x)
```

```
[1] 1 4 5 10
```

```
order(x)
```

```
[1] 3 4 1 2
```

Q13. What are the five least liked candy types in this set?

```
inds <- order( candy$winpercent )
head( candy[inds,], 5 )
```

	chocolate	fruity	caramel	peanut	almond	nougat
Nik L Nip	0	1	0		0	0
Boston Baked Beans	0	0	0		1	0
Chiclets	0	1	0		0	0
Super Bubble	0	1	0		0	0
Jawbusters	0	1	0		0	0

	crisped	rice	wafer	hard	bar	pluribus	sugar	percent	price	percent
Nik L Nip				0	0	0	1	0.197		0.976
Boston Baked Beans				0	0	0	1	0.313		0.511
Chiclets				0	0	0	1	0.046		0.325
Super Bubble				0	0	0	0	0.162		0.116
Jawbusters				0	1	0	1	0.093		0.511

	winpercent
Nik L Nip	22.44534
Boston Baked Beans	23.41782
Chiclets	24.52499
Super Bubble	27.30386
Jawbusters	28.12744

Q14. What are the top 5 all time favorite candy types out of this set?

```
inds <- order( candy$winpercent, decreasing = T )
head( candy[inds,], 5 )
```

	chocolate	fruity	caramel	peanut	almond	nougat
Reese's Peanut Butter cup	1	0	0		1	0
Reese's Miniatures	1	0	0		1	0
Twix	1	0	1		0	0
Kit Kat	1	0	0		0	0
Snickers	1	0	1		1	1

	crisped	rice	wafer	hard	bar	pluribus	sugar	percent
Reese's Peanut Butter cup				0	0	0	0	0.720
Reese's Miniatures				0	0	0	0	0.034
Twix				1	0	1	0	0.546
Kit Kat				1	0	1	0	0.313
Snickers				0	0	1	0	0.546


  

	price	percent	winpercent
Reese's Peanut Butter cup	0.651		84.18029

Reese's Miniatures	0.279	81.86626
Twix	0.906	81.64291
Kit Kat	0.511	76.76860
Snickers	0.651	76.67378


Make a bar plot with ggplot and order it by winpercent values.

```
ggplot(candy) +
  aes(winpercent, rownames(candy)) +
  geom_col()
```




```
ggplot(candy) +
  aes(winpercent, reorder( rownames(candy), winpercent )) +
  geom_col()
```






```
ggplot(candy) +
  aes(winpercent, reorder( rownames(candy), winpercent )) +
  geom_col(fill="red")
```




```
ggplot(candy) +
  aes(x=winpercent, y=reorder( rownames(candy), winpercent),
      fill=chocolate) +
  geom_col()
```



Here we want a custom color vector to color each bar the way we want - with `chocolate` and `fruity` candy together with whether it is a bar or not

```
mycols <- rep("gray", nrow(candy))
mycols[as.logical(candy$chocolate)] <- "chocolate"
mycols[as.logical(candy$fruity)] <- "pink"
mycols[as.logical(candy$bar)] <- "brown"

#mycols
ggplot(candy) +
  aes(winpercent, reorder( rownames(candy), winpercent )) +
  geom_col(fill=mycols)
```



```
ggsave("mybarplot.png", width=3, height=6)
```





Figure 1: My silly barplot image

#### 4. Winpercent vs Pricepercent

```
# Pink and gray is too light, let's change to red
library(ggrepel)
mycols <- rep("black", nrow(candy))
mycols[as.logical(candy$chocolate)] <- "chocolate"
mycols[as.logical(candy$fruity)] <- "red"
mycols[as.logical(candy$bar)] <- "brown"

# How about a plot of price vs win
ggplot(candy) +
  aes(winpercent, pricepercent, label=rownames(candy)) +
  geom_point(col=mycols) +
  geom_text_repel(col=mycols, size=3.3, max.overlaps = 5)
```

Warning: ggrepel: 65 unlabeled data points (too many overlaps). Consider increasing max.overlaps



## 5. Correlation Structure

```
cij <- cor(candy)
cij
```


	chocolate	fruity	caramel	peanutyalmondy	nougat
chocolate	1.0000000	-0.74172106	0.24987535	0.37782357	0.25489183
fruity	-0.7417211	1.00000000	-0.33548538	-0.39928014	-0.26936712
caramel	0.2498753	-0.33548538	1.00000000	0.05935614	0.32849280
peanutyalmondy	0.3778236	-0.39928014	0.05935614	1.00000000	0.21311310
nougat	0.2548918	-0.26936712	0.32849280	0.21311310	1.00000000
crispedricewafer	0.3412098	-0.26936712	0.21311310	-0.01764631	-0.08974359
hard	-0.3441769	0.39067750	-0.12235513	-0.20555661	-0.13867505
bar	0.5974211	-0.51506558	0.33396002	0.26041960	0.52297636
pluribus	-0.3396752	0.29972522	-0.26958501	-0.20610932	-0.31033884
sugarpercent	0.1041691	-0.03439296	0.22193335	0.08788927	0.12308135
pricepercent	0.5046754	-0.43096853	0.25432709	0.30915323	0.15319643
winpercent	0.6365167	-0.38093814	0.21341630	0.40619220	0.19937530
	crispedricewafer	hard	bar	pluribus	
chocolate	0.34120978	-0.34417691	0.59742114	-0.33967519	
fruity	-0.26936712	0.39067750	-0.51506558	0.29972522	
caramel	0.21311310	-0.12235513	0.33396002	-0.26958501	
peanutyalmondy	-0.01764631	-0.20555661	0.26041960	-0.20610932	
nougat	-0.08974359	-0.13867505	0.52297636	-0.31033884	
crispedricewafer	1.00000000	-0.13867505	0.42375093	-0.22469338	
hard	-0.13867505	1.00000000	-0.26516504	0.01453172	
bar	0.42375093	-0.26516504	1.00000000	-0.59340892	
pluribus	-0.22469338	0.01453172	-0.59340892	1.00000000	
sugarpercent	0.06994969	0.09180975	0.09998516	0.04552282	
pricepercent	0.32826539	-0.24436534	0.51840654	-0.22079363	
winpercent	0.32467965	-0.31038158	0.42992933	-0.24744787	
	sugarpercent	pricepercent	winpercent		
chocolate	0.10416906	0.5046754	0.6365167		
fruity	-0.03439296	-0.4309685	-0.3809381		
caramel	0.22193335	0.2543271	0.2134163		
peanutyalmondy	0.08788927	0.3091532	0.4061922		
nougat	0.12308135	0.1531964	0.1993753		
crispedricewafer	0.06994969	0.3282654	0.3246797		
hard	0.09180975	-0.2443653	-0.3103816		
bar	0.09998516	0.5184065	0.4299293		
pluribus	0.04552282	-0.2207936	-0.2474479		

sugarpercent	1.00000000	0.3297064	0.2291507
pricepercent	0.32970639	1.0000000	0.3453254
winpercent	0.22915066	0.3453254	1.0000000

```
library(corrplot)
```

```
corrplot 0.95 loaded
```

```
corrplot(cij)
```



Q22. Examining this plot what two variables are anti-correlated (i.e. have minus values)?

Chocolate and fruity are negatively correlated.

```
cij["chocolate", "fruity"]
```

```
[1] -0.7417211
```

Q23. Similarly, what two variables are most positively correlated?

Winpercent and chocolate are most positively correlated.

```
cij["winpercent", "chocolate"]
```

```
[1] 0.6365167
```

## Principal Component Analysis (PCA)

We need to be sure to scale our input `candy` data before PCA as we have the `winpercent` column on a different scale to all others in the dataset.

```
pca <- prcomp(candy, scale=T)
summary(pca)
```

Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	2.0788	1.1378	1.1092	1.07533	0.9518	0.81923	0.81530
Proportion of Variance	0.3601	0.1079	0.1025	0.09636	0.0755	0.05593	0.05539
Cumulative Proportion	0.3601	0.4680	0.5705	0.66688	0.7424	0.79830	0.85369


	PC8	PC9	PC10	PC11	PC12
Standard deviation	0.74530	0.67824	0.62349	0.43974	0.39760
Proportion of Variance	0.04629	0.03833	0.03239	0.01611	0.01317
Cumulative Proportion	0.89998	0.93832	0.97071	0.98683	1.00000

First main result figure is my “PCA plot”

```
#pca$x
ggplot(pca$x) +
  aes(PC1, PC2, label=rownames(pca$x)) +
  geom_point(col=mycols) +
  geom_text_repel(max.overlaps = 6, col=mycols) +
  theme_bw()
```


Warning: ggrepel: 67 unlabeled data points (too many overlaps). Consider increasing max.overlaps






The second main PCA result is in the `pca$rotation` we can plot this to generate a so-called “loadings” plot.

```
#pca$rotation
ggplot(pca$rotation) +
  aes(PC1, rownames(pca$rotation)) +
  geom_col()
```



```
ggplot(pca$rotation) +
  aes(PC1, reorder(rownames(pca$rotation), PC1), fill=PC1) +
  geom_col()
```



Q24. What original variables are picked up strongly by PC1 in the positive direction? Do these make sense to you?

Fruity, pluribus, and bar are originally picked up strongly by PC1 in the positive direction. These make sense to me.