

On Tseitin formulas, read-once branching programs and treewidth

Ludmila Glinskih, Dmitry Itsykson

St. Petersburg Department of V. A. Steklov Institute of Mathematics


24th Estonian Winter School in Computer Science
Palmse, Estonia
March 7, 2019

Outline

- ▶ Branching programs
- ▶ Tseitin formulas
- ▶ Treewidth and pathwidth
- ▶ Lower bounds

Branching program

- ▶ BP is a way to represent Boolean function
 - ▶ directed graph without cycles;
 - ▶ two sinks: labeled with 0 and 1, one source;
 - ▶ every internal vertex labeled with a variable and has two outgoing edges: labeled with 0 and 1;
 - ▶ the value of function equals label of the sink in the end of corresponding path.




Branching program

- ▶ BP is a way to represent Boolean function

- ▶ directed graph without cycles;
- ▶ two sinks: labeled with 0 and 1, one source;
- ▶ every internal vertex labeled with a variable and has two outgoing edges: labeled with 0 and 1;
- ▶ the value of function equals label of the sink in the end of corresponding path.


- ▶ BP size is between circuit and formula sizes

- ▶ $C(f) \leq 3BP(f) \leq O(L^{1+\epsilon}(f))$,
- ▶ $C(f)$ circuit complexity of f , $L(f)$ size of minimal formula, BP size of minimal branching program.




Non-deterministic Branching program

- ▶ NBP additionally has non-deterministic nodes:
 - ▶ non-deterministic nodes are unlabeled
 - ▶ the value equals 1 iff there exists a path to 1-sink




Non-deterministic Branching program

- ▶ NBP additionally has non-deterministic nodes:
 - ▶ non-deterministic nodes are unlabeled
 - ▶ the value equals 1 iff there exists a path to 1-sink
- ▶ BP corresponds to $L/poly$
- ▶ NBP corresponds to $NL/poly$




Non-deterministic Branching program

- ▶ NBP additionally has non-deterministic nodes:
 - ▶ non-deterministic nodes are unlabeled
 - ▶ the value equals 1 iff there exists a path to 1-sink
- ▶ BP corresponds to $L/poly$
- ▶ NBP corresponds to $NL/poly$
- ▶ k -(N)BP if for every path every variable occurs no more than k times



Non-deterministic Branching program


- ▶ NBP additionally has non-deterministic nodes:
 - ▶ non-deterministic nodes are unlabeled
 - ▶ the value equals 1 iff there exists a path to 1-sink
- ▶ BP corresponds to L/poly
- ▶ NBP corresponds to NL/poly
- ▶ $k\text{-(N)BP}$ if for every path every variable occurs no more than k times
- ▶ OBDD is 1-BP with fixed order for variables in every path



Tseitin formulas

- ▶ Tseitin $TS_{G,c}$ formula defined for a graph
 - ▶ every edge is labeled with a variable
 - ▶ every vertex has a 0-1 label: $c : V \rightarrow 0, 1$
 - ▶ $TS_{G,c}(x) = 1 \iff$


$$\bigwedge_{v \in V} \left(\sum_{e \text{ incident } v} x_e = c(v) \bmod 2 \right)$$



$$\begin{cases} x_1 + x_2 = 1 \\ x_1 + x_3 + x_4 = 0 \\ x_2 + x_4 = 1 \\ x_3 = 1 \end{cases}$$

Tseitin formulas


- ▶ Tseitin $TS_{G,c}$ formula defined for a graph
 - ▶ every edge is labeled with a variable
 - ▶ every vertex has a 0-1 label: $c : V \rightarrow 0, 1$
 - ▶ $TS_{G,c}(x) = 1 \iff \bigwedge_{v \in V} \left(\sum_{e \text{ incident } v} x_e = c(v) \bmod 2 \right)$
- ▶ A Tseitin formula is satisfiable iff for every connected component the sum of labels is even



$$\begin{cases} x_1 + x_2 = 1 \\ x_1 + x_3 + x_4 = 0 \\ x_2 + x_4 = 1 \\ x_3 = 1 \end{cases}$$

Tseitin formulas

- ▶ Tseitin $TS_{G,c}$ formula defined for a graph
 - ▶ every edge is labeled with a variable
 - ▶ every vertex has a 0-1 label: $c : V \rightarrow 0, 1$
 - ▶ $TS_{G,c}(x) = 1 \iff \bigwedge_{v \in V} \left(\sum_{e \text{ incident } v} x_e = c(v) \bmod 2 \right)$
- ▶ A Tseitin formula is satisfiable iff for every connected component the sum of labels is even
- ▶ Unsatisfiable Tseitin formulas are classical hard examples for proof systems:
 - ▶ Resolution [Tseitin, 1968], [Urquhart, 1987]
 - ▶ Bounded depth Frege [Ben-Sasson, 2002], [Pitassi, Rossman, Servedio, Tan, 2016]
 - ▶ Polynomial Calculus over field with char $\neq 2$ [Alekhnovich, Razborov, 2001]
 - ▶ Tree-like Lovasz-Schrijver [Itsykson, Kojevnikov, 2006]



$$\begin{cases} x_1 + x_2 = 1 \\ x_1 + x_3 + x_4 = 0 \\ x_2 + x_4 = 1 \\ x_3 = 1 \end{cases}$$

Tseitin formulas (2)

- ▶ (Itsykson, Knop, Romashchenko, Sokolov, 2017) Exponential lower bound for OBDD(join) proof systems for unsatisfiable Tseitin formulas
 - ▶ key step: satisfiable Tseitin formulas are hard for OBDDs.
 - ▶ $OBDD(TS_{G,c}) = 2^{\Omega(n)}$

Tseitin formulas (2)

- ▶ (Itsykson, Knop, Romashchenko, Sokolov, 2017) Exponential lower bound for OBDD(join) proof systems for unsatisfiable Tseitin formulas
 - ▶ key step: satisfiable Tseitin formulas are hard for OBDDs.
 - ▶ $OBDD(TS_{G,c}) = 2^{\Omega(n)}$
- ▶ OBDD is a partial case of 1-BP
 - ▶ (Bryant, 1991) (Sieling, Wegener, 1995)
 $f : OBDD(f) = 2^{\Omega(n)}, 1\text{-BP}(f) = \text{poly}(n)$

The goal: what is the size of 1-BP for satisfiable Tseitin formulas?

Lower bound for Tseitin formula on complete graph

Theorem:

1-NBP for satisfiable Tseitin formula on complete graph on n vertices has size $2^{\Omega(n)}$.

Lower bound for Tseitin formula on complete graph

Theorem:



1-NBP for satisfiable Tseitin formula on complete graph on n vertices has size $2^{\Omega(n)}$.

Generalized theorem: 1-NBP for a satisfiable Tseitin formula on a connected graph G has at least $2^{|V| - k_G(l) - k_G(|E| - l) + 1}$ nodes, where the value $k_G(l)$ denotes the maximal number of connected components that can be obtained from G by deletion of l edges.

Treewidth and pathwidth

Tree/path decomposition of a graph

$G = (V, E)$ is a graph T that is a tree/path





Treewidth and pathwidth

Tree/path decomposition of a graph

$G = (V, E)$ is a graph T that is a tree/path

1. Vertices of T (called "bags") correspond to sets of vertices of G





Treewidth and pathwidth

Tree/path decomposition of a graph

$G = (V, E)$ is a graph T that is a tree/path

1. Vertices of T (called "bags") correspond to sets of vertices of G
2. If $(a, b) \in E$ then there is a bag with a and b





Treewidth and pathwidth

Tree/path decomposition of a graph

$G = (V, E)$ is a graph T that is a tree/path

1. Vertices of T (called "bags") correspond to sets of vertices of G
2. If $(a, b) \in E$ then there is a bag with a and b
3. All bags with the same vertex form a tree/path





Treewidth and pathwidth

Tree/path decomposition of a graph

$G = (V, E)$ is a graph T that is a tree/path

1. Vertices of T (called "bags") correspond to sets of vertices of G
2. If $(a, b) \in E$ then there is a bag with a and b
3. All bags with the same vertex form a tree/path

Width of a decomposition is the size of a maximal bag in it minus one.



Treewidth and pathwidth



Tree/path decomposition of a graph

$G = (V, E)$ is a graph T that is a tree/path

1. Vertices of T (called "bags") correspond to sets of vertices of G
2. If $(a, b) \in E$ then there is a bag with a and b
3. All bags with the same vertex form a tree/path

Width of a decomposition is the size of a maximal bag in it minus one.

Treewidth/pathwidth of a graph is the minimal width among all its tree/path decompositions.



Lower and upper bounds

Lower bound via treewidth: 1-NBP for satisfiable Tseitin formula on a graph G has size $\Omega(2^{t^\delta})$, where t is a treewidth of G , $\delta < \frac{1}{36}$.

Lower and upper bounds

Lower bound via treewidth: 1-NBP for satisfiable Tseitin formula on a graph G has size $\Omega(2^{t^\delta})$, where t is a treewidth of G , $\delta < \frac{1}{36}$.

Upper bound via pathwidth: a satisfiable Tseitin formula on graph G can be computed by an OBDD of size $O(m2^{p+1})$, where m is the number of edges and p is the pathwidth of G .

Graph minor

Minor of a graph G is a graph that can be obtained from G by sequence of edge contractions, edge and vertex deletion.

Graph minor

Minor of a graph G is a graph that can be obtained from G by sequence of edge contractions, edge and vertex deletion.

Grid Minor Theorem [Robertson, Seymour 1986], [Chuzhoy 2015]:

Every graph G of a tree-width t has a grid minor of size $t^\delta \times t^\delta$, where $\delta < \frac{1}{36}$.

Lower bound: sketch of the proof

Lower bound via tree-width: 1-NBP for satisfiable Tseitin formula on a graph G has size $\Omega(2^{t^\delta})$, where t is a treewidth of G , $\delta < \frac{1}{36}$.

Lower bound: sketch of the proof

Lower bound via tree-width: 1-NBP for satisfiable Tseitin formula on a graph G has size $\Omega(2^{t^\delta})$, where t is a treewidth of G , $\delta < \frac{1}{36}$.

Sketch of the proof:

Lower bound: sketch of the proof

Lower bound via tree-width: 1-NBP for satisfiable Tseitin formula on a graph G has size $\Omega(2^{t^\delta})$, where t is a treewidth of G , $\delta < \frac{1}{36}$.

Sketch of the proof:

- ▶ Take the smallest 1-NBP for a Tseitin formula on a graph G

Lower bound: sketch of the proof

Lower bound via tree-width: 1-NBP for satisfiable Tseitin formula on a graph G has size $\Omega(2^{t^\delta})$, where t is a treewidth of G , $\delta < \frac{1}{36}$.

Sketch of the proof:

- ▶ Take the smallest 1-NBP for a Tseitin formula on a graph G
- ▶ Apply sequence of edge contractions, edge and vertex deletions that could lead only to a smaller 1-NBP

Lower bound: sketch of the proof

Lower bound via tree-width: 1-NBP for satisfiable Tseitin formula on a graph G has size $\Omega(2^{t^\delta})$, where t is a treewidth of G , $\delta < \frac{1}{36}$.

Sketch of the proof:

- ▶ Take the smallest 1-NBP for a Tseitin formula on a graph G
- ▶ Apply sequence of edge contractions, edge and vertex deletions that could lead only to a smaller 1-NBP
- ▶ Obtain 1-NBP for a $t^\delta \times t^\delta$ grid-minor of a graph G

Lower bound: sketch of the proof

Lower bound via tree-width: 1-NBP for satisfiable Tseitin formula on a graph G has size $\Omega(2^{t^\delta})$, where t is a treewidth of G , $\delta < \frac{1}{36}$.

Sketch of the proof:

- ▶ Take the smallest 1-NBP for a Tseitin formula on a graph G
- ▶ Apply sequence of edge contractions, edge and vertex deletions that could lead only to a smaller 1-NBP
- ▶ Obtain 1-NBP for a $t^\delta \times t^\delta$ grid-minor of a graph G
- ▶ Show that 1-NBP for $t^\delta \times t^\delta$ grid graph has size $\Omega(2^{t^\delta})$

Open questions

Open questions

- ▶ Currently we have a lower bound using tree-width and upper bound using path-width. Can we close this gap?
 - ▶ $tw(G) \leq pw(G) \leq tw(G) \cdot \log(tw(G))$

Open questions

- ▶ Currently we have a lower bound using tree-width and upper bound using path-width. Can we close this gap?
 - ▶ $tw(G) \leq pw(G) \leq tw(G) \cdot \log(tw(G))$
- ▶ Prove non-trivial lower bound for semantic 1-NBP
 - ▶ only consistent paths should contain every variable no more than once
 - ▶ that means that all paths that consist repeated variable should not be reachable by any assignment

Open questions

- ▶ Currently we have a lower bound using tree-width and upper bound using path-width. Can we close this gap?
 - ▶ $tw(G) \leq pw(G) \leq tw(G) \cdot \log(tw(G))$
- ▶ Prove non-trivial lower bound for semantic 1-NBP
 - ▶ only consistent paths should contain every variable no more than once
 - ▶ that means that all paths that consist repeated variable should not be reachable by any assignment



No exponential lower bounds for semantic 1-NBP for a boolean function

All known results with exponential lower bounds are for functions with non-boolean domain [Cook, Edmonds, Medabalimi, Pitassi, 2016], [Jukna, 2009]

Lower bound for a grid graph

We want to show that 1-NBP for $s \times s$ grid graph has size $2^{\Omega(s)}$



- Use theorem: 1-NBP for a satisfiable Tseitin formula on a graph G has at least $2^{|V| - k_G(I) - k_G(|E| - I) + 1}$ nodes



Lower bound for a grid graph

We want to show that 1-NBP for $s \times s$ grid graph has size $2^{\Omega(s)}$



- Use theorem: 1-NBP for a satisfiable Tseitin formula on a graph G has at least $2^{|V| - k_G(I) - k_G(|E| - I) + 1}$ nodes



Lower bound for a grid graph

We want to show that 1-NBP for $s \times s$ grid graph has size $2^{\Omega(s)}$



- ▶ Use theorem: 1-NBP for a satisfiable Tseitin formula on a graph G has at least $2^{|V| - k_G(I) - k_G(|E| - I) + 1}$ nodes
- ▶ We set $I = \frac{|E|}{2}$



Lower bound for a grid graph

We want to show that 1-NBP for $s \times s$ grid graph has size $2^{\Omega(s)}$

- ▶ Use theorem: 1-NBP for a satisfiable Tseitin formula on a graph G has at least $2^{|V| - k_G(I) - k_G(|E| - I) + 1}$ nodes
- ▶ We set $I = \frac{|E|}{2}$
- ▶ We want to show that the number of connected components that can be obtained after deletion of half of edges is at most $\frac{|V|}{2} - \varepsilon \sqrt{|V|}$
- ▶ After that we get a lower bound $2^{\Omega(2\varepsilon \sqrt{|V|})} = 2^{\Omega(s)}$



Reducing NBP to its minor

We need to check that all operations can only reduce the size of a 1-NBP

Reducing NBP to its minor

We need to check that all operations can only reduce the size of a 1-NBP

- ▶ **Deletion of edge** x_i : substitute any value that will not break the satisfiability of the formula

Reducing NBP to its minor

We need to check that all operations can only reduce the size of a 1-NBP

- ▶ **Deletion of edge x_i :** substitute any value that will not break the satisfiability of the formula
- ▶ **Deletion of vertex v :** delete all connected edges one-by-one

Reducing NBP to its minor

We need to check that all operations can only reduce the size of a 1-NBP

- ▶ **Deletion of edge x_i** : substitute any value that will not break the satisfiability of the formula
- ▶ **Deletion of vertex v** : delete all connected edges one-by-one
- ▶ **Contraction of edge x_i** : the same as the satisfiability of the formula $\exists x_i : Ts_{G,w}(x)$
 - ▶ in 1-NBP all nodes labeled with x_i should be changed to non-deterministic nodes

Upper bound using path-width

Theorem: a satisfiable Tseitin formula on graph G can be computed by an OBDD of size $O(m2^{p+1})$, where m is the number of edges and p is the path-width of G .

Upper bound using path-width

Theorem: a satisfiable Tseitin formula on graph G can be computed by an OBDD of size $O(m2^{p+1})$, where m is the number of edges and p is the path-width of G .

Plan of the proof: We build OBDD layer-by-layer, each layer corresponds to one edge. Each edge corresponds to the first bag where it appears in the decomposition.

Upper bound using path-width

Theorem: a satisfiable Tseitin formula on graph G can be computed by an OBDD of size $O(m2^{p+1})$, where m is the number of edges and p is the path-width of G .

Plan of the proof: We build OBDD layer-by-layer, each layer corresponds to one edge. Each edge corresponds to the first bag where it appears in the decomposition. For every edge and its bag we add at most 2^{p+1} vertices that correspond to all possible values of parity in vertices from that bag.

Known lower bounds for k-BPs

- ▶ (Borodin, Razborov, Smolensky, 1993)

$$CLIQUE_ONLY_n(G) : \{0, 1\}^{\frac{n \cdot (n-1)}{2}} \rightarrow \{0, 1\}$$

- ▶ $CLIQUE_ONLY_n(G) = 1$ iff graph G on n vertices is exactly $\frac{n}{2}$ -clique
- ▶ $1\text{-NBP}(CLIQUE_ONLY_n) = 2^{\Omega(n)}$
- ▶ $2\text{-BP}(CLIQUE_ONLY_n) = \text{poly}(n)$

Known lower bounds for k-BPs

- ▶ (Borodin, Razborov, Smolensky, 1993)
 $CLIQUE_ONLY_n(G) : \{0, 1\}^{\frac{n \cdot (n-1)}{2}} \rightarrow 0, 1$
 - ▶ $CLIQUE_ONLY_n(G) = 1$ iff graph G on n vertices is exactly $\frac{n}{2}$ -clique
 - ▶ $1\text{-NBP}(CLIQUE_ONLY_n) = 2^{\Omega(n)}$
 - ▶ $2\text{-BP}(CLIQUE_ONLY_n) = poly(n)$
- ▶ (Borodin, Razborov, Smolensky, 1993) first strongly exponential lower bound
 - ▶ f artificially constructed function
 - ▶ $1\text{-NBP}(f) = 2^{\Omega(n)}$

Known lower bounds for k-BPs

- ▶ (Borodin, Razborov, Smolensky, 1993)
 $CLIQUE_ONLY_n(G) : \{0, 1\}^{\frac{n \cdot (n-1)}{2}} \rightarrow 0, 1$
 - ▶ $CLIQUE_ONLY_n(G) = 1$ iff graph G on n vertices is exactly $\frac{n}{2}$ -clique
 - ▶ $1\text{-NBP}(CLIQUE_ONLY_n) = 2^{\Omega(n)}$
 - ▶ $2\text{-BP}(CLIQUE_ONLY_n) = \text{poly}(n)$
- ▶ (Borodin, Razborov, Smolensky, 1993) first strongly exponential lower bound
 - ▶ f artificially constructed function
 - ▶ $1\text{-NBP}(f) = 2^{\Omega(n)}$
- ▶ (Thathachar, 1998) an explicit functions f_k for every k :
 - ▶ $k\text{-NBP}(f_k) = 2^{\Omega(n^{1/k})}$
 - ▶ $(k+1)\text{-BP}(f_k) = O(n)$

Known lower bounds for k-BPs (2)

- ▶ (Duris, Hromkovic, Jukna, Sauerhoff, Schnitger, 2004)
 - ▶ \oplus parity of the number of triangles in a graph, $\overline{\Delta}$ is 1 iff graph has no triangles
 - ▶ $1\text{-NBP}(\oplus) = 2^{\Omega(n)}$
 - ▶ $1\text{-NBP}(\overline{\Delta}) = 2^{\Omega(n)}$

Known lower bounds for k-BPs (2)

- ▶ (Duris, Hromkovic, Jukna, Sauerhoff, Schnitger, 2004)
 - ▶ \oplus parity of the number of triangles in a graph, $\overline{\Delta}$ is 1 iff graph has no triangles
 - ▶ $1\text{-NBP}(\oplus) = 2^{\Omega(n)}$
 - ▶ $1\text{-NBP}(\overline{\Delta}) = 2^{\Omega(n)}$
- ▶ (Jukna, 1995)
 - ▶ f_k characteristic function of error-correcting codes
 - ▶ $k\text{-NBP}(f_k) = 2^{\Omega(\sqrt{n})}$

The comparison with other results

Satisfiable Tseitin formulas $TS_{G,w}$ on (n, d, α) -expander can be represented by:

- ▶ $\text{CNF}(TS_{G,c}) = O(n)$
- ▶ $1\text{-NBP}(\lceil TS_{G,c} \rceil) = O(n)$
- ▶ $2\text{-BP}(TS_{G,c}) = O(n)$
- ▶ $1\text{-NBP}(TS_{G,c}) = 2^{\Omega(n)}$

The comparison with other results

Satisfiable Tseitin formulas $TS_{G,w}$ on (n, d, α) -expander can be represented by:

- ▶ $\text{CNF}(TS_{G,c}) = O(n)$
- ▶ $1\text{-NBP}(\lceil TS_{G,c} \rceil) = O(n)$
- ▶ $2\text{-BP}(TS_{G,c}) = O(n)$
- ▶ $1\text{-NBP}(TS_{G,c}) = 2^{\Omega(n)}$

Best previously known gaps:

$\text{CNF}(f) = O(n^3)$	$1\text{-NBP}(f) = 2^{\Omega(n)}$ (Duris et al, 2004)
$1\text{-NBP}(\lceil f \rceil) = O(n)$	$1\text{-NBP}(f) = 2^{\Omega(\sqrt{n})}$ (Jukna, 2009) explicit construction
$1\text{-NBP}(\lceil f \rceil) = O(n)$	$1\text{-NBP}(f) = 2^{\Omega(n)}$ (Duris et al, 2004) probabilistic construction
$2\text{-BP}(f) = O(n)$	$1\text{-NBP}(f) = 2^{\Omega(\sqrt{n})}$ (Thathachar, 1998)

Idea of the proof

Theorem: 1-NBP for a satisfiable Tseitin formula on a connected graph G has at least $2^{|V| - k_G(l) - k_G(|E| - l) + 1}$ nodes, where the value $k_G(l)$ denotes the maximal number of connected components that can be obtained from G by deletion of l edges.

Idea of the proof

Theorem: 1-NBP for a satisfiable Tseitin formula on a connected graph G has at least $2^{|V| - k_G(l) - k_G(|E| - l) + 1}$ nodes, where the value $k_G(l)$ denotes the maximal number of connected components that can be obtained from G by deletion of l edges.

Idea of the proof:

Choose level l of the 1-NBP

Idea of the proof

Theorem: 1-NBP for a satisfiable Tseitin formula on a connected graph G has at least $2^{|V| - k_G(l) - k_G(|E| - l) + 1}$ nodes, where the value $k_G(l)$ denotes the maximal number of connected components that can be obtained from G by deletion of l edges.

Idea of the proof:

Choose level l of the 1-NBP

- ▶ show that there are at least $2^{C_1 n}$ different non-zero paths to level l

Idea of the proof

Theorem: 1-NBP for a satisfiable Tseitin formula on a connected graph G has at least $2^{|V|-k_G(l)-k_G(|E|-l)+1}$ nodes, where the value $k_G(l)$ denotes the maximal number of connected components that can be obtained from G by deletion of l edges.

Idea of the proof:

Choose level l of the 1-NBP

- ▶ show that there are at least $2^{C_1 n}$ different non-zero paths to level l
- ▶ show that for every node v on the l -th level only $2^{C_2 n}$ non-zero paths go from source to the same node v

Idea of the proof

Theorem: 1-NBP for a satisfiable Tseitin formula on a connected graph G has at least $2^{|V|-k_G(I)-k_G(|E|-I)+1}$ nodes, where the value $k_G(I)$ denotes the maximal number of connected components that can be obtained from G by deletion of I edges.

Idea of the proof:

Choose level I of the 1-NBP

- ▶ show that there are at least $2^{C_1 n}$ different non-zero paths to level I
- ▶ show that for every node v on the I -th level only $2^{C_2 n}$ non-zero paths go from source to the same node v

Get that at least $2^{(C_1-C_2)n}$ different nodes are on the I -th level, so is in the 1-NBP