

Inference for Linear Regression

November 6, 2019

Regression Example

Asking R for a summary of the regression model, we get the following:

```
lm(formula = eruptions ~ waiting)
```

Residuals:

Min	1Q	Median	3Q	Max
-1.29917	-0.37689	0.03508	0.34909	1.19329

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-1.874016	0.160143	-11.70	<2e-16 ***
waiting	0.075628	0.002219	34.09	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.4965 on 270 degrees of freedom
Multiple R-squared: 0.8115, Adjusted R-squared: 0.8108
F-statistic: 1162 on 1 and 270 DF, p-value: < 2.2e-16

Let's pick this apart piece by piece.

Regression Example

```
Call:
lm(formula = eruptions ~ waiting)

Residuals:
    Min       1Q   Median       3Q      Max
-1.29917 -0.37689  0.03508  0.34909  1.19329
```

- The first line shows the command used in **R** to run this regression model.
- The **Residuals** item shows a quartile-based summary of our residuals.

Regression Example

```
Residual standard error: 0.4965 on 270 degrees of freedom  
Multiple R-squared: 0.8115,    Adjusted R-squared: 0.8108  
F-statistic: 1162 on 1 and 270 DF,  p-value: < 2.2e-16
```

The **F-statistic** and **p-value** give information about the model overall.

- These are based on an F-distribution.
- The null hypothesis is that all of our model parameters are 0 (the model gives us no good info).
- Since $p\text{-value} < 2.2 \times 10^{-16} < \alpha = 0.05$, at least one of the parameters is nonzero (the model is useful).

Regression Example

```
Residual standard error: 0.4965 on 270 degrees of freedom  
Multiple R-squared: 0.8115,    Adjusted R-squared: 0.8108  
F-statistic: 1162 on 1 and 270 DF,  p-value: < 2.2e-16
```

- Multiple R-squared is our squared correlation coefficient R^2 .
- This tells us how good our fit is.
- Ignore the adjusted R-squared and residual standard error for now.

Regression Example

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-1.874016	0.160143	-11.70	<2e-16
waiting	0.075628	0.002219	34.09	<2e-16

Finally, the `Coefficients` section gives us several pieces of information:

- 1 `Estimate` shows the estimated parameters for each value.
- 2 `Std. Error` gives the standard error for each parameter estimate.
- 3 The `t values` are the test statistics for each parameter estimate.
- 4 Finally, `Pr(>|t|)` are the p-values for each parameter estimate.

Regression Example

The hypothesis test for each regression coefficient has hypotheses

$$H_0 : \beta_i = 0$$

$$H_A : \beta_i \neq 0$$

where $i = 0$ for the intercept and $i = 1$ for the slope.

Regression Example

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-1.874016	0.160143	-11.70	<2e-16
waiting	0.075628	0.002219	34.09	<2e-16

- $p\text{-value} < 2 \times 10^{-16}$ for b_0 so we can conclude that the intercept is nonzero.
- $p\text{-value} < 2 \times 10^{-16}$ for b_1 so we conclude that the intercept is also nonzero.
- This means that the intercept and slope both provide useful information when predicting values of $y = \text{eruptions}$.

Confidence Intervals for a Coefficient

We can construct confidence intervals similar to those for hypothesis tests. A $(1 - \alpha)100\%$ confidence interval for β_i is

$$b_i \pm t_{\alpha/2}(df) \times SE(b_i)$$

where the model df and SE can be found in the regression output.

Aside: ANOVA for Regression Models

- ANOVA will also play a role in regression.
- We can get the ANOVA table for a regression.

Aside: ANOVA for Regression Models

The ANOVA table in regression will look something like this:

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
faithful\$waiting	1	286.478	286.478	1162.1	< 2.2e-16
Residuals	270	66.562	0.247		

Example

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-1.874016	0.160143	-11.70	<2e-16
waiting	0.075628	0.002219	34.09	<2e-16

Find 95% confidence intervals for β_0 and β_1 .

Estimation and Prediction Using a Regression Line

We now know

- how to examine if a model is useful.
- how to confirm that our regression assumptions are satisfied.

Estimation and Prediction Using a Regression Line

Given a useful regression line, we want to

- estimate an average value of y for a given value of x .
- estimate a particular value of y for a given value of x .

Estimation and Prediction Using a Regression Line

We've already talked about using a regression line to make predictions.

$$\hat{y} = b_0 + b_1x$$

Plug in x and we get a good estimate for the *average* value of y at that point.

Estimation and Prediction Using a Regression Line

Point estimates are useful, but we want to consider variability!

- Recall: one of our regression assumptions is normally distributed errors.
- This means that the variability around the regression line should be approximately normal
 - with mean $\beta_0 + \beta_1 x$
 - and standard deviation σ^2 .

The Variability of \hat{y}

- Notice that \hat{y} is an estimator.
- The variability of an estimator is its standard error.
- Then σ^2 is well-approximated by

$$SE(\hat{y}) = \sqrt{\text{MSE} \left(\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{s_x} \right)}$$

The Variability of \hat{y}

Since we are working with a normal distribution, estimation and testing can be based on the test statistic

$$t = \frac{\hat{y} - y_0}{SE(\hat{y})}$$

which corresponds to a $t(n - 2)$ distribution.

Confidence Intervals for \bar{y}

A $(1 - \alpha)100\%$ confidence interval for the average value of y (measured by $\beta_0 + \beta_1 x$) when $x = x_0$ is

$$\hat{y} \pm t_{\alpha/2}(n - 2) \times SE(\hat{y})$$

or

$$\hat{y} \pm t_{\alpha/2}(n - 2) \times \sqrt{\text{MSE} \left(\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{s_x} \right)}$$

Prediction Intervals for \bar{y}

- So far, we've only considered *average* values of the outcome variable y .
- What if we wanted to predict a *particular* value of y ?

Prediction Intervals for \bar{y}

For a residual,

$$e = \epsilon + \text{error in estimating line}$$

- We don't know the true breakdown between these components.
- ...but we can use this concept to build a new standard error formula.

Prediction Intervals for \bar{y}

The standard error of $(y - \hat{y})$ is

$$SE(y - \hat{y}) = \sqrt{\text{MSE} \left(1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{s_x} \right)}$$

Prediction Intervals for \bar{y}

A $(1 - \alpha)100\%$ **prediction interval** for a specific value of y when $x = x_0$ is

$$\hat{y} \pm t_{\alpha/2}(n - 2) \times SE(y - \hat{y})$$

or

$$\hat{y} \pm t_{\alpha/2}(n - 2) \times \sqrt{\text{MSE} \left(1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{s_x} \right)}$$