(a) places: a frame, and the split frame of H_5^* (Prop. HL5)



cell and offset of c at one place a_c; layer $\lambda_c\mu_d/(a_c - a_d)$: Plücker relations $\Rightarrow$ never all in $\frac{1}{2}+\mathbb{Z}$ (Prop. P)

cell of c at v_c (odd, filled), offset at u_c (even, open); layer $\lambda'_c\mu_d/(u_c - v_d)$: all ϑ_{cd} odd $\Rightarrow$ $\frac{1}{2}+\mathbb{Z}$ possible (Lem. Hs)

(b) the geometric layer: $F_0[c][d] = \lambda'_c\mu_d/\vartheta_{cd} \in \frac{1}{2}+\mathbb{Z}$ $(c \neq d)$

$\vartheta_{cd} = u_c - v_d$ (all odd)

	0	1	2	3	4
0	−7	−9	−1	−5	−3
1	−5	−7	1	−3	−1
2	−3	−5	3	−1	1
3	−1	−3	5	1	3
4	3	1	9	5	7

$2F_0$ (all odd)

	0	1	2	3	4
0	−147	5	45	−9	3
1	3	−61	−15	−5	3
2	5	3	−53	−15	−3
3	−15	−5	3	−61	1
4	15	45	5	−9	−45

row c, column d; the diagonal of F_0 (shaded) is free, not of Cauchy form

(c) $\mathcal{H}^{5*}_{2,p}$, $p = 10007$: xy-slice; D_5^+ across carries



cell grid lattice image actual point jitter shortest lattice vectors