The great tit HapMap project: a continental-scale analysis of genomic variation in a songbird

Lewis G. Spurgin[1,2]*, Mirte Bosse[3,4]*, The Great Tit HapMap Consortium, Veronika N. Laine[3], Martien A.M. Groenen[4], Ben C. Sheldon[2], Marcel E. Visser[3,4], Kees van Oers[3] and Jon Slate[5]**

1 School of Biological Sciences, University of East Anglia, Norwich Research Park, United Kingdom

2 Edward Grey Institute, University of Oxford, United Kingdom

3 Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Wageningen, the Netherlands

4 Animal Breeding and Genomics Centre, Wageningen University, the Netherlands

5 Department of Animal and Plant Sciences, University of Sheffield, United Kingdom

* These authors contributed equally to this manuscript

** **Correspondence**: j.slate@sheffield.ac.uk

Abstract

A major aim of evolutionary biology is to understand why patterns of genomic variation among populations and species. Widespread continental species are interesting models for studying how evolution shapes patterns of genomic divergence, as they are associated with recent and rapid colonisations, large population sizes and high levels of gene flow. We studied patterns of genomic variation among 28 great tit (*Parus major*) populations in Europe. We found that genome-wide variation was consistent with a recent colonisation across Europe from a single refugium in the Balkans and/or Turkey. The highest levels of genome-wide structure were found between island populations in Crete, Sardinia and Corsica, most likely as a result of bottlenecks and reduced gene flow among these populations. We then found that differentiation across the genome was highly heterogeneous, with clear "islands of differentiation", even among the most recently separated populations. These islands of differentiation were consistently found in regions of low recombination, suggesting that background selection can rapidly promote population differentiation among recently colonised populations. We also detected outlier regions that were unique to peripheral great tit populations, most likely as a result of recent directional selection at the range edges of this species. These "unique" outlier regions contained candidate genes for morphology, stress response and colouration, which both support previous research in this species and provide avenues for future investigation. Together, our results suggest that examining differentiation landscapes in the context of population history is a useful way for determining how natural selection shapes genomic variation in the wild.

Introduction

Since the first studies of allozyme variation in humans [1] and *Drosophila* [2,3], there has been great interest in explaining how evolutionary and ecological processes shape the patterns of genetic variation observed within and among natural populations. One focus of research and debate in this area has been on quantifying the roles of adaptive and neutral processes in explaining observed levels of genetic variation [4]. However, adaptation does not occur in isolation, but against a background of recombination, gene flow, and genetic drift. More recently there has been increased effort in understanding how these fundamental evolutionary forces operate in concert to generate and maintain the levels of genetic diversity commonly observed in natural populations [5,6].

The arrival of high-throughput sequencing and subsequent characterisation of genome-wide variation across large numbers of individuals has revealed that at the genetic level, patterns of variation and divergence among natural populations and species are highly heterogeneous [7]. A feature of these "genomic landscapes"

of divergence that has received particular attention is the presence of so-called "islands of differentiation"; outlier regions of the genome with high levels of divergence estimated from statistics such as F_{ST} or d_{xy} [7–10]. Initially these regions were termed "islands of speciation", and were thought to arise as a result of reduced gene flow in genomic regions associated with reproductive isolation [7,11]. Subsequent modelling and analysis has revealed that highly heterogeneous patterns of genomic divergence can occur in the complete absence of gene flow, as a result of recombination rate variation and linked selection [12,13]. In genomic regions of low recombination, selection for beneficial mutations (positive selection), or against deleterious mutations (background selection), will impact relatively large genomic regions as a result of high levels of linkage disequilibrium (LD) among sites. Selection within these regions reduces diversity within populations, and increases differentiation among them, resulting in "islands" of increased differentation that persist over evolutionary time [13,14]

Comparing patterns of genomic differentiation among sets of populations and/or species with different different levels of genome-wide divergence is a powerful way of disentangling the forces that shape variation among populations. Martin et al. [15] showed that, across multiple *Heliconius* butterfly populations and species, patterns of genomic variation were shaped by a combination of gene flow and selection, particularly in genomic regions harbouring genes involved in wing patterning. In contrast, Renaut et al. [9] showed that in *Helianthus* sunflowers, genomic architecture was the main driver of genomic differentiation across sets of populations. Similarly, recent research in birds has revealed that differentiation landscapes are conserved across populations, species and even across avian families, with the same islands of differentiation arising among populations of distantly related species [16–18]. This latter pattern appears to have arisen, at least in part, as a result of a highly conserved recombination landscape in birds, with background selection in regions of low recombination producing recurrent islands of differentiation [19].

It is now clear that the recombination landscape and linked selection are key drivers of genomic variation within and among populations. However, we are only just beginning to understand how this linked selection interacts with other evolutionary forces to shape patterns of differentiation across natural populations and species [19–23]. A recent, large-scale analysis in sticklebacks showed that outlier genomic regions were more likely to arise in low recombination regions when gene flow occurred between populations [24]. There is also a significant impact of divergence time; in very recently separated populations the differentiation landscape is most likely to reflect selective sweeps, then as divergence accumulates, genomic architecture is expected to play an increasingly important role [19].

Widespread continental species are excellent models for studying how demography and the environment shape genetic and phenotypic variation among populations, due to their large effective population sizes and

ecologically varied ranges. Patterns of genome-wide variation in continental species is typically shaped by post-glacial colonisation routes and contemporary gene flow [25,26], although local adaptation can also play a role in shaping patterns of genetic diversity [27]. Genomic studies of continental species have enabled accurate assessments of colobisation routes, and revealed adaptation to environmental conditions REFS. However, to our knowledge, no studies have examined how the landscape of genomic differentiation changes in the context of postglacial recolonisation of continental environments. Large-scale studies of genomic differentiation across widespread species may inform how divergence accumulates in low recombining regions among recently separated populations, as well as how selection, gene flow and genomic architecture interact to shape genomic divergence in the early stages of population separation.

Insight into a species' evolutionary history can be gained if genetic variation in that species is characterized across much of its geographical range. Cross-population comparisons of genetic variation can then be used to make inferences about phylogeography, levels of gene flow between populations and how adaptation to different environmental and ecological conditions occurs. Whole-genome resequencing and customized SNP genotyping arrays have made studies that characterize all, or much, of the genome tractable. The first large-scale studies were performed in humans i.e. The HapMap Projects [28–30] which characterized human genetic variation on different continents, with a view to determining the feasibility of association mapping studies. Similar studies have been conducted in agriculturally important species and their progenitors[31–33], and in model organisms [34,35]. More recently, there is a growing appreciation that HapMap-type studies are useful for studying signatures of selection and adaptation in natural populations of species with large effective population sizes and high levels of gene flow [36–38].

The European great tit (Parus major major) is an excellent model for ecological and evolutionary study [39]. A wealth of ecological data exists across multiple great tit populations [40–42], enabling informed hypotheses about selection to be tested in this system. Phyleogeographic research using mitochondrial DNA suggests that this species has recently expanded across its European range, possibly from a single refugium in the Balkans [43]. Contemporary populations are characterised by large effective population sizes and high levels of gene flow among populations, resulting in low levels of genetic differentiation [44,45]. However, these previous cross-population molecular studies have relied on a modest number of microsatellite loci and mtDNA, making the detection of genomic regions under selection impossible. The genome of the great tit has recently been sequenced 19, and a high density panel of SNP markers has been developed [46]. A recent study of two European populations using this marker panel suggests that rapid adaptation has occurred at the genomic and phenotypic levels [47]. In this paper we describe how patterns of genetic variation and selection vary across populations by performing a HapMap study of over 600 unrelated individuals across 28 populations

using a high density SNP array with almost 500K SNPs.

In this study we examined how genomic architecture, natural selection and population history have shaped patterns of genomic variation across recently colonised European great tit populations. Using a large SNP panel typed across 28 populations (Fig. 1), we first characterise genome-wide patterns of variation within and among populations, in order to infer population history. We then examine how variation is partitioned across the genome, and test the hypothesis that highly divergent genomic regions have arisen in regions of low recombination [12,13]. Finally, we examine how genomic divergence accumulates along the colonisation route of this species, with the aim of inferring how recent natural selection and background selection drive variation across the genome in the wild.



Figure 1 Sampling locations of great tit populations used in the present study.

Results and Discussion

Genetic diversity and population history

Sampling locations and sample sizes for each population are given in Table 1. Levels of genetic diversity (π_{SNP}) were generally high, but we observed substantial differences among populations (Fig. 2A). Similarly, LD declined rapidly in all populations, reaching a baseline levels within ~5kb in all populations, but also varying among populations (Fig. S1). Highest levels of LD (and lowest levels of genetic diversity) were observed in the island populations of Crete (Greece), Sardinia (Italy), Corsica (France) and Vlieland (Netherlands) with lowest levels of LD in central Europe, Turkey and Spain (Fig. S1). This is consistent with reduced effective population size and reduced gene flow in these populations, either as result of the colonisation process, or more recent bottlenecks [48].

Genome-wide F_{ST} between European great tit populations was 0.007, with no significant pattern of isolation-by-distance (Mantel test; r = 0.14, p = 0.16). Instead, the highest levels of F_{ST} were found in comparisons involving the island populations of Corsica, Sardinia and Crete (Fig. S2). Admixture analysis was consistent with this pattern (Fig. S3); the $K = 2$ analysis assigned individuals in Sardinia and Corsica to one genetic

cluster, and the remaining populations to the second. Thus, it is likely that much of the genetic structure between European great tit populations is a result of genetic drift in small island populations. Admixture also revealed some structure between (mainly peripheral) mainland and larger island populations. At $K = 3$ (the model that best fitted the great tit data; Fig. S3), Spain was separated from the rest of mainland Europe. Increasing values of K resulted of the separation of populations in Scotland ($K = 4$), Sardinia (from Corsica; $K = 5$), France ($K = 6$), Crete ($K = 7$) and England ($K = 8$). The Admixture output at $K = 8$ is displayed in Fig. 2C; further increases in K did not generate patterns of structure that corresponded to geographical variation (Fig. S3), and were increasingly less well supported (Fig. S4). Thus, even with hundreds of thousands of markers Admixture was unable to separate many of the central European populations, confirming that levels of divergence are extremely low. PCA largely corroborated the Admixture results, with PC1 separating Corsical and Sardinia from the remaining populations, PC2 separating Spain, PC3 separating Corsical from Sardinina, and PC4 separating Scotland, England and France (Fig. S5).

Maximum likelihood analyses implemented in TreeMix showed that a model with no migration explained 97.8% of variance in relatedness between populations [49]; increasing the number of migration events substantially improved the percentage of relatedness explained, up to 99.7% when 10 migration events were fitted (Fig. S6). In Figure 3 we display the maximum likelihood tree with two migration events, after which the variance in relatedness explained plateaued when more migration events were added (Fig. S6). The tree was generally characterised by short branch lengths, with the exception of the island populations of Sardinia and Crete, which were grouped with the population from mainland Italy. Thus, the TreeMix analysis is consistent with the pattern of low overall genomic divergence, with the exception of island poppulations. However, much of the grouping that did occur among continental populations made geographical sense, with populations from Finland and Estonia grouped together, as were populations from Turkey and the Balkans, and populations from England and Scotland (Fig. 3). Interestingly, TreeMix grouped the Spanish and Corsican populations, which is consistent with previous subspecies descriptions of European great tits [50]. The two fitted migration edges both involved Sardinia, involving migration from eastern Europe, and to Corsica (Fig. 3).

We next tested the hypothesis that European great tits colonised from a single refugium in Turkey and the Balkans. This scenario has been suggested before [43], but due to the lwo number of genetic markers available there has been limited power with which to test this hypothesis. Using our genome-wide panel of SNP markers, we compared genetic and geographic distance between each population and the proposed refugial populations. Becasuse of the elevated structure in Corsica, Sardinia and Crete (Fig. S2), we excluded comparisons involving these populations. We found that F_{ST} was significantly related to distance from Turkey ($r = 0.84$, $p < 0.001$; Fig. 2C) and the Balkans ($r = 0.47$, $p < 0.001$). The same relationship was not

found for alternative potential refugial populations in Spain ($r = -0.07$, $p = 0.65$), or Italy ($r = 0.15$, $p = 0.49$). Our results therefore lend empirical support to the hypothesis [43] that great tits colonised Europe from a single refugium in the southeast.



Figure 2 Genetic diversity and structure in European Great tit populations. **A** Nucleotide diversity within each population. **B** Pairwise F_{ST} in relation to geographic distance from the Turkey, only including comparisons involving Turkey. **C** Output from admixture analysis at $K = 8$.



Figure 3 Maximum likelihood tree inferred by TreeMix, allowing ten migration events. Migrations events (arrows) are coloured according to their weight (red = higher migration), and horizontal branch lengths are proportional to the amount of genetic drift that has occurred along the branch.

Genomic landscapes of differentiation

It is likely that the many, and perhaps the majority, of wild populations are characterised by highly heterogeneous patterns of differentiation across the genome [23]. To examine how landscapes of genomic divergence have formed along the colonisation route of European great tits, we calculated windowed F_{ST} in 500kb bins between each population and the proposed refugial population in Turkey. We found that F_{ST} varied markedly across the genome in all comparisons (Fig. 4; Fig. S7). Outlier regions (windows with $zF_{ST} > 10$, where zF_{ST} is F_{ST} expressed relative to the genome-wide average, in standard deviations) were found in all comparisons apart from Crete and Sardinia, where overall levels of divergence were highest (Fig. 4). Our results suggest, therefore, that islands of differentiation arise even among populations that could be considered panmictic.

Genomic differentiation with Turkey was negatively related to recombination rate in almost all populations (Fig. 5). The relationship between F_{ST} and recombination rare was generally weak, with correlation coefficients < 0.1. In a handful of populations this relationship was substantially stronger - most notably in the island populations of Corsica, Sardinia and Crete, and in England, Scotland and Spain (Fig. 5). Outlier regions of very high differentiation ($zF_{ST} > 10$) almost exclusively occurred in relatively low recombining regions, and accordingly: the recombination rate of outlier regions was significantly lower than the genome-wide average (Wilcoxon test, P 0.130), suggesting that linked selection in regions of low recombination plays a key role in driving patterns of genomic differentiation even in the very earliest stages of population separation.

Of the 11 outlier regions, nine were found in only one or two comparisons, while the other two were found in 12 and 10 comparisons, respectively (Table S2). We hereafter refer to outlier regions found in one or two comparisons as "unique" outlier regions, and to those found in more than two comparisons as "shared" regions. Unique regions tended to be found in the most peripheral European great tits, with three found in Scotland, two in England, Spain and Finland; the remaining outlier regions were found in comparisons involving the Czech republic, Russia, Vlieland (Netherlands) and Belgium (Table S2). There appeared to be no difference in the recombination rate between shared and unique outlier regions (Fig. 5), although the small number of regions precluded testing this hypothesis formally.

It is difficult to differentiate between positive and background selection in producing heterogeneous patterns of differentiation across the genome. Correlated patterns of genomic differentiation among independent sets of populations suggests that background selection likely plays a role [16–18]. It appears that European great tits have colonised from a single refugial population, and as such truly independent comparisons are not available. Therefore an outlier region shared among multiple populations could represent either selection in

the ancestral population, or background selection. Given the pervasiveness of background selection in birds [17,18], and other organisms [51], and the overall negative relationship between F_{ST} and recombination rate in this study (Fig. 5), it is likely that this background selection is the driver of the shared outlier regions in great tits. Thus, it appears that background selection can generate islands of differentiation in the very earliest stages of population separation. This is not necessarily what we would expect - it is often assumed that peaks of high differentiation in recently separated populations are the product of selective sweeps, and that only after time do correlated patterns of genomic divergence arise [19,24]. Further research into genomic landscapes of differentiation among widespread continental species will shed further light on the role of background selection in shaping genomic divergence over short evolutionary timescales.



Figure 4 Landscapes of genomic differentiation in European great tit populations. F_{ST} across the genome is averaged in 500kb windows, with each panel displaying a pairwise comparison between with the proposed refugial population in Turkey. Red lines represent F_{ST} outliers (windows with mean F_{ST} values at least 10 standard deviations greater than the global mean for that comparison) shared across more than two comparisons (solid red lines), or unique to one or two comparisons (dashed red lines). Standardised differentiation (zF_{ST}) is displayed in Fig. S7.



Figure 5 Genomic differentiation and recombination rate variation in European great tit populations. Each point is a 500kb window, with each panel displaying a pairwise comparison between with the proposed refugial population in Turkey. Coloured points represent F_{ST} outliers (mean zF_{ST} values > 10) shared across more than two comparisons (red), or unique to one or two comparisons (dark blue).

To further explore how selection may have shaped variation in F_{ST} outlier regions, we estimated levels of nucleotide and haplotype diversity within these regions. Nucleotide diversity (π_{SNP}) in outlier regions varied from 0.28 to 0.41, and diversity in these regions was significantly lower than the genome-wide average (Wilcoxon test, $P = 0.007$; Fig. S8A). However, there appeared to be no difference in nucleotide diversity between shared and unique regions (Fig. S8A). Haplotype diversity varied substantially among regions, with haplotype richness ranging from 72 to 1033. Both haplotype richness and marker density in shared regions tended to be lower than those in unique regions (Fig. S8B). A detailed examination of haplotype structure in these regions is displayed in Figure S9. Although more distinct regions are needed to properly test underlying differences in haplotype structure, we provide a detailed examination of haplotype structure in two regions typically containing a signal in 1 or multiple regions (Fig S9). The frequency of a single haplotype was much higher in Oulu than in other populations, indicating a population-specific selective sweep. However, the region on chromosome 2 containing a signal in multiple populations is much less complex, demonstrating higher haplotype frequencies across a multitude of populations.

Our data suggest that examining patterns of nucleotide and haplotype diversity in outlier regions may help to separate recent episodes of positive selection from the gradual process of background selection (Figs S8, S9). Population comparisons allow for the identification of haplotypes at high frequency in single populations, a patterns that is unexpected by chance giving the high background in haplotype diversity in other populations. By contrast, the genomic architecture of regions that contain a high level of differentiation in multiple populations indeed seems to be less complex. We stress that marker density, haplotype richness and the frequency of the most common haplotype provide important insight into the efficiency of selection. Comparing these statistics in multiple populations enable the distinction between recent positive selection and drift effects. However, as we only detected a small number of outlier regions we are unable to make a detailed comparison between shared and unique regions. Further, simulations suggest that haplotype-frequency methods for detecting selection can be sensitive to background selection [52]. Research which combines multiple populations to allow shared and unique outlier regions to be identified, alongside a range of haplotype and SNP based methods for detecting selection, may allow us to identify the roles of positive and background selection in shaping variation.

Genes found within 500kb outlier windows are displayed in Table S2. Perhaps most notable among these is COL4A5, a gene found to be associated with bill length, and under selection between populations in England and the Netherlands, in a recent great tit study [47]. Here we found that the COL4A5 region is an F_{ST} outlier in England and Scotland, but not in any other European populations (Table S2). Another notable candidate gene potentially involved in beak morphology, and previously found to be under selection in UK

great tits is BMPR1A, which in this study was found in an outlier region in Scotland. Other genes in the unique outlier regions included potential candidates for morphology and obesity (PPP1CB [53], GHITM [54]), thermal stress response (CDKN1B [55]), malaria infection (MRPL33 [56]) and colour variation (SOX10 [57]).

Regions of high differentiation that are not shared among populations are more likely to be the result of recent positive selection [19]. We found evidence for such non-shared regions in European great tits, and that these tended to occur in peripheral populations. Observational and experimental research shows that adaptation at range edges is a key feature shaping divergence among recently colonised and expanding populations [58–60]. In great tits, previous research has shown evidence for adaptation in marginal populations. A comparison of Swedish and Finnish populations revealed genetic differences in basal metabolic rate, most likely an adaptive response to winter conditions [61]. Here we found outlier regions unique to populations in cold populations in Scotland, Finland and Russia, containing at least one candidate gene for thermal stress [55]. The detection of an outlier region containing the gene COL4A5 was particularly interesting, as this region was detected in a previous comparison of UK and Dutch great tits [47], in which significant associations with bill length and fitness were detected. Our results here are in accordance with the description of UK great tits as a separate subspecies [62], and combined with previous results suggest that this divergence is the result of recent natural selection in the UK [47]. Numerous other potential candidate genes were detected in these marginal populations, with potential roles in morphology, colouration, obesity and malaria infection. We are reluctant to speculate whether these candidate genes are genuine targets for natural selection, and more reluctant still to speculate as to how selection might be driving variation at these region. Regardless, these candidates will provide useful starting points for future genomic and ecological investigation.

Finally, it is worth noting that environmental association methods are highly suited method for detecting adaptation in widespread continental species [63,64], and further work will test how variation in the abiotic environment has shaped patterns of genomic variation in great tits. To our knowledge there has been little research on the potential for background selection to generate false positives in environmental association analyses, and given the pervasiveness of background selection, this should now be a priority. However, careful association of environmental and genotype data in species such as great tits, in which a wealth of ecological and genomic resources are available, is likely to generate interesting insights into the the genetic and phenotypic basis of natural selection.

Materials and Methods

Sampling and molecular methods

Samples were collected from 28 populations from 22 countries across Europe (Fig. 1; Table S1). Birds were trapped from nest boxes, or using mist nets, and ringed with a uniquely numbered aluminium ring. Blood was taken via brachial venipuncture, and stored in either 1 ml Cell Lysis Solution (Gentra Puregene Kit, Qiagen, USA) or Queen's buffer. All samples were genotyped using a custom-made Affymetrix® great tit 650K SNP chip at Edinburgh Genomics (Edinburgh, United Kingdom). After filtering, a total of 647 samples typed at 483888 SNPs were retained for analysis.

Analyses

Unless stated otherwise, all population genetic statistics were calculated in PLINK version 1.9 [65], and downstream analysis and plotting was carried out in R version 3.3 [66]. In each population, we estimated LD (R^2) for each pair of markers within 50kb on the same chromosome, and compared this to physical distance between marker pairs. For a simple comparison of short-range LD among populations, we also calculated, for each population, mean LD between all pairs of markers within 1kb. We calculated observed heterozygosity for each SNP and population using a reduced SNP dataset, which was pruned based on LD to remove all markers with $R^2 > 0.1$, then thinned with a probability of retaining each variant of 0.25.

We calculated genome-wide (mean) F_{ST} between each pair of populations using the pruned and thinned dataset described above. Pairwise F_{ST} was compared to geographic distance between populations using Mantel tests, implemented in the Ecodist package in R [67]. We tested whether genetic structure was related to distance from candidate refugial populations (in the Balkans, Turkey, Spain and Italy), using Pearson correlations. We also estimated population structure using Admixture version 1.3, with default settings [68]. We varied values of K from one to ten; by which point increasing values of K provided no informative information about population structure (see results). Model support for each value of K was estimated by calculating 5-fold cross-validation error. Finally, we visualised the evolutionary history among European great populations by generating a maximum likelihood tree in TreeMix version 1.13 [49]. We rooted the tree using a sample of *P. minor* individuals sampled from Amur, Russia. We fitted models allowing for range of migration events (0-10), and used a window size of 500 SNPs [49]. To assess model fit, we calculated the proportion of variance in relatedness between populations explained by each model [49].

We examined the genomic landscape of differentiation across European great tit populations by calculating F_{ST} in 500kb bins, using python scripts obtained from Github (https://github.com/simonhmartin/genomics_general). We did not estimate d_{xy}, as this parameter is difficult to estimate accurately from single SNP loci [13]. We also calculated standardised F_{ST} (hereafter ZF_{ST}) by mean-centring windowed values and dividing them by the standard deviation among windows. We defined outlier regions as 500kb bins with zF_{ST} values

greater than ten. We tested whether the landscape of genomic differentiation was related to recombination rate variation using a recombination map previously developed for the great tit using a 10K SNP chip [69], from which we estimated recombination rates in 500kb non-overlapping windows.

Acknowledgements

This work was supported by grants from the European Research Council (grant 202487 to J.S. and grant 339092 – E-Response to MEV) and Natural Environment Research Council (grant NE/J012599/1 to J.S.). LGS was supported by fellowships from the Edward Grey Institute for Ornithology and the BBSRC (BB/N011759/1). We thank Claire Bloor, Geoff Scopes and Alessandro Davassi of Affymetrix for their help during the chip design and genotyping calling processes. Richard Talbot and Alison Downing of Edinburgh Genomics provided the genotyping service.

References

1. Harris H. Enzyme polymorphisms in man. Proceedings of the Royal Society B: Biological Sciences. 1966;164: 298–310. doi:10.1098/rspb.1966.0032

2. Lewontin RC, Hubby JL. A molecular approach to the study of genic heterozygosity in natural populations. II. Amount of variation and degree of heterozygosity in natural populations of Drosophila pseudoobscura. Genetics. 1966;54: 595–609. doi:10.1111/j.1601-5223.1973.tb01163.x

3. Hubby JL, Lewontin RC. A molecular approach to the study of genic heterozygosity in natural populations. I. The number of alleles at different loci in Drosophila pseudoobscura. Genetics. 1966;54: 577–594.

4. Nei M. Selectionism and Neutralism in Molecular Evolution. Molecular Biology and Evolution. Oxford University Press; 2005;22: 2318–2342. doi:10.1093/molbev/msi242

5. Feder JL, Egan SP, Nosil P. The genomics of speciation-with-gene-flow. Trends in Genetics. Elsevier Current Trends; 2012;28: 342–350. doi:10.1016/J.TIG.2012.03.009

6. Lanfear R, Kokko H, Eyre-Walker A. Population size and the rate of evolution. Trends in Ecology & Evolution. Elsevier Current Trends; 2014;29: 33–41. doi:10.1016/J.TREE.2013.09.009

7. Turner TL, Hahn MW, Nuzhdin SV, Mitchison G, Spencer F. Genomic Islands of Speciation in Anopheles gambiae. Barton N, editor. PLoS Biology. Cambridge University Press; 2005;3: e285.

doi:10.1371/journal.pbio.0030285

8. Nadeau NJ, Whibley A, Jones RT, Davey JW, Dasmahapatra KK, Baxter SW, et al. Genomic islands of divergence in hybridizing Heliconius butterflies identified by large-scale targeted sequencing. Philosophical Transactions of the Royal Society of London B: Biological Sciences. 2011;367. Available: http://rstb.royalsocietypublishing.org/content/367/1587/343.short

9. Renaut S, Grassa CJ, Yeaman S, Moyers BT, Lai Z, Kane NC, et al. Genomic islands of divergence are not affected by geography of speciation in sunflowers. Nature Communications. 2013;4: 1827. doi:10.1038/ncomms2833

10. Poelstra JW, Vijay N, Bossu CM, Lantz H, Ryll B, Müller I, et al. The genomic landscape underlying phenotypic integrity in the face of gene flow in crows. Science. 2014;344. Available: http://science.sciencemag.org/content/344/6190/1410

11. Wolf JBW, Ellegren H. Making sense of genomic islands of differentiation in light of speciation. Nature Reviews Genetics. 2017;18: 87–100. doi:10.1038/nrg.2016.133

12. Noor MAF, Bennett SM. Islands of speciation or mirages in the desert? Examining the role of restricted recombination in maintaining species. Heredity. 2009;103: 439–444. doi:10.1038/hdy.2009.151

13. Cruickshank TE, Hahn MW. Reanalysis suggests that genomic islands of speciation are due to reduced diversity, not reduced gene flow. Molecular Ecology. 2014;23: 3133–3157. doi:10.1111/mec.12796

14. Turner TL, Hahn MW. Genomic islands of speciation or genomic islands and speciation? Molecular Ecology. 2010;19: 848–850. doi:10.1111/j.1365-294X.2010.04532.x

15. Martin SH, Dasmahapatra KK, Nadeau NJ, Salazar C, Walters JR, Simpson F, et al. Genome-wide evidence for speciation with gene flow in Heliconius butterflies. Genome Research. Cold Spring Harbor Laboratory Press; 2013;23: 1817–28. doi:10.1101/gr.159426.113

16. Burri R, Nater A, Kawakami T, Mugal CF, Olason PI, Smeds L, et al. Linked selection and recombination rate variation drive the evolution of the genomic landscape of differentiation across the speciation continuum of Ficedula flycatchers. Genome research. Cold Spring Harbor Laboratory Press; 2015;25: 1656–65. doi:10.1101/gr.196485.115

17. Vijay N, Weissensteiner M, Burri R, Kawakami T, Ellegren H, Wolf JBW. Genomewide patterns of variation in genetic diversity are shared among populations, species and higher-order taxa. Molecular Ecology.

2017;26: 4284–4295. doi:10.1111/mec.14195

18. Van Doren BM, Campagna L, Helm B, Illera JC, Lovette IJ, Liedvogel M. Correlated patterns of genetic diversity and differentiation across an avian family. Molecular Ecology. 2017;26: 3982–3997. doi:10.1111/mec.14083

19. Burri R. Dissecting differentiation landscapes: a linked selection's perspective. Journal of Evolutionary Biology. 2017;30: 1501–1505. doi:10.1111/jeb.13108

20. Ellegren H, Wolf JBW. Parallelism in genomic landscapes of differentiation, conserved genomic features and the role of linked selection. Journal of Evolutionary Biology. 2017;30: 1516–1518. doi:10.1111/jeb.13113

21. Jiggins CD, Martin SH. Glittering gold and the quest for Isla de Muerta. Journal of Evolutionary Biology. 2017;30: 1509–1511. doi:10.1111/jeb.13110

22. Lohse K. Come on feel the noise - from metaphors to null models. Journal of Evolutionary Biology. 2017;30: 1506–1508. doi:10.1111/jeb.13109

23. Ravinet M, Faria R, Butlin RK, Galindo J, Bierne N, Rafajlović M, et al. Interpreting the genomic landscape of speciation: a road map for finding barriers to gene flow. Journal of Evolutionary Biology. 2017;30: 1450–1477. doi:10.1111/jeb.13047

24. Samuk K, Owens GL, Delmore KE, Miller SE, Rennison DJ, Schluter D. Gene flow and selection interact to promote adaptive divergence in regions of low recombination. Molecular Ecology. 2017;26: 4378–4390. doi:10.1111/mec.14226

25. Hewitt GM. Some genetic consequences of ice ages, and their role in divergence and speciation. Biological Journal of the Linnean Society. 1996;58: 247–276. doi:10.1006/bijl.1996.0035

26. Hewitt GM. Postglacial re-colonisation of European biota. Biological Journal of the Linnean Society. 1999;68: 87–112.

27. Orsini L, Vanoverbeke J, Swillen I, Mergeay J, De Meester L. Drivers of population genetic differentiation in the wild: isolation by dispersal limitation, isolation by adaptation and isolation by colonization. Molecular ecology. 2013;22: 5983–99. doi:10.1111/mec.12561

28. Altshuler D, Donnelly P. A haplotype map of the human genome. Nature. Nature Publishing Group; 2005;437: 1299–1320. doi:10.1038/nature04226

29. Frazer KA, Ballinger DG, Cox DR, Hinds DA, Stuve LL, Gibbs RA, et al. A second generation human

haplotype map of over 3.1 million SNPs. Nature. 2007;449: 851–861. doi:10.1038/nature06258

30. Altshuler DM, Gibbs RA, Peltonen L, Altshuler DM, Gibbs RA, Peltonen L, et al. Integrating common and rare genetic variation in diverse human populations. Nature. 2010;467: 52–8. doi:10.1038/nature09298

31. Kijas JW, Lenstra JA, Hayes B, Boitard S, Porto Neto LR, San Cristobal M, et al. Genome-wide analysis of the world's sheep breeds reveals high levels of historic mixture and strong recent selection. Tyler-Smith C, editor. PLoS biology. 2012;10: e1001258. doi:10.1371/journal.pbio.1001258

32. Gibbs RA, Taylor JF, Van Tassell CP, Barendse W, Eversole KA, Gill CA, et al. Genome-wide survey of SNP variation uncovers the genetic structure of cattle breeds. Science (New York, NY). 2009;324: 528–32. doi:10.1126/science.1167936

33. Chia J-M, Song C, Bradbury PJ, Costich D, Leon N de, Doebley J, et al. Maize HapMap2 identifies extant variation from a genome in flux. Nature genetics. 2012;44: 803–7. doi:10.1038/ng.2313

34. Lindblad-Toh K, Wade CM, Mikkelsen TS, Karlsson EK, Jaffe DB, Kamal M, et al. Genome sequence, comparative analysis and haplotype structure of the domestic dog. Nature. 2005;438: 803–819. doi:10.1038/nature04338

35. Kirby A, Kang HM, Wade CM, Cotsapas C, Kostem E, Han B, et al. Fine mapping in 94 inbred mouse strains using a high-density haplotype resource. Genetics. 2010;185: 1081–95. doi:10.1534/genetics.110.115014

36. Horton MW, Hancock AM, Huang YS, Toomajian C, Atwell S, Auton A, et al. Genome-wide patterns of genetic variation in worldwide Arabidopsis thaliana accessions from the RegMap panel. Nature Genetics. 2012;44: 212–216. doi:10.1038/ng.1042

37. Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J, Borgwardt KM, et al. 1,135 Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis thaliana. Cell. 2016;166: 481–491. doi:10.1016/j.cell.2016.05.063

38. Williamson RJ, Josephs EB, Platts AE, Hazzouri KM, Haudry A, Blanchette M, et al. Evidence for widespread positive and negative selection in coding and conserved noncoding regions of Capsella grandiflora. Nachman MW, editor. PLoS genetics. 2014;10: e1004622. doi:10.1371/journal.pgen.1004622

39. Gosler A. The great tit. Hamlyn Species Guides; 1993.

40. Visser ME, Noordwijk AJv, Tinbergen JM, Lessells CM. Warmer springs lead to mistimed reproduction in great tits (Parus major). Proceedings of the Royal Society B: Biological Sciences. The Royal Society;

1998;265: 1867–1870. doi:10.1098/rspb.1998.0514

41. Charmantier A, McCleery RH, Cole LR, Perrins C, Kruuk LEB, Sheldon BC. Adaptive phenotypic plasticity in response to climate change in a wild bird population. Science (New York, NY). American Association for the Advancement of Science; 2008;320: 800–3. doi:10.1126/science.1157174

42. Dingemanse NJ, Bouwman KM, Pol M van de, Overveld T van, Patrick SC, Matthysen E, et al. Variation in personality and behavioural plasticity across four populations of the great tit Parus major. Journal of Animal Ecology. Blackwell Publishing Ltd; 2012;81: 116–126. doi:10.1111/j.1365-2656.2011.01877.x

43. Kvist L, Ruokonen M, Lumme J, Orell M. The colonization history and present-day population structure of the European great tit (Parus major major). Heredity. 1999;82: 495–502. Available: http://www.nature.com/hdy/journal/v82/n5/abs/6885130a.html

44. Kvist L, Martens J, Higuchi H, Nazarenko AA, Valchuk OP, Orell M. Evolution and genetic structure of the great tit (Parus major) complex. Proceedings Biological sciences / The Royal Society. 2003;270: 1447–54. doi:10.1098/rspb.2002.2321

45. Lemoine M, Lucek K, Perrier C, Saladin V, Adriaensen F, Barba E, et al. Low but contrasting neutral genetic differentiation shaped by winter temperature in European great tits. Biological Journal of the Linnean Society. 2016;118: 668–685. doi:10.1111/bij.12745

46. Kim J-M, Santure AW, Barton HJ, Quinn JL, Cole EF, Great Tit HapMap Consortium ME, et al. A high-density SNP chip for genotyping great tit (Parus major) populations and its application to studying the genetic architecture of exploration behaviour. Molecular ecology resources. 2018;18: 877–891. doi:10.1111/1755-0998.12778

47. Bosse M, Spurgin L, Laine V, Cole E, Firth J, Gienapp P, et al. Recent natural selection causes adaptive evolution of an avian polygenic trait. Science. 2017;358: 365–368. doi:10.1126/science.aal3298

48. James JE, Lanfear R, Eyre-Walker A. Molecular Evolutionary Consequences of Island Colonization. Genome Biology and Evolution. Oxford University Press; 2016;8: 1876–1888. doi:10.1093/gbe/evw120

49. Pickrell JK, Pritchard JK. Inference of Population Splits and Mixtures from Genome-Wide Allele Frequency Data. PLoS Genetics. 2012;8. doi:10.1371/journal.pgen.1002967

50. Clements J. The Clements checklist of the birds of the world. New York: Cornell University Press; 2007.

51. Comeron JM. Background selection as null hypothesis in population genomics: insights and challenges from Drosophila studies. Philosophical transactions of the Royal Society of London Series B, Biological

sciences. The Royal Society; 2017;372. doi:10.1098/rstb.2016.0471

52. Zeng K, Mano S, Shi S, Wu C-I. Comparisons of Site- and Haplotype-Frequency Methods for Detecting Positive Selection. Molecular Biology and Evolution. 2007;24: 1562–1574. doi:10.1093/molbev/msm078

53. Cho Y-L, Min J-K, Roh KM, Kim WK, Han BS, Bae K-H, et al. Phosphoprotein phosphatase 1CB (PPP1CB), a novel adipogenic activator, promotes 3T3-L1 adipogenesis. Biochemical and Biophysical Research Communications. Academic Press; 2015;467: 211–217. doi:10.1016/J.BBRC.2015.10.004

54. Migliano AB, Romero IG, Metspalu M, Leavesley M, Pagani L, Antao T, et al. Evolution of the Pygmy Phenotype: Evidence of Positive Selection from Genome-wide Scans in African, Asian, and Melanesian Pygmies. Human Biology. Wayne State University Press; 2013;85: 251–284. doi:10.3378/027.085.0313

55. Logan CA, Somero GN. Effects of thermal acclimation on transcriptional responses to acute heat stress in the eurythermal fish Gillichthys mirabilis (Cooper). American Journal of Physiology-Regulatory, Integrative and Comparative Physiology. American Physiological Society Bethesda, MD; 2011;300: R1373–R1383. doi:10.1152/ajpregu.00689.2010

56. Videvall E, Cornwallis CK, Palinauskas V, Valkiūnas G, Hellgren O. The Avian Transcriptome Response to Malaria Infection. Molecular Biology and Evolution. Oxford University Press; 2015;32: 1255–1267. doi:10.1093/molbev/msv016

57. Gunnarsson U, Kerje S, Bed'hom B, Sahlqvist A-S, Ekwall O, Tixier-Boichard M, et al. The Dark brown plumage color in chickens is caused by an 8.3-kb deletion upstream of SOX10. Pigment Cell & Melanoma Research. 2011;24: 268–274. doi:10.1111/j.1755-148X.2011.00825.x

58. Sexton JP, McIntyre PJ, Angert AL, Rice KJ. Evolution and Ecology of Species Range Limits. Annual Review of Ecology, Evolution, and Systematics. Annual Reviews; 2009;40: 415–436. doi:10.1146/annurev.ecolsys.110308.120317

59. Hill JK, Griffiths HM, Thomas CD. Climate Change and Evolutionary Adaptations at Species' Range Margins. Annual Review of Entomology. Annual Reviews; 2011;56: 143–159. doi:10.1146/annurev-ento-120709-144746

60. Weiss-Lehman C, Hufbauer RA, Melbourne BA. Rapid trait evolution drives increased speed and variance in experimental range expansions. Nature Communications. Nature Publishing Group; 2017;8: 14303. doi:10.1038/ncomms14303

61. Broggi J, Hohtola E, Orell M, Nilsson J-A. Local adaptation to winter conditions in a passerine spreading north: a common-garden approach. Evolution; international journal of organic evolution. 2005;59: 1600–3.

Available: http://www.ncbi.nlm.nih.gov/pubmed/16153046

62. Gosler AG. A comment on the validity of the British Great Tit Parus major newtoni. Bulletin of The British Ornithologists' Club. 1999;119: 47–55. Available: http://biostor.org/reference/111964

63. Coop G, Witonsky D, Di Rienzo A, Pritchard JK. Using environmental correlations to identify loci underlying local adaptation. Genetics. Genetics Society of America; 2010;185: 1411–23. doi:10.1534/genetics.110.114819

64. Frichot E, Schoville SD, Bouchard G, François O. Testing for associations between loci and environmental gradients using latent factor mixed models. Molecular biology and evolution. Oxford University Press; 2013;30: 1687–99. doi:10.1093/molbev/mst063

65. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, et al. PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. The American Journal of Human Genetics. 2007;81: 559–575. doi:10.1086/519795

66. R Development Core Team. R: A Language and Environment for Statistical Computing [Internet]. Team RDC, editor. 2011. doi:10.1007/978-3-540-74686-7

67. Goslee SC, Urban DL. The ecodist package for dissimilarity-based analysis of ecological data. Journal Of Statistical Software. 2007;22: 1–19. doi:citeulike-article-id:12008924

68. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in unrelated individuals. Genome Research. 2009;19: 1655–1664. doi:10.1101/gr.094052.109

69. Oers K van, Santure AW, De Cauwer I, Bers NE van, Crooijmans RP, Sheldon BC, et al. Replicated high-density genetic maps of two great tit populations reveal fine-scale genomic departures from sex-equal recombination rates. Heredity. Nature Publishing Group; 2014;112: 307–316. doi:10.1038/hdy.2013.107