

1 **Polyandry promotes successful colonisation in novel thermal envi-**  
2 **ronments**

3 Rebecca C. Lewis\*, Michael D. Pointer\*, Lucy Friend, Ramakrishnan Vasudeva, James Bemrose, Mathew  
4 J.G. Gage and Lewis G. Spurgin\*\*

5 School of Biological Sciences, University of East Anglia, Norwich Research Park, United Kingdom

6 \* These authors contributed equally to this work

7 \*\* **Correspondence:** L.Spurgin@uea.ac.uk

## Abstract

It is increasingly recognised that polyandry and sexual selection may enable successful colonisation of novel environments. In addition to the benefits provided by sexual selection, polyandry can promote successful colonisation through minimising the impact of mating with unsuitable males, and by reducing levels of inbreeding in founder populations. Here we show that polyandry provides substantial benefits in the colonisation of novel, and harsh, thermal environments. Using colonisation experiments with the model beetle *Tribolium castaneum*, we founded populations at increased temperature using either singly or doubly mated females, and followed population dynamics for ten generations. We found that extinction rates were significantly lower (9%) in polyandrous-founded populations compared to monogamous-founded populations (34%). Further, we found significant and consistently higher population sizes in polyandrous compared to monogamous-founded populations. Importantly, we identified distinct short-term and delayed benefits of multiple mating for population persistence and fitness, suggesting that polyandry acts to both buffer against mating with infertile males, and reduce inbreeding depression. Our results therefore suggest that polyandry provides substantial benefits for colonising populations, on top of those provided by sexual selection.

**Key words:** colonisation, extinction, population dynamics, sexual selection, *Tribolium*

## 23 Introduction

24 Global warming, when combined with other anthropogenic pressures, is placing a range of ecological and  
25 evolutionary stresses on wild populations [1,2]. In particular, a combination of habitat clearance and changing  
26 temperatures will result in populations simultaneously going through bottlenecks, while needing to respond  
27 to novel habitats and thermal regimes [3]. Understanding the behavioural, ecological, and genetic processes  
28 by which bottlenecked populations respond to novel thermal environments is therefore of relevance to  
29 understanding - and ultimately predicting - organismal responses to global warming [4]. Disentangling these  
30 processes requires a combination of laboratory and field research at a wide range of organisational levels [5].

31 One way that populations may be buffered against demographic and environmental change is through female  
32 multiple mating, or polyandry [6,7]. In addition to the direct benefits that polyandry may provide to females  
33 and their offspring (e.g. increased resources from males) [8], multiple mating can provide genetic benefits [9].  
34 Polyandry facilitates sexual selection, in which females may provide genetic benefits to their descendants  
35 through biasing paternity towards specific males [10]. There are multiple mechanisms by which sexual  
36 selection may improve population fitness [11], including offspring inheriting either “good” or “compatible”  
37 genes as a result of polyandry [12], or through selection for increased genetic diversity in offspring, which may  
38 increase adaptability to fluctuating environments [13].

39 Polyandry may also provide benefits to individuals and populations even in the absence of sexual selection.  
40 The best studied mechanism through which this may occur is bet-hedging [14]. If mate-choice is unreliable  
41 and there is variance in male fitness, indiscriminate mating with more than one male may be an effective  
42 strategy. It is still debated whether the fitness gains derived from bet-hedging are sufficient to drive the  
43 evolution of polyandry [15]. However, bet-hedging may play an important role in the dynamics of small  
44 colonising populations, in which the consequences of mating with a single, low-quality male are expected to  
45 be particularly severe [16].

46 Finally, polyandry may enable colonisation of new environments through short-term reductions in inbreeding.  
47 Cornell and Tregenza [17] developed a model showing that because offspring of polyandrous females contain  
48 half-sibs, inbreeding depression in future generations will be reduced, leading to greater probability of  
49 colonisation success. This theory received empirical support in a recent study of seed beetles (*Callosobruchus*  
50 *maculatus*), in which populations founded by polyandrous females had increased fitness after five generations  
51 compared to monogamous females [18]. Interestingly, despite the benefits of polyandry for individual fitness,  
52 Power and Holman [18] found no effect of mating treatment on extinction rates, which were low throughout  
53 the experiment. It is therefore not yet known how much of a benefit polyandry may provide in terms of

establishment probability in harsh environments. Given that inbreeding depression is most severe in poor environmental conditions [19], this is an important area for future investigation.

The red flour beetle *Tribolium castaneum* is an ideal model to test how polyandry influences colonisation success in a new environment. A pest of stored products, the ecology of *T. castaneum* is characterised by colonisation of empty habitats (e.g grain stores), presumably often by a small number of founders [20]. Experimental studies in this species have shown that founder effects have pronounced costs as a result of genetic and demographic effects, and that colonising populations are able to rapidly adapt to novel environments [21,22]. Further, this species is highly promiscuous and experimental evolution studies have shown that a history of sexual selection results in decreased risk of extinction under inbreeding, and in improved invasion of competitor populations [23,24]. A substantial proportion of matings in this species fail to result in offspring being produced, and there is some evidence to suggest that these costs are reduced when females mate multiply [25]. However, we do not yet know how polyandry affects colonisation success in a novel environment.

The aim of this study is to test how polyandry affects colonisation dynamics in a novel thermal environment. We placed singly and doubly mated *T. castaneum* females into an empty habitat at high temperature, and allowed populations to grow for ten generations. We first tested the hypotheses that, i) populations founded from polyandrous females were less likely to go extinct, and ii) polyandrous populations exhibited larger population sizes due to increased reproductive fitness. We use these findings to determine how mating strategy and inbreeding interact to affect colonisation dynamics in novel thermal environments.

## Materials and Methods

### Experimental protocols

All beetles were of our Krakow Super-strain (KSS) [26] and were maintained both before and throughout the experiment on a fodder medium consisting of 90% organic strong white bread flour mixed with 10% brewer's yeast, and topped with a thin layer of oats for traction.

Founding females and their mates were reared separately and mated under standard conditions of 30°C and 60% humidity. To allow matings to occur, pairs were placed into small (7ml) screw-top vials containing 1.5g of fodder. All females received two mating opportunities, each lasting 24 hours. In the first round of pairings, virgin females were randomly paired with virgin males (aged ~7 days post-eclosion). In the second round of pairings, half of the females from the polyandrous treatment were paired with a second, non-virgin male (hereafter referred to as the 'polyandrous' treatment). The remaining females were assigned to a 'monogamous'

84 treatment, in which they were re-paired with the same male, who, for consistency between treatments, was  
85 briefly removed from the dish before being replaced.

86 Using the above mating treatments, we established populations in a novel thermal environment for the  
87 remainder of the experiment. Specifically, after 48 hours of mating with either one or two males, females were  
88 transferred to a population container (100 ml PVC screw-cap containers, with the caps pierced for ventilation,  
89 containing 70 ml fodder) and left to oviposit for 7 days at 38°C and 60% humidity, after which they were  
90 removed and the offspring left to develop. This temperature is at the upper limit at which *T. castaneum*  
91 can reproduce, and constitutes a highly stressful environment [27]. All population containers post-mating  
92 were marked only with a randomised ID number so that experimental treatment was unknown by researchers  
93 during subsequent handling. Twenty-eight days after females were removed, the first generation of offspring  
94 were separated from the fodder by sieving, the fodder was discarded and the container and sieve cleaned with  
95 ethanol between replicates. The number of live adults was counted and placed into fresh fodder to seed the  
96 next generation. If >100 individuals were present, 100 were retained to mate in the next generation and the  
97 remainder discarded after counting (in order to minimise density-dependent effects). This new generation of  
98 adults was allowed to mate and lay on the fresh fodder for seven days, after which adults were removed by  
99 sieving and the offspring again left to develop for 28 days. This process was repeated for 10 generations, all  
100 at 38°C.

## 101 Statistical analyses

102 All analyses were carried out using R version 3.3.3 [28]. We separately modelled how the experimental mating  
103 treatment affected i) the probability of extinction over ten generations, and ii) changes in population size  
104 over the same period. For the extinction analysis, we used Cox proportional hazards models, implemented in  
105 the Survival package [29] in R. Because some populations went extinct in the first generation, possibly as a  
106 result of failure to mate and/or conceive, we ran the survival models both with and without populations that  
107 went extinct in the first generation.

108 To model how population size changed over time, we used generalised linear mixed models, implemented  
109 using the glmmADMB package [30] in R. For this analysis we only included population counts above  
110 zero. Offspring number per generation was modeled as a response variable with a negative binomial error  
111 distribution, and generation and experimental treatment (monogamous vs polyandrous founder) were fitted as  
112 explanatory variables, alongside the interaction between treatment and generation. To account for potential  
113 non-linear changes in population size over time, we fitted changes in offspring numbers over generations as  
114 i) a continuous variable, and ii) a third-order polynomial. We fitted random slopes models, which allowed

variation among individual populations over generations. Finally, we tested for a difference in population size between experimental treatments in the first and last generation, using two separate generalized linear models (as above but with no random effects), implemented using the MASS package [31] in R.

## Results

We tracked a total of 53 monogamous and 55 polyandrous *T. castaneum* populations in a novel thermal environmental until extinction, or for 10 generations. Overall dynamics of the populations are given in Figure 1; though there was a general trend for increasing population size, there were substantial fluctuations over generations, with decreases in population size between generations three and four, and between generations six and seven (Fig. 1). Despite this, we observed a clear and consistent trend of larger adult population size in populations founded by polyandrous compared to monogamous females (Fig. 1).

For statistical comparison of our experimental treatments, we separately tested for differences in extinction rates (i.e. did time to extinction differ between treatments), and population size changes over time. In the first generation, six populations founded by monogamous females went extinct (11%), while no populations founded by polyandrous females went extinct. By generation ten, 18 monogamous populations (34%) and five polyandrous populations (9%) were extinct (Fig. 2A). The effect of treatment on time to extinction was significant (Cox proportional hazards; hazard ratio = 0.256; 95% CIs = 0.102, 0.642;  $P = 0.004$ ). This effect remained significant after removal of populations that went extinct in the first generation (hazard ratio = 0.361; 95% CIs = 0.137, 0.950;  $P = 0.039$ ).

We next tested how founder mating regime affected fitness and population trajectories. Excluding extinctions, we found no significant difference in the number of offspring produced by monogamous or polyandrous females in the first generation (GLM,  $P = 0.503$ ). Considering all generations, however, we found that populations founded from polyandrous females had higher overall population sizes than populations founded by monogamous females (Fig. 2B, Table 1). We also found that, when generation modeled as a continuous variable, population size increased over time, but that there was no interaction between treatment and generation (Fig. 2B, Table 1). The effect of experimental treatment was also significant when generation was modeled as a third-order polynomial ( $P = 0.019$ ). Finally, considering only populations that survived all ten generations, population size in polyandrous-founded populations in generation ten was significantly higher in monogamous-founded populations (GLM,  $P = 0.004$ ).

## Discussion

The costs of mating with multiple males are likely to outweigh any direct benefits gained, and as such the evolution and maintenance of polyandry are generally viewed as an evolutionary puzzle [32]. Potential indirect benefits of polyandry include i) enabling sexual selection, ii) enabling bet-hedging, and iii) reducing inbreeding load. Previous research using the *T. castaneum* model has found that long-term sexual selection results in increased population fitness [23,24]. Here we show that polyandry has substantial benefits in colonising populations, even when the opportunity for pre-copulatory sexual selection is reduced.

Perhaps the best-studied way in which females can increase the fitness of their offspring through polyandry is through bet-hedging [16]. By mating with multiple males, females may reduce the risk of being fertilised by an unsuitable male, and therefore increase offspring fitness. Bet-hedging is likely to be most beneficial when i) there is a substantial proportion of highly unsuitable (e.g. infertile) males in the population, and ii) the population is small [16]. In our study, populations were founded by a single female, and as such there is clear potential for polyandry to provide benefits. We found that 11% of monogamous females produced no offspring in the first generation, while all polyandrous females produced offspring. This is consistent with a situation in which a failure to produce offspring is the result of male infertility, under which we would only have expected 1.28% of random pairs of males to both be infertile. The mating conditions we imposed were short and artificial, and studies of natural populations are required. Nevertheless, previous research in *T. castaneum* has found that a substantial proportion of matings fail to result in offspring production [25,33], and across insects, high levels of male infertility have been observed in the wild [34]. It is therefore likely that multiple mating is an important mechanism for increasing short-term establishment probability in newly-colonised populations.

An additional mechanism through which polyandry can benefit colonising populations is through reducing levels of inbreeding in subsequent generations [17]. Consistent with this hypothesis, we found significantly lower population sizes and higher extinction rates in monogamous-founded populations over the full duration of our experiment. Population sizes fluctuated substantially over the course of our experiment, likely a result of density-dependent processes which are well-documented in *T. castaneum* [35]. Though the higher population sizes in polyandrous-founded populations were generally consistent over time, it is notable that the difference between treatments was highest when population sizes were high (i.e. in generations 2,3,8 and 9), and lowest when population sizes were reduced (i.e. generations 4 and 7). It is possible that a bet-hedging like scenario could explain these longer-term benefits of polyandry if there was substantial variation in fitness among fertile males, as multiple-mating would increase the chances of mating with at least one suitable

male [16]. However, this is unlikely to explain our results, as we found no difference in population size between treatments in the first generation, but instead only when considering later generations. Similarly, if post-copulatory sexual selection could explain some of the differences observed between our experimental treatments, we would expect to observe at least some differences in offspring fitness in the first generation. We therefore suggest that in *T. castaneum* and similar systems, polyandry benefits colonising populations through i) guarding against male infertility and enabling establishment, and ii) reducing inbreeding and enabling longer-term population persistence.

Our results are broadly consistent with a recent study in *C. maculatus*, in which the increased fitness in polyandrous-founded populations were observed in F4 and F5, but not F1-F3, generations [18]. It is worth noting that, unlike in our study, Power and Holman [18] found no effect of mating treatment on extinction, likely because of the environment was relatively benign, or because there was insufficient time for extinctions to occur. Here, through a longer-term experiment, we demonstrate that the benefits of polyandry can persist for substantial periods of time, and show that they are likely to be important for populations colonising novel environments.

Finally, it is worth noting that our study does not aim to address whether the benefits gained from either bet-hedging or inbreeding avoidance are sufficient to explain the evolution of polyandry [11,17]. Rather, we suggest that in the specific context of colonising novel environments, the benefits of these processes are likely to be substantial [16]. Future climate change is expected to result in species shifting their ranges and undergoing changes in population size, and there is increasing realisation that evolutionary processes need to be incorporated into predictive models of population and species responses to climate change [36]. The multiple, interacting benefits of polyandry should certainly be incorporated into such models in order to improve predictive power.

## Acknowledgements

This research was funded by a BBSRC Future Leader fellowship to LGS (BB/N011759/1), a UEA-funded PhD studentship to RL, BBSRC NRP DTP funded PhD studentships to MDP and JB. We thank Andreas Sutter for useful discussions about this work.



## References

1. Hoffmann AA, Sgrò CM. Climate change and evolutionary adaptation. *Nature*. nature.com; 2011;470: 479–485.
2. Pecl GT, Araújo MB, Bell JD, Blanchard J, Bonebrake TC, Chen I-C, et al. Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science*. 2017;355.
3. Hill JK, Griffiths HM, Thomas CD. Climate change and evolutionary adaptations at species' range margins. *Annu Rev Entomol*. 2011;56: 143–159.
4. Hoelzel AR. Looking backwards to look forwards: Conservation genetics in a changing world. *Conserv Genet*. 2010;11: 655–660.
5. Scheffers BR, De Meester L, Bridge TCL, Hoffmann AA, Pandolfi JM, Corlett RT, et al. The broad footprint of climate change from genes to biomes to people. *Science*. 2016;354.
6. Candolin U, Heuschele J. Is sexual selection beneficial during adaptation to environmental change? *Trends Ecol Evol*. 2008;23: 446–452.
7. Parrett JM, Knell RJ. The effect of sexual selection on adaptation and extinction under increasing temperatures. *Proc Biol Sci*. 2018;285.
8. Fedorka KM, Mousseau TA. Material and genetic benefits of female multiple mating and polyandry. *Anim Behav*. 2002;64: 361–367.
9. Zeh JA, Zeh DW. Reproductive mode and the genetic benefits of polyandry. *Anim Behav*. 2001;61: 1051–1063.
10. Andersson M, Iwasa Y. Sexual selection. *Trends Ecol Evol*. 1996;11: 53–58.
11. Yasui Y. The “genetic benefits” of female multiple mating reconsidered. *Trends Ecol Evol*. 1998;13: 246–250.
12. Neff BD, Pitcher TE. Genetic quality and sexual selection: An integrated framework for good genes and compatible genes. *Mol Ecol*. 2005;14: 19–38.
13. Garcia-Gonzalez F, Yasui Y, Evans JP. Mating portfolios: Bet-hedging, sexual selection and female multiple mating. *Proc Biol Sci*. 2015;282: 20141525.
14. Watson PJ. Multiple paternity as genetic bet-hedging in female sierra dome spiders, *linyphia litigiosa*

(linpyhiidae). Anim Behav. 1991;41: 343–360.

15. Holman L. Bet hedging via multiple mating: A meta-analysis. Evolution. 2016;70: 62–71.

16. Yasui Y, Garcia-Gonzalez F. Bet-hedging as a mechanism for the evolution of polyandry, revisited. Evolution. 2016;70: 385–397.

17. Cornell SJ, Tregenza T. A new theory for the evolution of polyandry as a means of inbreeding avoidance. Proc Biol Sci. 2007;274: 2873–2879.

18. Power DJ, Holman L. Polyandrous females found fitter populations. J Evol Biol. 2014;27: 1948–1955.

19. Armbruster P, Reed DH. Inbreeding depression in benign and stressful environments. Heredity. 2005;95: 235–242.

20. Dawson PS. Life history strategy and evolutionary history of tribolium flour beetles. Evolution. [Society for the Study of Evolution, Wiley]; 1977;31: 226–229.

21. Szucs M, Melbourne BA, Tuff T, Hufbauer RA. The roles of demography and genetics in the early stages of colonization. Proc Biol Sci. 2014;281.

22. Szucs M, Melbourne BA, Tuff T, Weiss-Lehman C, Hufbauer RA. Genetic and demographic founder effects have long-term fitness consequences for colonising populations. Ecol Lett. 2017;20: 436–444.

23. Lumley AJ, Michalczyk Ł, Kitson JJN, Spurgin LG, Morrison CA, Godwin JL, et al. Sexual selection protects against extinction. Nature. 2015;522: 470–473.

24. Godwin JL, Spurgin LG, Michalczyk Ł, Martin OY, Lumley AJ, Chapman T, et al. Lineages evolved under stronger sexual selection show superior ability to invade conspecific competitor populations. Evol Lett. 2018;2: 511–523.

25. Pai A, Bennett L, Yan G. Female multiple mating for fertility assurance in red flour beetles (tribolium castaneum). Can J Zool. NRC Research Press; 2005;83: 913–919.

26. Dickinson M. The impacts of heat-wave conditions on reproduction in a model insect, tribolium castaneum. PhD thesis, University of East Anglia; ueaeprints.uea.ac.uk. 2018.

27. Howe RW. THE EFFECTS OF TEMPERATURE AND HUMIDITY ON THE RATE OF DEVELOPMENT AND THE MORTALITY OF TRIBOLIUM CONFUSUM DUVAL (COLEOPTERA, TENEBRIONIDAE). Ann Appl Biol. 1960;48: 363–376.

28. Development Core Team RR. R: A language and environment for statistical computing. R foundation for

255 statistical computing ...; 2011;

256 29. Therneau TM. A package for survival analysis in s [Internet]. 2015. Available: [https://CRAN.R-project.](https://CRAN.R-project.org/package=survival)

257 [org/package=survival](https://CRAN.R-project.org/package=survival)

258 30. Fournier DA, Skaug HJ, Ancheta J, Ianelli J, Magnusson A, Maunder M, et al. AD Model Builder: Using

259 automatic differentiation for statistical inference of highly parameterized complex nonlinear models. *Optim*

260 *Methods Softw.* 2012;27: 233–249.

261 31. Venables WN, Ripley BD. Modern applied statistics with s [Internet]. Fourth. New York: Springer; 2002.

262 Available: <http://www.stats.ox.ac.uk/pub/MASS4>

263 32. Simmons LW. The evolution of polyandry: Sperm competition, sperm selection, and offspring viability.

264 *Annu Rev Ecol Evol Syst.* Annual Reviews; 2005;36: 125–146.

265 33. Tyler F, Tregenza T. Why do so many flour beetle copulations fail? *Entomol Exp Appl.* 2013;146:

266 199–206.

267 34. García-González F. Infertile matings and sperm competition: The effect of “nonsperm representation” on

268 intraspecific variation in sperm precedence patterns. *Am Nat.* The University of Chicago Press; 2004;164:

269 457–472.

270 35. Mertz DB. The tribolium model and the mathematics of population growth. *Annu Rev Ecol Syst.* Annual

271 Reviews; 1972;3: 51–78.

272 36. Lavergne S, Mouquet N, Thuiller W, Ronce O. Biodiversity and climate change: Integrating evolutionary


273 and ecological responses of species and communities. *Annu Rev Ecol Evol Syst.* Annual Reviews; 2010;41:

274 321–350.


# Tables and Figures

**Table 1** Summary of results from a generalised linear mixed model of population dynamics of experimental *T. castaneum* populations founded from monogamous or polyandrous females. Here, the ‘treatment’ estimate refers to the effect of polyandrous relative to monogamous females, and generation was modeled as a linear effect. As random effects we modeled population ID (Var = 0.12, SD = 0.35), Generation () and a random slope of population ID ().

	Estimate	SE	P
(Intercept)	4.719	0.083	< 0.001
Treatment	0.229	0.091	0.012
GenN	0.063	0.011	< 0.001



**Figure 1** Colonisation dynamics of experimental *T. castaneum* populations founded from monogamous or polyandrous females.



**Figure 2** Extinction and population dynamics of experimental *T. castaneum* populations founded from monogamous or polyandrous females. **A** Proportion of populations surviving over time; **B** number of adults in experimental populations. In **B**, thin lines represent individual populations, while the thick lines represent fitted values from a negative-binomial GLM.