

Relations

CS application: relational databases

student id	first name	last name
123	Bob	Smith

student id	passed course
123	CSCI 246
123	CSCI 127

Questions about data stored in relational databases can be posed precisely using the language of relations.

SQL (structured query language)

Def The cartesian product of two sets A, B is

$$A \times B = \{ (a, b) : a \in A \wedge b \in B \}$$

lists/tuples[↑]/arrays — order matters

ex

$\mathbb{R} \times \mathbb{R} = 2d \text{ plane, Cartesian plane}$

$$\{\text{red, blue}\} \times \{1, 2, 3\} = \{(\text{red}, 1), (\text{red}, 2), (\text{red}, 3), (\text{blue}, 1), (\text{blue}, 2), (\text{blue}, 3)\}$$

Q what is $|A \times B|$? $|A|, |B|$

$$|A| \cdot |B|$$

$$\mathbb{R} \times \mathbb{R} = \mathbb{R}^2$$

$$\mathbb{R} \times \mathbb{R} \times \mathbb{R} = \mathbb{R}^3$$

Def A binary relation R on sets A, B is a subset $R \subseteq A \times B$.

We write $(x, y) \in R$ as $x R y$

$(x, y) \notin R$ as $x \not R y$

examples

① R , "is (blood) related to" is a binary relation on people.

let P be the set of all people

"is blood related to" is

$$\{(x, y) : x \in P \wedge y \in P \wedge x \text{ is related to } y\}$$

$(\text{serena williams}, \text{Venus williams}) \in R,$

$(\text{Lucy williams}, \text{serena williams}) \notin R.$

② $<$ on $A = \{1, 2, 3, 4\}$

$< = \{ (1, 2), (1, 3), (1, 4), (2, 3), (2, 4), (3, 4) \}$

$1 < 2$ but $3 \nless 2$

③ let $f: A \rightarrow B$ be a function

$\{ (a, f(a)) : a \in A \} \subseteq A \times B$, so it is
a relation

Q Is the converse true? let R
be a binary relation on A, B .

$\{ (x, y) : x \in A \wedge y \in B \wedge x R y \}$

$\Rightarrow f: A \rightarrow B$ s.t. $f(x) = y$

is a function

→ true or false?

④ let $A = \text{months}$, $B = \text{number of days}$

Relation: month, its # days

$\{ (\text{Jan}, 31), (\text{Feb}, 28), (\text{Feb}, 29), (\text{Mar}, 31) \dots \}$

Jan	31
Feb	28
Feb	29
Mar	31
\vdots	\vdots

Jan	\rightarrow 31
Feb	\rightarrow 28
Feb	\rightarrow 29
Mar	\rightarrow 31
\vdots	\vdots


⑤ $A = \{1, 2, 3, 4, 5\}$

$(1, 1) \in R_2$

$(2, 4) \in R_2$

$(3, 2) \notin R_2$

$R_2:$



Properties of relations

let $R \subseteq A \times A$, so R is a relation on A


$R: a_1 \rightarrow a_2$
 $\quad \searrow$
 $\quad a_3$

R is reflexive if $\forall a \in A: a R a$
all nodes have self-loops

R is irreflexive if $\forall a \in A: a \not R a$
no nodes have self-loops

R is symmetric if




$$\forall a_1, a_2 \in A: a_1 R a_2 \Rightarrow a_2 R a_1$$



Whenever we have a forward edge, we have the backward edge.

R is anti-symmetric if $\{(a,b), (b,c), (c,a)\}$

$$\forall a_1, a_2 \in A: (a_1 R a_2 \wedge a_2 R a_1) \Rightarrow a_1 = a_2$$




never have backwards edges, but self-loops okay.

R is transitive if

$$\forall \underline{a}, \underline{b}, \underline{c} \in A : (\overset{T}{\underline{a} R \underline{b} \wedge \underline{b} R \underline{c}}) \Rightarrow (\overset{F}{\underline{a} R \underline{c}})$$

$\underline{a_1 R a_2 \wedge a_2 R a_1}$ $\underline{a_1 R a_1}$




Shortcut edges always exist

Q Is $a_1 \rightarrow a_2$ transitive? $a_1 \neq a_2$

Let $\underline{a = a_1}$

$b = a_2$

$\underline{c = a_1}$



Q Is a_1 transitive?

$$\overset{T}{(a_1 R a_1 \wedge a_1 R a_1)} \Rightarrow \overset{T}{(a_1 R a_1)}$$

Relations review




let A, B be sets. $(a, b) \in R$
 $a R b$

$R \subseteq A \times B$ is a binary relation

often, we are concerned with relations over a single set:

$R \subseteq S \times S$ " R is a relation on S "

Properties of relations on single sets:

- reflexive: $\forall a \in A: a R a$ 
- irreflexive: $\forall a \in A: a \not R a$
- symmetric: $\forall a_1, a_2 \in A: a_1 R a_2 \Rightarrow a_2 R a_1$ 
- anti-symmetric: $\forall a_1, a_2 \in A: (a_1 R a_2 \wedge a_2 R a_1) \Rightarrow a_1 = a_2$ 
- transitive: $\forall a_1, a_2, a_3 \in A: (a_1 R a_2 \wedge a_2 R a_3) \Rightarrow a_1 R a_3$

a

b

symmetric ✓
anti-symmetric ✓

$$A = \{a, b\}$$

$$\mathcal{R} \subseteq A \times A$$

$$B \subseteq \text{People} \times \text{People}$$

$$(\text{Lucy}, \text{Britney Spears}) \in B$$

Lucy B Britney Spears

Lucy $\not B$ Braeden

$$> \subseteq \mathbb{R} \times \mathbb{R}$$

$$2 > 1.5$$

$$2 \not> 2$$

ex relation $<$ on \mathbb{Z} :

- reflexive?

no - disproof by counterexample.

$$1 \in \mathbb{Z}. \quad 1 \not< 1$$

- irreflexive? yes.

$$\forall a \in \mathbb{Z}: a \not< a.$$

let $a \in \mathbb{Z}$. $a \not< a$ because no integer is less than itself. \square

- symmetric?

disproof by counterexample:

$$1 < 2 \text{ but } 2 \not< 1.$$

- antisymmetric? $\forall a_1, a_2 \in A:$
 $(a_1 < a_2 \wedge a_2 < a_1) \Rightarrow a_1 = a_2$

let $a_1, a_2 \in \mathbb{Z}$. Assume $a_1 < a_2$ and $a_2 < a_1$.

Since no a_1, a_2 satisfy $a_1 < a_2$ and $a_2 < a_1$,
 $(a_1 < a_2 \wedge a_2 < a_1) \Rightarrow a_1 = a_2$ is
vacuously true.

• transitive?

$$\forall a_1, a_2, a_3 \in A: (a_1 R a_2 \wedge a_2 R a_3) \Rightarrow a_1 R a_3$$

Proof Assume $a_1, a_2, a_3 \in \mathbb{Z}$ and
 $a_1 < a_2$ and $a_2 < a_3$. By the
def. of $<$, $a_1 < a_3$

	$< \subseteq \mathbb{R} \times \mathbb{R}$	Same Bday $\subseteq P \times P$	Subset $\subseteq P(S) \times P(S)$
reflexive			
irreflexive			
symmetric			
anti-symmetric			
transitive			
equivalence relation			
partial order			
strict partial order			
total order			
strict total order			