

Relations

CS application: relational databases

student id	first name	last name
123	Bob	Smith

student id	passed course
123	CSCI 246
123	CSCI 127

Questions about data stored in relational databases can be posed precisely using the language of relations.

SQL (structured query language)

Def The cartesian product of two sets A, B is

$$A \times B = \{ (a, b) : a \in A \wedge b \in B \}$$

lists/tuples[↑]/arrays — order matters

ex

$\mathbb{R} \times \mathbb{R} = 2d \text{ plane, Cartesian plane}$

$$\{\text{red, blue}\} \times \{1, 2, 3\} = \{(\text{red}, 1), (\text{red}, 2), (\text{red}, 3), (\text{blue}, 1), (\text{blue}, 2), (\text{blue}, 3)\}$$

Q what is $|A \times B|$? $|A|, |B|$

$$|A| \cdot |B|$$

$$\mathbb{R} \times \mathbb{R} = \mathbb{R}^2$$

$$\mathbb{R} \times \mathbb{R} \times \mathbb{R} = \mathbb{R}^3$$

Def A binary relation R on sets A, B is a subset $R \subseteq A \times B$.

We write $(x, y) \in R$ as $x R y$

$(x, y) \notin R$ as $x \not R y$

examples

① R , "is (blood) related to" is a binary relation on people.

let P be the set of all people

"is blood related to" is

$$\{(x, y) : x \in P \wedge y \in P \wedge x \text{ is related to } y\}$$

$(\text{serena williams}, \text{Venus williams}) \in R,$

$(\text{Lucy williams}, \text{serena williams}) \notin R.$

② $<$ on $A = \{1, 2, 3, 4\}$

$< = \{ (1, 2), (1, 3), (1, 4), (2, 3), (2, 4), (3, 4) \}$

$1 < 2$ but $3 \nless 2$

③ let $f: A \rightarrow B$ be a function

$\{ (a, f(a)) : a \in A \} \subseteq A \times B$, so it is
 $\uparrow \quad \uparrow$
a relation

Q Is the converse true? let R
be a binary relation on A, B .

$\{ (x, y) : x \in A \wedge y \in B \wedge x R y \}$

$\Rightarrow f: A \rightarrow B$ s.t. $f(x) = y$

is a function

→ true or false?

④ let $A = \text{months}$, $B = \text{number of days}$

Relation: month, its # days

$\{ (\text{Jan}, 31), (\text{Feb}, 28), (\text{Feb}, 29), (\text{Mar}, 31) \dots \}$

Jan	31
Feb	28
Feb	29
Mar	31
⋮	⋮

Jan	→	31
Feb	→	28
Feb	→	29
Mar	→	⋮
⋮		⋮


⑤ $A = \{1, 2, 3, 4, 5\}$

$(1, 1) \in R_2$

$(2, 4) \in R_2$

$(3, 2) \notin R_2$

$R_2:$



Properties of relations

let $R \subseteq A \times A$, so R is a relation on A


$R: a_1 \rightarrow a_2$
 $\quad \searrow$
 $\quad a_3$

R is reflexive if $\forall a \in A: a R a$
all nodes have self-loops

R is irreflexive if $\forall a \in A: a \not R a$
no nodes have self-loops

R is symmetric if



$$\forall a_1, a_2 \in A: a_1 R a_2 \Rightarrow a_2 R a_1$$



Whenever we have a forward edge, we have the backward edge.

R is anti-symmetric if $\{(a,b), (b,c), (c,a)\}$

$$\forall a_1, a_2 \in A: (a_1 R a_2 \wedge a_2 R a_1) \Rightarrow a_1 = a_2$$



$$a \rightarrow b \rightarrow c \rightarrow a$$

never have backwards edges, but self-loops okay.

R is transitive if

$$\forall a, b, c \in A : (a R b \wedge b R c) \Rightarrow (a R c)$$

$a \rightarrow b \rightarrow c$ shortcut edges always exist \checkmark

~~$a R b \wedge c R b$~~

$a \rightarrow b \leftarrow c$

Q Is $a_1 \begin{matrix} \curvearrowright \\ \curvearrowleft \end{matrix} a_2$ transitive?