The *Catalogue Astrochimique de Sections Efficaces d'Ionisation* (CASEI): A Database Of Low Energy Total Ionisation Cross-Section For Astrochemical Purpose

Lilian Heydel[1]

[1]Centre de Recherche sur les Ions, les Matériaux et la Photonique Université Caen Normandie, GANIL, ENSICAEN, CNRS, CEA, Normandie Univ, CIMAP UMR6252, F-14000 Caen, France. **lilian.heydel@ganil.fr**

Total Ionisation Cross Section (TICS) reprents a certainly important factor for a large number of physical domains.

1 Introduction

Electron-driven ionisation plays a big role in planetary and interplanetary medium. The result of it the production of reactive radicals. One of the main parameter to quantify this phenomena is the cross-section of ionisation of the molecule, either partial (Partial Ionisation Cross Section, **PICS**) or total (Total Ionisation Cross Section, **TICS**). While it can be mesured experimentally (Rapp and Englander-Golden, 1965; Jiao, DeJoseph, and Garscadden, 2007, and reference therein), this is not necesseraly the case for every molecules. In fact, some molecules are either absent or under covered litterature-wise. Those include molecule of interest for planetary and exoplanetary science, like phosphine PH_3, found in the atmosphere of gas giants and in the interstellar medium (Burgasser et al., 2025; Larson et al., 1980; Ridgway, Wallace, and Smith, 1976), propaenitril C_3H_5N, found in Titan atmosphere (cite), or even brome bearing molecules such as ..., ..., ... The same can be said for isomers, as not all isomer of a define specie have the TICS calculated for all its isomer. It is, however, possible to compute this parameter. Total Ionisation Cross-Section (TICS) database established using computing method were already done, such as ALeCS (Gaches et al., 2024), NIFS (citer) or NIST (citer). They, howerever, lack the presence of some of those *exotic* specie, or are either complicated to acces for general public.
In this work, I present a database of TICS for 274 neutral and stable molecules made with open-source tools. This database is also available on a website hosted on GitHub and is publicly available[1]. All the content of the website can be downloaded freely.

2 Methodology

The methodology employed in this work is based around the 3-dimensionnal geometry of each molecules.

2.1 Molecular geometry pipeline

One of the most extensive database for chemical species is PubChem (Kim et al., 2025). Practically all of the molecular species present in the database have 3D conformers available to download. The files associated to those conformers, `.xyz` files, consits in the space position of each atom in 3D space. The code employed in this work make a request at PubChem server with the name of the searched molecule and fetch for the 3D structure as for other properties such as the formula. The 3D conformer catalogue of Pubchem was, as their documentation suggests[2], calculated using advanced quantum computational methods of atomic-force fields repultation, with the assurance of a great accuracy between experimental, theoritical and modeled 3D conformers (see Bolton, Kim, and Bryant, 2011). In this case, atomic coordinates should already be optimized and no further optimization is needed (see appendix **??**).

One problem emerge regarding molecules with no 3D conformer available. While they represent a very small fraction of this database, and appears as *niche* astrochemically speaking at first glance, they do represents molecules with a certain degre of interest. For instance, Germane GeH_4 could be present in the atmosphere of giant planets (Corice and Fox, 1972; Moses et al., 2020) as well as arsenic bearing molecules (Bezard et al., 1989; Roederer and Lawler, 2012). First calculations were initally made using 2D structure (*e.g* with x and y positions and z as vector fullfilled with zeroes), but it was quickly proven to be a wrong approach. While it could be possible to get 3D conformers from other sources (like Jmol ;Hanson, 2010), it is easier for automated workflow to keep the same source of data.




Figure 1: Overview of the molecular geometry pipeline for molecules without 3D conformer available. Germane GeH_4 is used as an example.

Thankfully, it is possible to obtain 3D geometry starting from PubChem 2D structures. For that, we used the SMILES id of each molecule. Using the `strainedSMILES2xyz` python package (Demuth, Schnizer, and Svatunek, 2026), we can retrieve a coherent 3D structure from the SMILES structure. While this packages is orignially aimed at strained ring system in complex molecules, it was proven to be very effective at converging to accurate 3D geometry (see the discussion section of Demuth, Schnizer, and Svatunek, 2026). It uses tools from the `RDKit` python library

[1]https://lh67000.github.io/casei

[2]https://pubchem.ncbi.nlm.nih.gov/docs/pubchem3d




Figure 2: Overview of the general workflow and methodology employed in this work.

(Landrum et al., 2025) at the start. Namely, the *Experimental-torsion-knowledge distance geometry* (ETKDG) method proposed by Riniker and Landrum (2015) which relies on atomic distances in order to find equilibrum in force field from each atoms. While `strainedSMILES2xyz` uses ORCA quantum model (Neese, 2012) in order to optimize obtained geometry, we choosed to rely on open-source tools to do so. As such, we used the `geometric` optimization package by Wang and Song (2016) for several cycles, in order to get a geometry that takes into account atomic translations and rotations. This optimization is done during PySCF calculations using the Hartree-Fock (HF) method, as described in subsection 2.3. Figure 1 recapitulates the overall workflow employed for those molecules. The molecules that use this pipeline are indicated in grey in table 1. However, an exception was made for sulfur hexafluoride SF_6, as it was proven to be very difficult to converge to a stable geometry that was coherent. As such, and regarding its importance for astrochemistry, the 3D conformer of SF_6 was downloaded from Jmol and used for calculations in that specific case.

2.2 The BEB method for TICS calculation

The *Binary Encounter Bethe* (BEB) method was proposed by Kim and Rudd (1994) in order to provide a simplier model than the *Binary Encounter Dipole* (BED) for ionisation cross-sections calculation. The BEB method estimates the cross-section of total ionisation using the electron energy of the Molecular Orbital (MO) and the kinetic energy of the electron whithin. In application, the BEB method can expressed as, for a distinct MO i:

$$\sigma_i(T) = \frac{S}{t+u+1}\left[\frac{\ln t}{2}\left(1-\frac{1}{t^2}\right)+1-\frac{1}{t}-\frac{\ln t}{t+1}\right] \quad (1)$$

with $t = \frac{T}{B}$, $u = \frac{U}{B}$, where T the ionisation particle energy and B and U are the aforementioned MO electrons and kinetic energy, and S is expressed as:

$$S = 4\pi a_0^2 N_i \left(\frac{R}{B}\right)^2 \quad (2)$$

with a_0 the Bohr radius (0.5292 $\mathring{A}$), R the Rydberg constant (13.6 eV) and N_i the number of electrons in the MO. For neutral stable species, N is always equal to 2. This, however, gives the cross section for only one MO. To get the total ionisation cross section, we need to sum all the $\sigma_i(T)$ of each MO, which gives us:

$$\sigma_{TICS}(T) = \sum_{i=1}^{n} \sigma_i(T) \quad (3)$$

Concerning T, this parameter is kept at relatively low energy. We considered 10^3 values in a logscale array with $T \in [10^1; 10^5]$ eV.

2.3 Quantum chemistry parameters calculations

As such, BEB calculations requires the MO electrons energies (B) as well as their kinetic energies (U) . While NIST database propose this parameter for several molecule[3], this is limited in terms of selection. One way is to calculate these parameter. To do that, PySCF quantum chemistry model (Sun et al., 2017) was employed. Using the downloaded geometries, we can input them into the model and obtain the required paramaters. This model is based on the Self Consistent Field (SCF) method of quantum chemistry for the HF approach. The basis-set family used in this work is the Karlsruhe `def2-TZVPPD` basis-set – valence triple-zeta with sets of polarization and diffuse functions (Weigend and Ahlrichs, 2005; Rappoport and Furche, 2010). Diffusion function are rather useful to describe electron affinity (Morrison, 2025). Although a quadruple-zeta (QVZ) basis set could have been used, we used a triple-zeta (TVZ) basis set in this work for computational reasons.

[3] https://physics.nist.gov/PhysRefData/Ionization/molTable.html

Table 1: List of molecules present in the database. Superscript number indicates the number of isomers. Grey molecules indicates that no 3D conformer was available on PubChem and that the geometry was generated from SMILES structure.

Number of atoms													
1	2	3	4	5	6	7	8	9	10	11	12	13	≥ 14
	AlCl	CHN^2	AsH_3	C_2H_2O	C_2Cl_4	$C_2H_3NO^2$	$C_2H_4N_2$	$C_2H_5NO^2$	$C_2H_5NO_2$	C_2H_7NO	$C_3H_6O_3$	$C_3H_8O_2^2$	$C_{17}H_9N^3$
	AlF	CHP	C_2H_2	C_2H_2S	$C_2H_2O_2$	$C_2H_4O^3$	$C_2H_4O_2^4$	$C_2H_6O^2$	$C_2H_6O_2^2$	$C_3H_6O_2^2$	C_3H_7NO	$C_3H_9N^3$	$C_{16}H_{10}$
	Br_2	CO_2	$C_2N_2^2$	C_2HNO	C_2H_3Cl	$C_2H_4S^2$	$C_2H_5N^2$	$C_2H_6S^2$	C_2H_6OS	C_3H_7N	$C_3H_8O^3$	C_3H_9P	$C_{14}H_{10}$
	BrH	COS	CH_2O	C_3HN^2	$C_2H_3N^3$	$C_3H_2N_2$	C_2H_6	$C_3H_3N_3$	$C_2H_7N^2$	C_3H_8	$C_4H_6O_2^2$	$C_4H_8O^2$	$C_{11}H_{12}N_2O_2$
	C_2	CS_2	CH_2S	CCl_4	C_2H_4	C_3H_3N	$C_3H_6NO^3$	$C_3H_4N_2^2$	C_3H_5NO	C_4H_6O	$C_4H_7N^3$	C_4H_8S	$C_{11}H_7N^2$
	CO	ClNO	$CHNO^3$	CF_4	C_2HCl_3	$C_3H_4^3$	C_3H_3NS	C_3H_5N	$C_3H_6O^4$	C_5H_5N	$C_4H_8^5$	$C_5H_8^5$	$C_{10}H_8$
	CS	FNO	$CHNS^3$	CH_2Cl_2	$C_3H_2O^4$	C_3N_4	$C_3H_4O^2$	$C_3H_6^2$	$C_4H_4N_2^2$	C_5H_6	$C_6H_6^3$	C_7H_5N	C_9H_8
	Cl_2	H_2O	H_2O_2	CH_2F_2	C_4H_2	C_5HN	$C_4H_3N^3$	C_4H_4S	$C_4H_5N^2$		C_6N_6		$C_8H_{10}^3$
	ClH	H_2S	H_2S_2	$CH_2N_2^2$	C_4N_2	CH_4Si_2	C_4H_4	$C_5H_4^2$	$C_4H_6^4$				C_8H_8
	F_2	H_2Se	H_3N	CH_2O_2	CH_2O_3	CH_5N	C_6H_2						$C_7H_8^2$
	FH	HNO	H_3P	CH_2S_2	CH_3NO	N_2O_5	C_6N_2						C_6H_{10}
	H_2	N_2O	HNO_2	CH_3Br	CH_3NS		CH_3ClO_2S						$C_6H_{12}^9$
	N_2	O_2S	HNOS	CH_3Cl	CH_3NSi		CH_4N_2O						$C_6H_{14}^4$
	OSi	O_3	O_3S	CH_3F	CH_4O		H_6Si_2						C_6N_8
				CH_3I	CH_4S								$C_5H_{10}^3$
				CH_3P	H_4N_2								C_5H_{12}
				CH_4	N_2O_4								C_5H_8O
				$CHCl_3$									$C_4H_{10}^2$
				CHF_3									$C_4H_{10}O^4$
				GeH_4									$C_4H_{10}S$
				H_3NO									$C_4H_8O_2^3$
				H_4Si									$C_3H_8O_3$
													C_3N_{12}

This type of basis, in particular the `def2-TZVPP` basis, is proven to be reliable for HF modeling for a majority of the periodic table, from H up to Rn (Hellweg and Rappoport, 2015; Zheng, Xu, and Truhlar, 2011). Several BEB-related papers use either the cross-correlated Dunning Dunning (1989) `cc-` base (Gaches et al., 2024; Kumar, 2014) or Pope (Krishnan et al., 1980) gaussian `6-311G` basis (Ali et al., 1997; Hwang, Kim, and Rudd, 1996; Scott and Irikura, 2005). At our knowledge, this is the first use of a `def2-` basis for TICS calculations. Only Restricted Hartree-Fock (RHF) calculations were performed, as all molecules are neutral and stable and thus have closed shell.

Regarding the energies output of the model, we assume that B is the absolute value of the lowest electron binding energy of the orbital, *e.g* the first ionisation potential (IP) which satisfies the Koopmans theorem (Koopmans, 1934) $B = |-\epsilon_i|$ with ϵ_i the ionisation energy of an orbital i. Whilst this is a good approach that is easy to implement and do not require any calculations, this does involve some simplifications as the orbitals are considered as "frozen", *e.g* that the other electrons do not react to the removal of ionized electron and that the post-ionization orbitals cannot be relaxed. This leads to overestimated first ionization potential for each orbitals (Colonna et al., 2019; Wang, Batool, and Wang, 2026; Jeseněk, Luque, and Lehtinen, 2025). However, alternative approach such as outervalence Green's function (OVGF, von Niessen, Schirmer, and Cederbaum, 1984), coupled cluster (CC, generally single-double-triple CCSDT; Čížek, 1966) require higher computational effiort. While some TICS-related papers use those approach (Scott and Irikura, 2005; Gaches et al., 2024), the BEB method was historically built around HF method and B, as well as IP, were either experimental values or theoritical values. Furthermore, first ionization potential approach is widely used in other BEB-related papers (Ali et al., 1997; Hwang, Kim, and Rudd, 1996; Graves, Cooper, and Tennyson, 2021) and do provide accurate results. One way to overcome the sturdyness of HF results is to use post-HF methods. We already discussed of complex method such as CCSSD(T) before, but such implementation require large amount of computational power. A less demanding method is to use Density Functionnal Theory (DFT) to overcome the assumptions. We can use the ΔDFT method to calculate the first ionisation potential of an orbital, as, for a given orbital i of an atom N (Wang and Vasilyev, 2025):

$$\Delta DFT = E_{i,N^+} - E_{i,N} = IP \tag{4}$$

with E_{i,N^+} the energy of a OM of the cation of the molecule and $E_{i,N}$ the MO energy of the neutral. Calculation are done using the Coulomb-Attenuating Method (CAM) with the B3LYP functional (Yanai, Tew, and Handy, 2004). The basis-set used is the same as for HF calculations, *i.e* `def2-TZVPPD`. See table 3 in appendix for comparison of our calculated IP with IP obtained though other means. Regarding kinetic energy U, this not a direct output of the model and it needs to be derived. As such, U is calculated as:

$$U_i = \mathbf{Tr}\, \mathbf{C}_i^{\mathbf{T}} \mathbf{T} \mathbf{C}_i \tag{5}$$

with $\mathbf{C}_i$ the MO coefficient matrix of the orbital i in the Linear Combinaison of Atomic Orbital (LCAO)[4] formalism and $\mathbf{T}$ the kinetic energy matrix that is composed by elements as

$$T_{xy} = \langle \chi_x | -\frac{1}{2}\nabla^2 | \chi_y \rangle \tag{6}$$

The BEB method, by itself, has limitations for heavy elements, with quantum number $n > 2$ or $Z > 10$ (Graves, Cooper, and Tennyson, 2021; Huo and Kim, 2000; Ali and Kim, 2008; Scott and Irikura, 2005). The discrepancy arises because the BEB method uses, as shown in Eq. 1, the kinetic energy of each molecular orbital as a variable. This is problematic for heavier elements, as inner electronic layers of this type have very high kinetic energies. However, ionisation should occur when the electron is far from the

[4] For a given guess wave function $\phi_j(\mathbf{r})$: $\phi_j = C_{\mu j}\chi_\mu(\mathbf{r})$ with $C_{\mu j}$ the expansion coefficient and χ_μ the atom centered Gaussian basis function.

centre of the shell, i.e. when it is slower and thus has a lower kinetic energy. An overestimation of the kinetic energy leads to an underestimation of σ_{TICS}. The solution proposed by Huo and Kim (2000), and employed in several BEB-related studies on heavy elements (Graves, Cooper, and Tennyson, 2021; Dolg and Cao, 2012; Scott and Irikura, 2005), is to use the effective core potential (ECP) method. An ECP limits the interaction of valence electrons with the core shell by replacing the latter with analytical functions that represents the interaction between the two. These functions appear as effective potential, while the valence electrons are described by pseudo-valence orbitals that do not penetrate the inner region of the nucleus. In a nut*shell*, the ECP enables the removal of high kinetic energies of core electrons from the BEB sum, as core electrons do not intervene in electronic bonds, which would otherwise artificially minimalize σ_{TICS}. The ECP method can be used in PySCF and, in order to include as many molecules as possible in the database, was implemented in this work. This, however, requires a different basis than the all-electrons Karlsruhe `def2-`, because its related ECP basis only start at Rb ($Z \geq 37$) and thus don't work for lighter elements, *e.g* $3p$ and $4p$ elements. The only basis that has an ECP basis that supports these that is inherently available in PySCF and in the Basis Set Exchange database[5] (Pritchard et al., 2019) is the `crenbl` basis set (Hurley et al., 1986; Fernandez Pacios and Christiansen, 1985). More modern basis are available on the open-source Pseudopotentiallibrary[6], as it provides Dunning derived basis combo for both $3p$ and $4p$ elements. For Al, Si, P, S, Cl and Ar bearing molecules, we used the `aug-cc-pVTZ` orbital basis along with its associated ECP basis proposed by (Bennett et al., 2018), whereas for Ge, As, Se and Br bearing molecules we used the same orbital basis and the ECP basis proposed by (Wang et al., 2019). The impact of ECP on TICS calculation is discussed in subsection 3.3. Overall, the whole methodology employed in this work is summarized on figure 2

3 Results

3.1 Overview

260 total ionisation cross section of unique molecules have been calculated, and up to 9 isomer for some formula such as C_6H_{12}. The complete list is visible on table 1. On the database website, the user can find the TICS table of each molecule, along with a table with the calculated B and U values for each molecular orbital following the chosen basis.

3.2 HF method agains post-HF methods

3.3 ECP impact on TICS calculation

On table 2 we compared the mean Δ of TICS between ECP and non-ECP calculations for $3p$ and $4p$ elements bearing molecules. We compared the results of the `cc-ECP` basis aswell as with the `crenbl` basis with the `def2-` basis. The mean Δ percentage is calculated as:

$$\overline{\Delta\sigma}(\%) = \left\langle \frac{\sigma_{\text{def2-}} - \sigma_{\text{ECP}}}{\sigma_{\text{ECP}}} \times 100 \right\rangle \tag{7}$$

with σ_{ECP} the TICS calculated with either `cc-ECP` or `crenbl`. Figure 3 shows this comparison for several molecules. As expected, and as proposed un Huo and Kim (2000) as well as in other BEB-related studies on heavy elements (Graves, Cooper, and Tennyson, 2021; Dolg and Cao, 2012; Scott and Irikura, 2005), the use of ECP basis leads to a higher TICS value. As seen in the table, we

Table 2: Mean Δ of TICS between ECP and non-ECP calculations for $3p$ and $4p$ elements bearing molecules.

	$\overline{\Delta\sigma}$ (%)	
Molecule	cc-ECP	CRENBL
H_3P	13.996	14.568
H_4Si	6.505	7.311
H_6Si_2	9.242	9.283
Geh_4	27.548	32.212
H_2Se	27.384	31.159
H_2S	18.474	16.958
Ch_3Br	20.059	22.418
Brh	24.582	27.666
Br_2	26.276	26.496
Cs	11.996	0.948
Cl_2O	18.123	12.592
Osi	5.703	6.007
Csi	7.079	7.372
Ch_4Si_2	5.093	7.995
Ch_3Nsi	2.749	3.753
$C_4H_{10}S$	9.545	9.718

3.4 Impact of isomery on TICS calculation

4 Discussion

4.1 Comparison with literature

In order to validate our results, we compared our calculated TICS with several experimental and theoretical data from the literature.

Hydrocarbons

The Des soucis avec les hydrocarbures saturés ? Le propene par ex

As pointed out by Scott and Irikura (2005), Koopmans (i.e., Hartree-Fock) binding energies are too approximate for this purpose. par exemple utilise le outervalence Green's function (OVGF) Wang and Song (2016) utilisent aussi et disent que Koopmans c la merde

Nitrogen bearing molecules

Oxygen bearing molecules

Sulfur and phosphorus bearing molecules

Heavy elements bearing molecules

First and foremost, experimental B values tend to be smaller than theoritical/computed B values (Ali et al., 1997; Hwang, Kim, and Rudd, 1996). As such, the cross section obtained through experimental B values tend to be higher,≈10-15% at the cross section peak. This was observed several time in the plots of figures 1-3, where theoritical curves were generally smaller than those from experimental papers. This is not systemic.

[5] https://basissetexchange.org/
[6] https://pseudopotentiallibrary.org/




Figure 3: Comparison of TICS results for several molecule with and without ECP enabled

4.2 Limits

Whereas this paper serves as a database presentation, some results do harbour scientific interest, as some molecule haven't got a dedicated TICS paper yet. blalbalbalb
As shown in several studies, isomery play a minor role regarding σ_{TICS}. dshfdv

Ici évoque le SF6, qui a un petit intérêt astro mais que j'arrive pas a calcluer la géométrie

4.3 Applications to mass spectrometry

5 Conclusion

Pouvoir utiliser un spectre complet de section efficace d'ionisation comme ceux que j'ai calculé pour calculer la section effiace d'ionisation partielle ? Pouvoir brancher cela a des autres modeles (comme celui de Gautier 2020) ? Faire du calcul de flux d'ionisation, jpeux essayer de lier ca a triton

Multiple outlook could be imaginated:

- Adapt the pipeline for radicals (*i.e* molecules with spin $\neq 0$) and ions; This would require to do calculation using the UHF method of SCF;
- Refine and optimize the geometry of molecules -if possible. This could be useful for molecules taken from a 2D structure.
- Adapt and create a custom `def2-` ECP basis for $3d$ and $4p$ elements in order to homogenize the database as much as possible protocol-wise;

Acknowledgement

Appendix

Optimization of PubChem geometries

Energy of first ionisation IP

References

Ali, M A and Yong-Ki Kim (July 2008). "Ionization Cross Sections by Electron Impact on Halogen Atoms, Diatomic Halogen and Hydrogen Halide Molecules". In: *Journal of Physics B: Atomic, Molecular and Optical Physics* 41.14, p. 145202. ISSN: 0953-4075, 1361-6455. DOI: 10.1088/0953-4075/41/14/145202. (Visited on 10/28/2026).

Ali, M. A. et al. (June 1997). "Electron-Impact Total Ionization Cross Sections of Silicon and Germanium Hydrides". In: *The Journal of Chemical Physics* 106.23, pp. 9602–9608. ISSN: 0021-9606, 1089-7690. DOI: 10.1063/1.473842. (Visited on 10/28/2026).

Bennett, M. Chandler et al. (Sept. 2018). "A New Generation of Effective Core Potentials from Correlated Calculations: 2nd Row Elements". In: *The Journal of Chemical Physics* 149.10, p. 104108. ISSN: 0021-9606. DOI: 10.1063/1.5038135. (Visited on 10/29/2026).

Bezard, B. et al. (Nov. 1989). "Detection of Arsine in Saturn". In: *The Astrophysical Journal* 346, p. 509. ISSN: 0004-637X. DOI: 10.1086/168032. (Visited on 11/05/2026).

Bolton, Evan E, Sunghwan Kim, and Stephen H Bryant (Dec. 2011). "PubChem3D: Conformer Generation". In: *Journal of Cheminformatics* 3.1, p. 4. ISSN: 1758-2946. DOI: 10.1186/1758-2946-3-4. (Visited on 11/05/2026).

Burgasser, Adam J. et al. (Nov. 2025). "Observation of Undepleted Phosphine in the Atmosphere of a Low-Temperature Brown Dwarf". In: *Science* 390.6774, pp. 697–701. ISSN: 0036-8075, 1095-9203. DOI: 10.1126/science.adu0401. arXiv: 2510.03916 [astro-ph.SR]. (Visited on 10/28/2026).




Figure 4: Calculted TICS for several hydrocarbons. In those plots, σ_{TICS} as been cropped at 10^{-8}. Scatter plot mean experimental data and line plot mean theoritical data. Those data come from Rapp and Englander-Golden (1965), Jiao, DeJoseph, and Garscadden (2007), Chatham et al. (1984), and Ali et al. (1997). The references marked with an * where obtained through lxcat.com database (Dutton, 2026; Carbone et al., 2021; Pitchford et al., 2017; Pancheshnyi et al., 2012). Those references include

.




Figure 5: Calculted TICS for several organic and non-organic molecules. In those plots, σ_{TICS} as been cropped at 10^{-8}. Scatter plot mean experimental data and line plot mean theoritical data. Those data come from Rapp and Englander-Golden (1965), Jiao, DeJoseph, and Garscadden (2007), Chatham et al. (1984), Ali et al. (1997), and Kumar (2014). The references marked with an * where obtained through lxcat.com database. Those references include

Table 3: Comparison of the energy of the ionization potential between this work (where $B = -\epsilon$) and Gaches et al. (2024) CCSD(T) and CAM-B3LYP calculations.

Molecule	This work	IP (CCSD(T))	IP (CAM)
CO	14,84	14,13	14,24
H_2	16,23	16,45	
H_2S	10,45	10,52	10,43
H_2O	13,87	12,8	12,69
O_3	13,26	13,02	12,77
C_2H_2	13,48	11,54	11,28
C_2N_2	13,48	14,74	13,25
C_3H_2O	11,59		10,59
CH_2O	12,15	11,04	10,88
CH_2S	9,61	9,5	9,3
NH_3	11,73	11,04	10,22
CH_4	14,80		12,72
C_2H_4	10,25		10,4
N_2H_4	10,46		6,82
C_4N_2	12,08		11,67
$c-H_2C3O$	10,54		9,54
CH_3NS	9,04		9,45
C_4H_2	10,14		9,96
C_2H_6	13,19		11,59
C_6N_2	11,25		10,69
C_4H_4	9,39		9,32
C_5H_4	9,64		10,03
C_3H_8	12,74		10,99
C_4H_{10}	12,28		10,89
$c-C_6H_6$	9,16		9,17
$C_{14}H_{10}$	6,74		7,21
C_5H_{12}	11,95		10,76
C_6H_{14}	11,69		10,77

Carbone, Emile et al. (Mar. 2021). "Data Needs for Modeling Low-Temperature Non-Equilibrium Plasmas: The LXCat Project, History, Perspectives and a Tutorial". In: *Atoms* 9.1, p. 16. ISSN: 2218-2004. DOI: 10.3390/atoms9010016. (Visited on 11/05/2026).

Chatham, H. et al. (Aug. 1984). "Total and Partial Electron Collisional Ionization Cross Sections for CH4, C2H6, SiH4, and Si2H6". In: *The Journal of Chemical Physics* 81.4, pp. 1770–1777. ISSN: 0021-9606. DOI: 10.1063/1.447848. (Visited on 10/22/2026).

Čížek, Jiří (Dec. 1966). "On the Correlation Problem in Atomic and Molecular Systems. Calculation of Wavefunction Components in Ursell-Type Expansion Using Quantum-Field Theoretical Methods". In: *The Journal of Chemical Physics* 45.11, pp. 4256–4266. ISSN: 0021-9606. DOI: 10.1063/1.1727484. (Visited on 11/09/2026).

Colonna, Nicola et al. (Mar. 2019). "Koopmans-Compliant Functionals and Potentials and Their Application to the GW100 Test Set". In: *Journal of Chemical Theory and Computation* 15.3, pp. 1905–1914. ISSN: 1549-9618, 1549-9626. DOI: 10.1021/acs.jctc.8b00976. (Visited on 11/09/2026).

Corice, Raymond J. and Kenneth Fox (Apr. 1972). "The Hypothetical Chemical and Spectroscopic Activity of Germane in the Atmosphere of Jupiter". In: *Icarus* 16.2, pp. 388–391. ISSN: 0019-1035. DOI: 10.1016/0019-1035(72)90082-6. (Visited on 11/05/2026).

Demuth, Tori, Julian Schnizer, and Dennis Svatunek (May 2026). "strainedSMILES2xyz: A Workflow for Reliable 3D Structures of Strained Molecules from SMILES". In: *Journal of Cheminformatics* 18.1, p. 80. ISSN: 1758-2946. DOI: 10.1186/s13321-026-01219-0. (Visited on 10/30/2026).

Dolg, Michael and Xiaoyan Cao (Jan. 2012). "Relativistic Pseudopotentials: Their Development and Scope of Applications". In: *Chemical Reviews* 112.1, pp. 403–480. ISSN: 0009-2665. DOI: 10.1021/cr2001383. (Visited on 10/28/2026).

Dunning, Thom H. (Jan. 1989). "Gaussian Basis Sets for Use in Correlated Molecular Calculations. I. The Atoms Boron through Neon and Hydrogen". In: *The Journal of Chemical Physics* 90.2, pp. 1007–1023. ISSN: 0021-9606, 1089-7690. DOI: 10.1063/1.456153. (Visited on 10/28/2026).

Dutton (2026). *LXCAT Database*. www.lxcat.net.

Fernandez Pacios, Luis and P. A. Christiansen (Mar. 1985). "Ab Initio Relativistic Effective Potentials with Spin-orbit Operators. I. Li through Ar". In: *The Journal of Chemical Physics* 82.6, pp. 2664–2671. ISSN: 0021-9606. DOI: 10.1063/1.448263. (Visited on 10/28/2026).

Gaches, Brandt A. L. et al. (Apr. 2024). "The Astrochemistry Low-Energy Electron Cross-Section (ALeCS) Database - I. Semi-Empirical Electron-Impact Ionization Cross-Section Calculations and Ionization Rates". In: *Astronomy & Astrophysics* 684, A41. ISSN: 0004-6361, 1432-0746. DOI: 10.1051/0004-6361/202348293. (Visited on 10/21/2026).

Graves, Vincent, Bridgette Cooper, and Jonathan Tennyson (Mar. 2021). "The Efficient Calculation of Electron Impact Ionization Cross Sections with Effective Core Potentials". In: *The Journal of Chemical Physics* 154.11, p. 114104. ISSN: 0021-9606, 1089-7690. DOI: 10.1063/5.0039465. (Visited on 10/28/2026).

Hanson, Robert M. (Oct. 2010). "*Jmol* – a Paradigm Shift in Crystallographic Visualization". In: *Journal of Applied Crystallography* 43.5, pp. 1250–1260. ISSN: 0021-8898. DOI: 10.1107/S0021889810030256. (Visited on 11/05/2026).

Hellweg, Arnim and Dmitrij Rappoport (Jan. 2015). "Development of New Auxiliary Basis Functions of the Karlsruhe Segmented Contracted Basis Sets Including Diffuse Basis Functions (Def2-SVPD, Def2-TZVPPD, and Def2-QVPPD) for RI-MP2 and RI-CC Calculations". In: *Physical Chemistry Chemical Physics* 17.2, pp. 1010–1017. ISSN: 1463-9076. DOI: 10.1039/c4cp04286g. (Visited on 10/28/2026).

Huo, Winifred M and Yong-Ki Kim (Mar. 2000). "Use of Relativistic Effective Core Potentials in the Calculation of Total Electron-Impact Ionization Cross-Sections". In: *Chemical Physics Letters* 319.5-6, pp. 576–586. ISSN: 00092614. DOI: 10.1016/S0009-2614(00)00150-0. (Visited on 10/28/2026).

Hurley, M. M. et al. (June 1986). "Ab Initio Relativistic Effective Potentials with Spin-orbit Operators. II. K through Kr". In: *The Journal of Chemical Physics* 84.12, pp. 6840–6853. ISSN: 0021-9606. DOI: 10.1063/1.450689. (Visited on 10/29/2026).

Hwang, W., Y.-K. Kim, and M. E. Rudd (Feb. 1996). "New Model for Electron-Impact Ionization Cross Sections of Molecules". In: *The Journal of Chemical Physics* 104.8, pp. 2956–2966. ISSN: 0021-9606, 1089-7690. DOI: 10.1063/1.471116. (Visited on 10/28/2026).

Jeseněk, Anthony, Alejandro Luque, and Nikolai Lehtinen (Aug. 2025). "Partial Ionisation Cross Sections for the Binary-Encounter Bethe Model". In: *Plasma Sources Science and Technology* 34.8, p. 085015. ISSN: 0963-0252. DOI: 10.1088/1361-6595/adfc09. (Visited on 11/10/2026).

Jiao, C Q, C A DeJoseph, and A Garscadden (Jan. 2007). "Electron Impact Ionization and Ion Reactions in N-Butane". In: *Journal of Physics D: Applied Physics* 40.2, p. 409. ISSN: 0022-3727. DOI: 10.1088/0022-3727/40/2/018. (Visited on 10/22/2026).

Kim, Sunghwan et al. (Jan. 2025). "PubChem 2025 Update". In: *Nucleic Acids Research* 53.D1, pp. D1516–D1525. ISSN: 1362-4962. DOI: 10.1093/nar/gkae1059. (Visited on 11/05/2026).

Kim, Yong-Ki and M. Eugene Rudd (Nov. 1994). "Binary-Encounter-Dipole Model for Electron-Impact Ionization". In: *Physical Review A* 50.5, pp. 3954–3967. ISSN: 1050-2947, 1094-1622. DOI: 10.1103/PhysRevA.50.3954. (Visited on 10/28/2026).

Koopmans, T. (Jan. 1934). "Über Die Zuordnung von Wellenfunktionen Und Eigenwerten Zu Den Einzelnen Elektronen Eines Atoms". In: *Physica* 1, pp. 104–113. DOI: 10.1016/S0031-8914(34)90011-2. (Visited on 11/09/2026).

Krishnan, R. et al. (Jan. 1980). "Self-Consistent Molecular Orbital Methods. XX. A Basis Set for Correlated Wave Functions". In: *The Journal of Chemical Physics* 72.1, pp. 650–654. ISSN: 0021-9606, 1089-7690. DOI: 10.1063/1.438955. (Visited on 10/28/2026).

Kumar, Rajeev (Aug. 2014). "Electron Ionization Cross Sections for the PH3 Molecule". In: *Chemical Physics Letters* 609, pp. 108–112. ISSN: 00092614. DOI: 10.1016/j.cplett.2014.06.034. (Visited on 10/22/2026).

Landrum, Greg et al. (June 2025). "Rdkit/Rdkit: 2025_03_4 (Q1 2025) Release". In: DOI: 10.5281/zenodo.15773589. (Visited on 11/06/2026).

Larson, H. P. et al. (Aug. 1980). "The Middle-Infrared Spectrum of Saturn - Evidence for Phosphine and Upper Limits to Other Trace Atmospheric Constituents". In: *The Astrophysical Journal* 240, pp. 327–337. ISSN: 0004-637X. DOI: 10.1086/158236. (Visited on 10/28/2026).

Morrison, Robert C. (Oct. 2025). "The Limits of Accuracy of the Extended Koopmans' Theorem for Electron Affinities". In: *The Journal of Chemical Physics* 163.14, p. 144102. ISSN: 0021-9606, 1089-7690. DOI: 10.1063/5.0292192. (Visited on 11/09/2026).

Moses, J. I. et al. (Dec. 2020). "Atmospheric Chemistry on Uranus and Neptune". In: *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 378.2187, p. 20190477. ISSN: 1364-503X, 1471-2962. DOI: 10.1098/rsta.2019.0477. (Visited on 11/02/2026).

Neese, Frank (Jan. 2012). "The ORCA Program System". In: *WIREs Computational Molecular Science* 2.1, pp. 73–78. ISSN: 1759-0876, 1759-0884. DOI: 10.1002/wcms.81. (Visited on 11/06/2026).

Pancheshnyi, Sergey et al. (2012). "The LXCat Project : Electron Scattering Cross Sections and Swarm Parameters for Low Temperature Plasma Modeling". In: *Chemical Physics* 398, pp. 148–153. DOI: 10.1016/j.chemphys.2011.04.020. (Visited on 11/05/2026).

Pitchford, Leanne C. et al. (Jan. 2017). "LXCat: An Open-Access, Web-Based Platform for Data Needed for Modeling Low Temperature Plasmas". In: *Plasma Processes and Polymers* 14.1-2. ISSN: 1612-8850. DOI: 10.1002/ppap.201600098. (Visited on 11/05/2026).

Pritchard, Benjamin P. et al. (Nov. 2019). "New Basis Set Exchange: An Open, Up-to-Date Resource for the Molecular Sciences Community". In: *Journal of Chemical Information and Modeling* 59.11, pp. 4814–4820. ISSN: 1549-9596. DOI: 10.1021/acs.jcim.9b00725. (Visited on 10/22/2026).

Rapp, Donald and Paula Englander-Golden (Sept. 1965). "Total Cross Sections for Ionization and Attachment in Gases by Electron Impact. I. Positive Ionization". In: *The Journal of Chemical Physics* 43.5, pp. 1464–1479. ISSN: 0021-9606. DOI: 10.1063/1.1696957. (Visited on 10/27/2026).

Rappoport, Dmitrij and Filipp Furche (Oct. 2010). "Property-Optimized Gaussian Basis Sets for Molecular Response Calculations". In: *The Journal of Chemical Physics* 133.13, p. 134105. ISSN: 0021-9606, 1089-7690. DOI: 10.1063/1.3484283. (Visited on 11/06/2026).

Ridgway, S. T., L. Wallace, and G. R. Smith (Aug. 1976). "The 800 - 1200 Inverse Centimeter Absorption Spectrum of Jupiter." In: *The Astrophysical Journal* 207, pp. 1002–1006. ISSN: 0004-637X. DOI: 10.1086/154570. (Visited on 10/28/2026).

Riniker, Sereina and Gregory A. Landrum (Dec. 2015). "Better Informed Distance Geometry: Using What We Know To Improve Conformation Generation". In: *Journal of Chemical Information and Modeling* 55.12, pp. 2562–2574. ISSN: 1549-9596, 1549-960X. DOI: 10.1021/acs.jcim.5b00654. (Visited on 11/06/2026).

Roederer, Ian U. and James E. Lawler (May 2012). "DETECTION OF ELEMENTS AT ALL THREE *r* -PROCESS PEAKS IN THE METAL-POOR STAR HD 160617". In: *The Astrophysical Journal* 750.1, p. 76. ISSN: 0004-637X, 1538-4357. DOI: 10.1088/0004-637X/750/1/76. (Visited on 11/05/2026).

Scott, Gregory E. and Karl K. Irikura (Nov. 2005). "Electron-Impact Ionization Cross Sections of Molecules Containing Heavy Elements (Z > 10)". In: *Journal of Chemical Theory and Computation* 1.6, pp. 1153–1161. ISSN: 1549-9618. DOI: 10.1021/ct050077j.

Sun, Qiming et al. (Mar. 2017). *The Python-Based Simulations of Chemistry Framework (PySCF)*. DOI: 10.48550/arXiv.1701.08223. arXiv: 1701.08223 [physics.chem-ph]. (Visited on 10/28/2026).

von Niessen, W., J. Schirmer, and L. S. Cederbaum (Apr. 1984). "Computational Methods for the One-Particle Green's Function". In: *Computer Physics Reports* 1.2, pp. 57–125. ISSN: 0167-7977. DOI: 10.1016/0167-7977(84)90002-9. (Visited on 11/09/2026).

Wang, Feng, Fozia Batool, and James Wang (May 2026). *Ioniza on Poten al of JP-10 Fuel: A First-Principles Study Addressing a 50-Year Gap in Experimental Data*. DOI: 10.26434/chemrxiv.15003977/v1. (Visited on 11/09/2026).

Wang, Feng and Vladislay Vasilyev (Aug. 2025). "Fe 3d Orbital Evolution in Ferrocene Ionization: Insights from ΔSCF, EOES, and Orbital Momentum Distribution". In: *Molecules* 30.17, p. 3541. ISSN: 1420-3049. DOI: 10.3390/molecules30173541.

Wang, Guangming et al. (Oct. 2019). "A New Generation of Effective Core Potentials from Correlated Calculations: 4s and 4p Main Group Elements and First Row Additions". In: *The Journal of Chemical Physics* 151.14, p. 144110. ISSN: 0021-9606. DOI: 10.1063/1.5121006. (Visited on 10/29/2026).

Wang, Lee-Ping and Chenchen Song (June 2016). "Geometry Optimization Made Simple with Translation and Rotation Coordinates". In: *The Journal of Chemical Physics* 144.21, p. 214108. ISSN: 0021-9606. DOI: 10.1063/1.4952956. (Visited on 10/30/2026).

Weigend, Florian and Reinhart Ahlrichs (2005). "Balanced Basis Sets of Split Valence, Triple Zeta Valence and Quadruple Zeta Valence Quality for H to Rn: Design and Assessment of Accuracy". In: *Physical Chemistry Chemical Physics* 7.18, p. 3297. ISSN: 1463-9076, 1463-9084. DOI: 10.1039/b508541a. (Visited on 10/28/2026).

Yanai, Takeshi, David P Tew, and Nicholas C Handy (July 2004). "A New Hybrid Exchange–Correlation Functional Using the Coulomb-Attenuating Method (CAM-B3LYP)". In: *Chemical Physics Letters* 393.1-3, pp. 51–57. ISSN: 00092614. DOI: 10.1016/j.cplett.2004.06.011. (Visited on 11/09/2026).

Zheng, Jingjing, Xuefei Xu, and Donald G. Truhlar (Feb. 2011). "Minimally Augmented Karlsruhe Basis Sets". In: *Theoretical Chemistry Accounts* 128.3, pp. 295–305. ISSN: 1432-881X, 1432-2234. DOI: 10.1007/s00214-010-0846-z. (Visited on 10/28/2026).