

Figure 1. Composition-dependent A–B score comparison

Points show supplied aggregate mean differences, not individual unit observations. Lower score is better: negative A–B favors A, positive A–B favors B. The record contains eight independent paired units, four in g1 and four in g2; technical repetitions were averaged within unit/method after conversion. g1 is -1.00 score and g2 is $+3.00$. The equal sampled mixture (0.5/0.5) is $+1.00$; the target mixture (0.8/0.2, fixed before measurement) is -0.20 . The horizontal interval on the target point is the supplied 95% within-stratum paired-unit bootstrap percentile interval $[-0.60, 0.20]$, with paired units resampled within stratum and target weights reapplied. No stratum or equal-mixture intervals were supplied, so their points have no error bars. The zero reference marks equality of scores, not a tested equivalence margin. No causal or equivalence interpretation and no significance test or symbols are asserted. The figure is redrawn from the authoritative version 2 result memo; numerical source data and bootstrap outputs are unavailable, so this is a display of reported results rather than independent replication.

Alt text: Four labeled points share an A–B score axis with a zero reference. g1 is at -1.00 , g2 at $+3.00$, equal sampled mixture at $+1.00$, and target mixture at -0.20 . The target's 95% bootstrap percentile interval extends from -0.60 to $+0.20$ and crosses zero. The two aggregate points lie on opposite sides of zero.