

Essential Summary

Generalized Padeé Approximation Method for Homoclinic Orbits of Strongly Nonlinear Autonomous Oscillators

Zhenbo Li, Jiashi Tang, Ping Cai

Chinese Journal of Theoretical and Applied Mechanics

Vol. 45, No. 3, 461–464 (2013) DOI: 10.6052/0459-1879-12-277

This is an author-prepared essential summary, not the original paper.
Please cite the published version.

1 Background and Motivation

The determination of homoclinic orbits is central to understanding global bifurcations, chaos, and solitary wave solutions in nonlinear dynamics. Existing Padeé-based methods share a limitation: approximants are restricted to polynomial numerators and denominators, reducing flexibility for exponential-decay homoclinic behavior.

This paper introduces the **generalized Padeé approximation (GPA) method**, extending numerator and denominator from polynomials to series of *any* function type. This unifies prior Padeé-type variants as special cases. A novel approximant using hyperbolic functions is constructed for solving homoclinic orbits of strongly nonlinear autonomous oscillators.

2 Core Innovation

2.1 The Generalized Padeé Definition

Let $\hat{H}_m = \{P : P(z) = \sum_{i=0}^m a_i g_i(z)^i\}$ and $G(m, n) = \{R = P/Q : P \in \hat{H}_m, Q \in \hat{H}_n \setminus \{0\}\}$. The GPA $[L/M]_f$ is $P_L/Q_M \in G(L, M)$ satisfying:

$$f(z) - \frac{P_L(z)}{Q_M(z)} = \mathcal{O}(z^{L+M+1}). \quad (1)$$

When $g_i(z) = z^i$, the GPA reduces to the classical Padeé approximant.

2.2 Homoclinic Orbit via Hyperbolic Functions

For $\ddot{x} + g(x) = 0$ with saddle $H(h, 0)$, the power series $x(t) = \sum_{i=0}^{\infty} a_i t^i$ is obtained by substitution. The initial amplitude a is from $\int_h^a g(x) dx = 0$. The key step: construct a GPA using $\text{sech}^i t$ that automatically satisfies the boundary condition $x(\pm\infty) = h$:

$$\text{GPA}[L/M] = \frac{\sum_{i=0}^L \alpha_i \text{sech}^i t}{1 + \sum_{i=1}^M \beta_i \text{sech}^i t}. \quad (2)$$

The $L + M + 1$ coefficients $\{\alpha_i, \beta_i\}$ are determined by Taylor-expanding at $t = 0$ and matching with the power series $\{a_i\}$ — a purely algebraic procedure.

3 Key Results

3.1 Example 1: Strongly Asymmetric Helmholtz–Duffing

$$\ddot{x} + c_1 x + c_2 x^2 = 0, \quad c_1 = -3, \quad c_2 = 100. \quad (3)$$

With $L = M = 2$, GPA[2/2] gives:

$$\text{GPA}[2/2] = \frac{0.005 \operatorname{sech} t + 0.07 \operatorname{sech}^2 t}{1 + 0.611111 \operatorname{sech} t + 0.055555 \operatorname{sech}^2 t}. \quad (4)$$

The phase portrait matches Runge–Kutta with excellent agreement despite $c_2 = 100 \gg 1$.




Figure 1: **Figure 1.** (Original Fig. 1) Homoclinic orbit for $c_1 = -3, c_2 = 100$. Solid: Runge–Kutta. Dotted: GPA[2/2].

3.2 Example 2: Cubic–Quadratic Helmholtz–Duffing

$$\ddot{x} + c_1 x + c_2 x^2 + c_3 x^3 = 0, \quad c_1 = -2, \quad c_2 = 5, \quad c_3 = 15. \quad (5)$$

Both left and right homoclinic orbits are obtained simultaneously with $L = M = 2$, showing the method handles asymmetric nonlinearities without modification.

3.3 Example 3: Duffing–Harmonic Oscillator

$$\ddot{x} + \frac{\lambda x + \mu x^3}{1 + \nu x^2} = 0, \quad \lambda = -1, \quad \mu = 1, \quad \nu = 1. \quad (6)$$

With $L = M = 3$, the GPA captures homoclinic orbits of this rational-restoring-force oscillator — a system whose exact solution is not expressible in elementary or classical elliptic functions.

4 Significance

1. **Theoretical unification:** The GPA definition unifies classical Padeé and all quasi-Padeé variants under a single framework.




Figure 2: **Figure 2.** (Original Fig. 2) Homoclinic orbits for $c_1 = -2, c_2 = 5, c_3 = 15$. Solid: Runge-Kutta. Dotted: GPA[2/2].




Figure 3: **Figure 3.** (Original Fig. 3) Homoclinic orbits for $\lambda = -1, \mu = 1, \nu = 1$. Solid: Runge-Kutta. Dotted: GPA[3/3].

2. **Computational efficiency:** $\operatorname{sech}^i t$ has simpler Taylor coefficients than exponentials, reducing matching cost at the same order.
3. **Broad applicability:** Handles polynomial $g(x)$ of arbitrary degree and rational $g(x)$ without modification.

5 Limitations

- Conservative systems only ($\varepsilon = 0$).
- Approximation order chosen empirically; no a priori error bound.
- Single-degree-of-freedom only.

6 Connection to the GHFP/MGHFP Series

This paper and its English version (Chin. Phys. B 2014) and the GHFP/MGHFP series approach nonlinear oscillators via complementary routes: GPA directly constructs homoclinic solutions through hyperbolic-function rational approximants; GHFP/MGHFP uses angular-domain perturbation theory for amplitude–parameter relationships. Both share the mathematical insight of rational function approximation.

Author: Zhenbo Li (lizhenbo@usc.edu.cn)
 University of South China, School of Mathematics and Physics
 Hunan Key Laboratory of Mathematical Modeling and Scientific Computing