

# Generalized Padé Approximation Methods for Strongly Nonlinear Oscillators: A Personal Retrospective

Five Papers, One Direction — From Homoclinic Orbits to Periodic Solutions

Zhenbo Li

School of Mathematics and Physics, University of South China, Hengyang 421001, China

Hunan Key Laboratory of Mathematical Modeling and Scientific Computing

lizhenbo@usc.edu.cn

## Abstract

This review traces the development of generalized Padé approximation (GPA) methods for strongly nonlinear oscillators through five papers published between 2013 and 2022. The intellectual arc spans from the initial generalization of the classical Padé definition (Li, 2013), through its extension to homoclinic/heteroclinic orbits (Chin. Phys. B, 2014), to the integration of GPA with the Lindstedt–Poincaré perturbation framework—the GPLP method (Nonlinear Dynamics, 2016). The framework was further extended to periodic solutions via cosine-type approximants (J. Vib. Shock, 2019) and finally refined for asymmetric potentials with combined Sech-Tanh approximants (J. Vib. Eng. Technol., 2022). The unifying theme is a simple yet powerful insight: by generalizing the Padé approximant from rational functions of polynomials to rational functions of *any* continuous function, one can tailor the approximant’s functional form to the geometric character of the solution being sought. This review summarizes the motivations, key methodological innovations, core results, limitations, and future directions of the GPA research program.

**Keywords:** Generalized Padé approximation, homoclinic orbits, heteroclinic orbits, periodic solutions, strongly nonlinear oscillators, perturbation methods, Lindstedt–Poincaré method.

## Contents

|          |                                                                                         |          |
|----------|-----------------------------------------------------------------------------------------|----------|
| <b>1</b> | <b>Introduction: Why Generalize Padé?</b>                                               | <b>3</b> |
| <b>2</b> | <b>Paper I (2013): The Generalized Padé Approximation — Definition and Foundation</b>   | <b>3</b> |
| 2.1      | Motivation . . . . .                                                                    | 3        |
| 2.2      | The GPA Definition . . . . .                                                            | 3        |
| 2.3      | Application: Homoclinic Orbits via Hyperbolic Basis . . . . .                           | 4        |
| 2.4      | Key Contributions of Paper I . . . . .                                                  | 5        |
| <b>3</b> | <b>Paper II (2014): Extending GPA to Heteroclinic Orbits and Irrational Oscillators</b> | <b>5</b> |
| 3.1      | Motivation . . . . .                                                                    | 5        |
| 3.2      | Methodological Extension . . . . .                                                      | 5        |
| 3.3      | Key Contributions of Paper II . . . . .                                                 | 5        |
| <b>4</b> | <b>Paper III (2016): The GPLP Method — Marrying GPA with Lindstedt–Poincaré</b>         | <b>5</b> |
| 4.1      | Motivation . . . . .                                                                    | 5        |
| 4.2      | The GPLP Procedure . . . . .                                                            | 6        |

|          |                                                                                |           |
|----------|--------------------------------------------------------------------------------|-----------|
| 4.3      | Three Approximant Types, By Design . . . . .                                   | 7         |
| 4.4      | Validation Across Four Oscillator Classes . . . . .                            | 7         |
| 4.5      | Why $\varepsilon = 50$ Works . . . . .                                         | 7         |
| 4.6      | Contributions of Paper III . . . . .                                           | 8         |
| <b>5</b> | <b>Paper IV (2019): Cosine GPA — Extending the Reach to Periodic Solutions</b> | <b>8</b>  |
| 5.1      | Motivation: The Periodicity Gap . . . . .                                      | 8         |
| 5.2      | The Cosine-Type Generalized Padé Approximant . . . . .                         | 8         |
| 5.3      | Two Improvement Schemes . . . . .                                              | 8         |
| 5.4      | Results Across Three Oscillator Classes . . . . .                              | 9         |
| 5.5      | Contributions of Paper IV . . . . .                                            | 9         |
| <b>6</b> | <b>Paper V (2022): Refining GPLP for Asymmetric Potentials</b>                 | <b>10</b> |
| 6.1      | Motivation . . . . .                                                           | 10        |
| 6.2      | Improvement 1: $\omega$ -Parameterized Homoclinic Approximant . . . . .        | 10        |
| 6.3      | Improvement 2: Type A and Type B Heteroclinic Approximants . . . . .           | 11        |
| 6.4      | Improvement 3: Rational Restoring Force Coverage . . . . .                     | 11        |
| 6.5      | Accuracy Gains: 2022 vs. 2016 . . . . .                                        | 11        |
| 6.6      | Contributions of Paper V . . . . .                                             | 12        |
| <b>7</b> | <b>The Unifying Theme: Basis-Geometry Correspondence</b>                       | <b>12</b> |
| <b>8</b> | <b>Limitations and Open Problems</b>                                           | <b>13</b> |
| 8.1      | Intrinsic Limitations of the GPA Framework . . . . .                           | 13        |
| 8.2      | Open Problems and Future Directions . . . . .                                  | 13        |
| <b>9</b> | <b>Conclusion</b>                                                              | <b>14</b> |

## 1 Introduction: Why Generalize Padé?

Classical Padé approximation [1] constructs a rational function—a ratio of two polynomials—whose Taylor expansion matches that of a given power series to the highest possible order. For over two centuries, it has been an indispensable tool in computational mathematics, quantum mechanics, field theory, and control theory.

When applied to nonlinear dynamics, however, classical Padé approximation encounters a fundamental limitation. Consider the general nonlinear oscillator:

$$\ddot{x} + g(x) = \varepsilon f(x, \dot{x}) \quad (1)$$

where  $g(x)$  is the restoring force and  $\varepsilon f(x, \dot{x})$  is a perturbation (damping, forcing, etc.). When  $\varepsilon = 0$  (conservative limit), Padé-based methods can construct highly accurate homoclinic and heteroclinic (H/H) orbits by approximating the power series solution [2,3]. However, when  $\varepsilon \neq 0$ , classical Padé approximants—being rational functions with limits at infinity—cannot adequately represent the complex geometry of damped H/H orbits or periodic solutions.

Meanwhile, perturbation-based methods (hyperbolic perturbation, elliptic Lindstedt–Poincaré, generalized harmonic function perturbation) can handle  $\varepsilon \neq 0$ , but at a heavy cost: each higher perturbation order demands increasingly complex derivation and integration. For oscillators with high-order polynomial, rational, or irrational restoring forces, these integrations quickly become analytically intractable.

The five papers reviewed here address this dilemma from a single conceptual direction: **generalize the definition of Padé approximation itself**. Once the numerator and denominator are freed from the constraint of being polynomial functions, the approximant can be tailored to the specific geometric and asymptotic properties of the solution being sought. This insight, first articulated in the Chinese-language foundational paper of 2013 [4], has evolved through four subsequent works into a versatile, integration-free framework for strongly nonlinear oscillator analysis.

## 2 Paper I (2013): The Generalized Padé Approximation — Definition and Foundation

### 2.1 Motivation

Classical Padé approximation [1] defines the approximant as:

$$[L/M]f(z) = \frac{\sum_{i=0}^L a_i z^i}{1 + \sum_{i=1}^M b_i z^i}, \quad f(z) - [L/M]f(z) = O(z^{L+M+1}) \quad (2)$$

While various “quasi-Padé” variants had been proposed—using exponentials [5,6], hyperbolic functions [2], or mixed bases—these modifications changed the *form* of the approximant but not the *definition*. The authors recognized that a deeper generalization was possible.

### 2.2 The GPA Definition

The core conceptual leap of [4] is to redefine the function spaces from which the numerator and denominator are drawn:

$$\hat{H}_m = \left\{ P : P(z) = \sum_{i=0}^m a_i g_i(z), \quad g_i : \mathbb{C} \rightarrow \mathbb{C}, \quad a_i \in \mathbb{C} \right\} \quad (3)$$

$$G(m, n) = \left\{ R = P/Q : P \in \hat{H}_m, \quad Q \in \hat{H}_n \setminus \{0\} \right\} \quad (4)$$

**Definition 1 (Generalized Padé Approximation).** Let  $f(z) = \sum_{i=0}^{\infty} a_i z^i$ . If there exist  $P_L(z)/Q_M(z) \in G(L, M)$  and an auxiliary operator  $\Gamma(z)$  such that:

$$\boxed{f(z) - \Gamma(z) \frac{P_L(z)}{Q_M(z)} = O(z^{L+M+1})} \quad (5)$$

then  $\Gamma(z)P_L(z)/Q_M(z)$  is a generalized Padé approximant of  $f(z)$ , denoted  $\text{GPA}[L/M]f(z)$ .

**Remarks.**

1. If  $g_i(z) = z^i$  and  $\Gamma(z) = 1$ , Definition 1 reduces exactly to the classical Padé definition.
2. All known quasi-Padé variants become special cases for appropriate choices of  $g_i(z)$  and  $\Gamma(z)$ .
3. The crucial freedom is the choice of **basis functions**  $g_i(z)$ —this is what enables the method to be adapted to specific solution geometries throughout the subsequent papers.

### 2.3 Application: Homoclinic Orbits via Hyperbolic Basis

To demonstrate the method's power, the authors construct a **hyperbolic-function-based** generalized Padé approximant for solving homoclinic orbits of strongly nonlinear conservative oscillators:

$$\text{GPA}[2/2]x(t) = \frac{\alpha_0 + \alpha_1 \text{Sech } t + \alpha_2 \text{Sech}^2 t}{1 + \beta_1 \text{Sech } t + \beta_2 \text{Sech}^2 t} \quad (6)$$

Why Sech? Because homoclinic orbits satisfy  $x(\pm\infty) = h_i$  (the saddle coordinate) and  $\dot{x}(\pm\infty) = 0$ —exactly the asymptotic behavior of the hyperbolic secant function. By matching the Taylor expansion of Eq. (6) with the power series solution  $\sum a_i t^i$  of the conservative equation  $\ddot{x} + g(x_0) = 0$ , the coefficients  $\alpha_i, \beta_i$  are determined by solving a system of algebraic equations—no integration required.




Figure 1: Homoclinic orbit of the Duffing oscillator obtained via  $\text{GPA}[2/2]$  (2013). Solid: Runge–Kutta. Dotted: GPA. The hyperbolic secant basis captures the orbit geometry with only 2nd-order approximants. From [4].

## 2.4 Key Contributions of Paper I

- **Generalized definition** that unifies classical Padé and all quasi-Padé variants under one framework.
- **First demonstration** that a non-polynomial, purpose-built basis (Sech) can dramatically improve accuracy for specific solution types.
- **Established the design principle** that drives all subsequent papers: *choose  $g_i(z)$  to match the asymptotic/geometric character of the solution.*

## 3 Paper II (2014): Extending GPA to Heteroclinic Orbits and Irrational Oscillators

### 3.1 Motivation

Paper I only addressed **homoclinic** orbits of oscillators with **polynomial** restoring forces. Two natural extensions were needed: (1) heteroclinic orbits connecting two different saddle points, and (2) oscillators with irrational restoring forces such as the smooth and discontinuous (SD) oscillator.

### 3.2 Methodological Extension

For heteroclinic orbits, the geometric requirements differ fundamentally from the homoclinic case:

- A homoclinic orbit approaches the *same* saddle as  $t \rightarrow +\infty$  and  $t \rightarrow -\infty$  — a symmetric asymptotic behavior naturally captured by Sech.
- A heteroclinic orbit approaches *two different* saddles — an asymmetric asymptotic behavior naturally captured by Tanh.

Accordingly, a Tanh-based approximant is constructed:

$$\text{GPA}[L/M]x(t) = \frac{\sum_{i=0}^L \alpha_i \text{Tanh}^i t}{1 + \sum_{i=1}^M \beta_i \text{Tanh}^i t} \quad (7)$$

For the SD oscillator  $\ddot{x} + x(1 - 1/\sqrt{x^2 + \alpha^2}) = 0$ , the irrational restoring force arising from a geometric constraint is handled without special treatment—the power series solution is simply substituted into the GPA matching procedure. The method successfully predicts both homoclinic and heteroclinic orbits for cubic-quintic Duffing,  $\Phi^6$ , and Duffing-harmonic oscillators.

### 3.3 Key Contributions of Paper II

- Extended the GPA method from homoclinic-only to both homoclinic and heteroclinic orbits.
- Established the **basis-geometry correspondence**: Sech  $\leftrightarrow$  homoclinic; Tanh  $\leftrightarrow$  heteroclinic.
- Demonstrated applicability to **irrational restoring forces** (SD oscillator).
- Proved that GPA handles polynomial, rational, and irrational  $g(x)$  within the same framework.

## 4 Paper III (2016): The GPLP Method — Marrying GPA with Lindstedt–Poincaré

### 4.1 Motivation

Papers I–II addressed the conservative case ( $\varepsilon = 0$ ) brilliantly. But to predict **bifurcations**—the critical parameter values at which limit cycles collide with saddle points to form H/H orbits—one




Figure 2: Homoclinic and heteroclinic orbits of the cubic-quintic Duffing oscillator (2014). Solid: Runge–Kutta. Dotted: GPA. Both orbit types are captured by choosing appropriate basis functions (Sech for homoclinic, Tanh for heteroclinic). From [7].

must handle the damped case ( $\varepsilon \neq 0$ ). This is where classical perturbation methods enter, and where their integration bottleneck becomes acute.

The key insight of Paper III [8] is that the perturbation equations arising from the Lindstedt–Poincaré expansion **do not need to be integrated**. Instead, each can be solved by constructing an appropriate generalized Padé approximant and matching coefficients—exactly as was done for the conservative case in Papers I–II.

## 4.2 The GPLP Procedure

The standard L–P expansion yields a hierarchy of perturbation equations:

$$\varepsilon^0 : \ddot{x}_0 + g(x_0) = 0 \quad (8)$$

$$\varepsilon^1 : \ddot{x}_1 + g'(x_0)x_1 = f(\mu_{c0}, x_0, \dot{x}_0) \quad (9)$$

$$\varepsilon^2 : \ddot{x}_2 + g'(x_0)x_2 + \frac{1}{2}g''(x_0)x_1^2 = \dots \quad (10)$$

$$\varepsilon^3 : \dots$$

In classical L–P, each  $x_n$  is obtained via the integral formula:

$$x_n = \dot{x}_0 \int \frac{1}{\dot{x}_0^2} \left[ \int \dot{x}_0 \cdot (\text{forcing terms}) dt \right] dt \quad (11)$$

GPLP replaces each integration with a tailored GPA:

**Step I**  $x_0$ : Solve via Sech-GPA (homoclinic) or Tanh-GPA (heteroclinic), as in Papers I–II.

**Step II**  $x_1$  and  $\mu_{c0}$ : Use a Tanh-prefixed polynomial approximant:

$$\text{GPA}[L_1/M_1]x_1(t) = \text{Tanh}(t) \frac{\sum_{i=0}^{L_1} \eta_i t^i}{1 + \sum_{i=1}^{M_1} \xi_i t^i} \quad (12)$$

The Tanh prefactor enforces  $x_1(\pm\infty) = 0$ . The condition  $\eta_{L_1}(\mu_{c0}) = 0$  yields the **zero-order critical bifurcation parameter**. Remarkably, this condition is equivalent to the classical Melnikov condition—providing rigorous theoretical grounding.

**Step III**  $x_2$  and  $d_0$ : Polynomial-based GPA;  $\lambda_{L_2} = 0$  determines the initial value correction.

**Step IV**  $x_3$  and  $\mu_{c2}$ : Polynomial-based GPA;  $\sigma_{L_3}(\mu_{c2}) = 0$  gives the second-order bifurcation parameter correction.

The assembled third-order solution:

$$x(t) = \sum_{i=0}^3 \varepsilon^i \text{GPA}[L_i/M_i]x_i(t) + O(\varepsilon^4), \quad \mu_c = \mu_{c0} + \varepsilon^2 \mu_{c2} + O(\varepsilon^3) \quad (13)$$

### 4.3 Three Approximant Types, By Design

A critical—and initially counterintuitive—design choice in GPLP is the use of **three different approximant forms** for different orders, rather than a single unified form. The rationale:

- $x_0$  uses Sech/Tanh because the basis must satisfy H/H asymptotic boundary conditions *exactly*.
- $x_1$  uses a Tanh-prefixed polynomial form because  $x_1(\pm\infty) = 0$  must be enforced.
- $x_2, x_3$  use pure polynomial GPA because asymptotics are already embedded in lower orders, and polynomial forms offer simpler coefficient matching.

The paper explicitly compares this mixed strategy against a single-form approach (Example 5), confirming that the tailored strategy yields substantially higher accuracy.




Figure 3: Homoclinic orbits via GPLP for the Helmholtz–Duffing–Van der Pol oscillator at  $\varepsilon = 2$  and 3 (2016). Solid: Runge–Kutta. Dash-dot: GPLP ( $\varepsilon^3$ ). From [8].

### 4.4 Validation Across Four Oscillator Classes

GPLP is validated on oscillators spanning all canonical restoring-force types:

1. **Helmholtz–Duffing–Van der Pol** (cubic polynomial):  $\mu_{c0} \approx -6.3941$ , accurate at  $\varepsilon = 2, 3$ .
2. **Duffing-harmonic–Van der Pol** (rational):  $\mu_{c0} = -1.9026$ .
3. **SD oscillator** (irrational): accurate at  $\varepsilon = 50, 60$ —remarkably large perturbation parameters.
4.  **$\Phi^6$ -Van der Pol** (quintic): simultaneous homoclinic and heteroclinic bifurcation prediction.

### 4.5 Why $\varepsilon = 50$ Works

This is a frequently asked question. The answer has two parts. First, the zero-order GPA solution for the conservative system already closely matches the exact orbit—often within 1%. The perturbation corrections  $\varepsilon x_1, \varepsilon^2 x_2, \varepsilon^3 x_3$  modify this nearly-exact conservative orbit incrementally. Since the zero-order baseline is so accurate, even large  $\varepsilon$  does not cause large absolute errors. Second, the GPA approximants at each order are “global” rational functions, not local Taylor expansions—they capture the full orbit geometry rather than a local neighborhood.




Figure 4: SD oscillator homoclinic orbits at  $\varepsilon = 50$  and  $60$  (2016). Solid: Runge-Kutta. Dash-dot: GPLP. The  $\varepsilon^3$  solution remains accurate at perturbation parameters two orders of magnitude beyond typical “weakly nonlinear” regimes. From [8].

#### 4.6 Contributions of Paper III

- **GPLP: the first integration-free perturbation method** for H/H bifurcation prediction.
- **Melnikov equivalence:** the GPLP condition  $\eta_{L_1}(\mu_{c0}) = 0$  is proved equivalent to the classical Melnikov condition.
- **Third-order accuracy at  $\varepsilon = 50$ – $60$ :** unprecedented for a perturbation-based method.
- **Simultaneous H/H prediction:** both critical values from a single framework.

### 5 Paper IV (2019): Cosine GPA — Extending the Reach to Periodic Solutions

#### 5.1 Motivation: The Periodicity Gap

Papers I–III all addressed H/H orbits—solutions that are non-periodic and approach saddle points asymptotically. But the vast majority of vibration problems involve **periodic solutions**. Classical Padé approximants, being rational functions with limits at infinity, are fundamentally unsuited for representing periodic functions. No prior work had directly applied Padé approximation to periodic solution problems.

The question posed by Paper IV [9] is: can we design a generalized Padé approximant that is *itself* periodic?

#### 5.2 The Cosine-Type Generalized Padé Approximant

The answer is elegantly simple: construct the numerator and denominator from cosine functions:

$$x(t) = \frac{\sum_{i=0}^m \alpha_i \cos(i\omega t)}{\sum_{j=0}^n \beta_j \cos(j\omega t)} \quad (14)$$

Since each  $\cos(i\omega t)$  is  $2\pi/\omega$ -periodic, any rational function of cosines inherits this periodicity. The intrinsic contradiction between Padé approximation and periodic solutions is resolved not by modifying the approximation *procedure*, but by modifying the *mathematical structure* of the approximant itself.

#### 5.3 Two Improvement Schemes

Classical Padé matching (expand approximant as Taylor series; compare with power series of the solution) encounters a difficulty here: the unknown frequency  $\omega$  makes the matching equations nonlinearly coupled. Two improvement schemes are developed:



**Scheme I (Active frequency iteration):** Treat  $\omega$  as an active variable with increment  $\Delta\omega$ , iterating from  $\omega_0$ . At each step, solve for  $\alpha_i, \beta_j$ . Terminate when the target function  $\Gamma = x(\pi/\omega) - x_1$  (where  $x_1$  is the other  $x$ -axis intersection of the orbit) satisfies  $|\Gamma| < 10^{-(p+4)}$ .

**Scheme II (Derivative-based target + piecewise description):** For rational restoring forces with longer-period orbits, switch the target function to the derivative at half-period and use piecewise matching at the half-period point. This restores accuracy for cases where Scheme I underperforms.

## 5.4 Results Across Three Oscillator Classes

The cosine GPA method is demonstrated on:

- **Duffing oscillator** (polynomial):  $\ddot{x} + x + x^3 = 0$ ,  $A = 2$ . Absolute error  $< 10^{-3}$ .
- **Duffing-harmonic oscillator** (rational):  $\ddot{x} + \frac{x+x^3}{1+x^2} = 0$ ,  $A = 3$ .
- **SD oscillator** (irrational):  $\ddot{x} + x(1 - 1/\sqrt{x^2 + \alpha^2}) = 0$ ,  $\alpha = 1$ ,  $A = 2.5$ .




Figure 5: Phase portrait, time history, and absolute error for the Duffing oscillator with  $A = 2$  (2019). The cosine GPA method maintains error below  $10^{-3}$  across the full period. From [9].

## 5.5 Contributions of Paper IV

- **First direct Padé approximation of periodic solutions:** Achieved by constructing a *periodic* approximant rather than modifying the approximation procedure.
- **Methodological generality:** Covers polynomial, rational, and irrational restoring forces.
- **Period prediction:** The frequency  $\omega$  obtained from the matching procedure yields the analytic period  $T = 2\pi/\omega$ .
- **Conceptual bridge:** Completed the GPA framework by extending it from non-periodic H/H orbits to periodic solutions.




Figure 6: Duffing-harmonic oscillator (rational restoring force),  $A = 3$  (2019). Scheme II restores accuracy for longer-period orbits. From [9].

## 6 Paper V (2022): Refining GPLP for Asymmetric Potentials

### 6.1 Motivation

The 2016 GPLP method handled **symmetric** potentials ( $c_1x + c_3x^3 + c_5x^5$ ) admirably. However, three practical limitations remained:

1. Realistic oscillators often have **all five polynomial terms** ( $x^2$  and  $x^4$  included), creating asymmetric multi-well potentials.
2. The 2016 heteroclinic approximant (pure Tanh) showed 5–8% error under strong asymmetry.
3. Rational restoring forces had not been tested in the GPLP framework.

Paper V [10] addresses all three with three targeted improvements.

### 6.2 Improvement 1: $\omega$ -Parameterized Homoclinic Approximant

The 2016 version hard-coded  $\omega = 1$ :

$$x(t) = \frac{\sum \alpha_i \operatorname{Sech}^i t}{1 + \sum \beta_i \operatorname{Sech}^i t} \quad \longrightarrow \quad \boxed{x(t) = \frac{\sum_{i=0}^L \alpha_i \operatorname{Sech}^i(\omega t)}{1 + \sum_{i=1}^M \beta_i \operatorname{Sech}^i(\omega t)}} \quad (15)$$

The free parameter  $\omega$  adapts to different potential well shapes—critical when left and right wells have different curvatures. This single change reduces homoclinic error by 40–60% for asymmetric cases.

### 6.3 Improvement 2: Type A and Type B Heteroclinic Approximants

**Type A** (rational Tanh, with free  $\omega$ ):

$$\text{GPA}[L/M]x(t) = \frac{\sum_{i=0}^L \alpha_i \text{Tanh}^i(\omega t)}{1 + \sum_{i=1}^M \beta_i \text{Tanh}^i(\omega t)} \quad (16)$$

Sufficient for pure heteroclinic problems.

**Type B** (combined Sech-Tanh) — the paper’s most important innovation:

$$\text{GPA}[L, M]x(t) = \frac{\sum_{i=0}^{L_1} \alpha_i \text{Sech}^i(\omega t) + \sum_{j=0}^{L_2} \gamma_j \text{Tanh}^j(\omega t)}{1 + \sum_{i=1}^{M_1} \beta_i \text{Sech}^i(\omega t) + \sum_{j=1}^{M_2} \delta_j \text{Tanh}^j(\omega t)} \quad (17)$$

The geometric motivation behind Type B is central to understanding the entire GPA research program. In an asymmetric potential, the higher saddle’s homoclinic orbit has a **Sech-like** symmetric approach, while the heteroclinic orbit connecting unequal saddles has a **Tanh-like** asymmetric character. When both coexist at the same saddle, a pure Tanh approximant cannot simultaneously capture the homoclinic orbit’s symmetry. Type B resolves this by leveraging both basis functions simultaneously—reducing heteroclinic error from 5–8% to  $< 1.5\%$ .




Figure 7: Homo-heteroclinic orbits via GPLP with Type B approximant at  $\varepsilon = 0.8$  and 1 (2022). Dashed: GPLP. Solid: Runge–Kutta. The combined Sech-Tanh approximant captures both orbit types simultaneously. From [10].

### 6.4 Improvement 3: Rational Restoring Force Coverage

The generalized Duffing-Harmonic-Van der Pol oscillator:

$$\ddot{x} + \frac{\lambda x + \delta x^3}{1 + \nu x^2} = \varepsilon(\mu_c + \mu_1 x + \mu_2 x^2)\dot{x} \quad (18)$$

is solved within the GPLP framework **without any procedural modification**. Only the coefficients in the perturbation equations change. Validated from  $\varepsilon = 0.5$  to  $\varepsilon = 2$ .

### 6.5 Accuracy Gains: 2022 vs. 2016

Table 1: Accuracy comparison between the 2016 and 2022 GPLP versions.

| Scenario                                   | 2016 Error   | 2022 Error | Improvement    |
|--------------------------------------------|--------------|------------|----------------|
| Symmetric potential, small $\varepsilon$   | $\sim 1\%$   | $\sim 1\%$ | Similar        |
| Asymmetric homoclinic, $\varepsilon = 1$   | $\sim 2.5\%$ | $< 1\%$    | 40–60%         |
| Asymmetric heteroclinic, $\varepsilon = 1$ | 5–8%         | $< 1.5\%$  | 70–80%         |
| Rational restoring force                   | N/A          | $< 2\%$    | New capability |

## 6.6 Contributions of Paper V

- **Full  $\Phi^6$  coverage:** GPLP extended to the full five-term quintic polynomial with asymmetric multi-well potentials.
- **Combined Sech-Tanh approximant (Type B):** 70–80% heteroclinic accuracy improvement in asymmetric regimes.
- **Rational restoring force coverage:** DH-VDP oscillator solved without procedural modification.
- **Completes the GPLP framework** for polynomial and rational restoring forces.

## 7 The Unifying Theme: Basis-Geometry Correspondence




Figure 8: Technical roadmap of the generalized Padé approximation (GPA) research program. The five papers progress from the foundational definition (2013) through method extension (2014), integration with perturbation theory—GPLP (2016), extension to periodic solutions (2019), and asymmetric refinement (2022). The core design principle—matching basis functions to solution geometry (left sidebar)—guides every methodological choice. The bottom timeline summarizes the basis-geometry correspondence across the entire series.

Looking across all five papers, a single design principle emerges that explains every major methodological choice:

**The basis functions  $g_i(z)$  of the generalized Padé approximant should match the asymptotic/geometric character of the solution being approximated.**

This principle manifests differently across the papers:

Table 2: Evolution of the basis-geometry correspondence across the five papers.

| Paper | Year | Solution Type       | Basis Functions                                                |
|-------|------|---------------------|----------------------------------------------------------------|
| I     | 2013 | Homoclinic orbits   | $\text{Sech}^i t$ (hyperbolic secant)                          |
| II    | 2014 | Heteroclinic orbits | $\text{Tanh}^i t$ (hyperbolic tangent)                         |
| III   | 2016 | H/H bifurcations    | $\text{Sech}/\text{Tanh}$ (order 0) + polynomial (orders 1–3)  |
| IV    | 2019 | Periodic solutions  | $\cos(i\omega t)$ (cosine series)                              |
| V     | 2022 | Asymmetric H/H      | $\text{Sech}^i(\omega t) + \text{Tanh}^j(\omega t)$ (combined) |

The progression from Sech to Tanh to mixed Sech-Tanh to cosine is not arbitrary—each choice is justified by the geometry of the target solution:

- **Sech:**  $x(\pm\infty) = h$  (symmetric approach to a single saddle).
- **Tanh:**  $x(+\infty) = h_i$ ,  $x(-\infty) = h_j$ ,  $i \neq j$  (asymmetric connection between two saddles).
- **Cosine:**  $x(t + 2\pi/\omega) = x(t)$  (periodicity).
- **Combined Sech-Tanh:** Coexisting homoclinic and heteroclinic orbits at a single saddle.

This is not merely a “choice of basis” in the linear algebra sense—it is a deliberate exploitation of the generalized Padé definition’s flexibility to encode physical knowledge about the solution directly into the approximant’s functional form.

## 8 Limitations and Open Problems

### 8.1 Intrinsic Limitations of the GPA Framework

1. **Pre-set parameters:** System parameters must be specified before calculation. Unlike algebraic methods (e.g., MGHP), GPA cannot trace continuous parameter-solution dependencies.
2. **Manual order selection:** The approximation orders  $L_i, M_i$  must be chosen by the user based on desired accuracy—no automated selection criterion exists.
3. **Algebraic complexity at high orders:** Although integration is avoided, the Taylor expansion and coefficient matching for the perturbation expansion (Eq. 20 in [8]) remains algebraically heavy at orders beyond  $\varepsilon^3$ .
4. **Convergence theory:** No rigorous convergence proof exists for the generalized Padé approximants constructed in these papers. Their accuracy is validated empirically (vs. Runge–Kutta), not theoretically guaranteed.

### 8.2 Open Problems and Future Directions

1. **Quasi-periodic and chaotic solutions:** Can mixed-basis approximants (e.g., cosine + exponential) capture quasi-periodic or weakly chaotic orbits? The GPA definition permits such constructions, but no implementation exists.
2. **Automated basis selection:** Given an oscillator and a target solution type, can the optimal basis functions  $g_i(z)$  be determined algorithmically rather than by the analyst’s intuition?
3. **Extension to PDEs:** The GPA framework is formulated for ODEs. Can it be adapted to nonlinear PDEs describing continuous structures (beams, plates, shells)?
4. **Integration with machine learning:** The coefficient matching problem is a nonlinear algebraic system. Could neural networks accelerate the solution of these systems, enabling real-time parametric studies?
5. **Multi-degree-of-freedom systems:** All five papers address single-DOF oscillators. The extension to coupled oscillators remains largely unexplored.

## 9 Conclusion

The generalized Padé approximation research program, spanning five papers from 2013 to 2022, demonstrates the power of a single conceptual insight—generalizing the definition of Padé approximation to allow arbitrary continuous basis functions—when systematically pursued across different problem domains. The arc from homoclinic orbits (Paper I) to heteroclinic orbits (Paper II), to integration-free bifurcation prediction (Paper III), to periodic solutions (Paper IV), and finally to asymmetric-potential refinement (Paper V) represents a rare example in the nonlinear dynamics literature of a sustained methodological research program built on a coherent design philosophy.

The practical value of this framework is twofold. For the analyst, it provides a family of methods—GPA for conservative systems, GPLP for damped bifurcation analysis, cosine GPA for periodic solutions—all sharing the same conceptual foundation and the same “no-integration” workflow. For the methodologist, it illustrates a general strategy: when an existing approximation framework proves inadequate for a new problem class, consider whether *generalizing the definition of the approximant itself*—rather than patching the procedure—might provide a more elegant and powerful solution.

## References

- [1] G.A. Baker Jr. and P. Graves-Morris, *Padé Approximants*, 2nd ed., Cambridge University Press, 1996.
- [2] Y.V. Mikhlin, “Construction of homoclinic and heteroclinic trajectories of two and three-dimensional dynamical systems by the Padé approximation,” *J. Sound Vib.*, vol. 230, pp. 889–913, 2000.
- [3] G.V. Manucharyan and Y.V. Mikhlin, “Construction of homoclinic and heteroclinic trajectories of non-autonomous Duffing equation,” *Int. J. Non-Linear Mech.*, vol. 37, pp. 529–540, 2002.
- [4] Z. Li, J. Tang, and P. Cai, “Generalized Padé approximation method for homoclinic orbits of strongly nonlinear autonomous oscillators” (in Chinese), *Chin. J. Theor. Appl. Mech.*, vol. 45, no. 3, pp. 461–464, 2013.
- [5] J.F. Martin and G.A. Baker Jr., “Quasi-Padé approximants,” *J. Math. Phys.*, vol. 32, pp. 1470–1477, 1991.
- [6] Q.C. Zhang, J.J. Feng, and W. Wang, “Application of quasi-Padé approximation in two-dimensional vibration systems” (in Chinese), *Chin. J. Theor. Appl. Mech.*, vol. 43, no. 4, pp. 658–664, 2011.
- [7] Z. Li, J. Tang, and P. Cai, “A generalized Padé approximation method of solving homoclinic and heteroclinic orbits of strongly nonlinear autonomous oscillators,” *Chin. Phys. B*, vol. 23, no. 12, 120501, 2014.
- [8] Z. Li and J. Tang, “A generalized Padé–Lindstedt–Poincaré method for predicting homoclinic and heteroclinic bifurcations of strongly nonlinear autonomous oscillators,” *Nonlinear Dyn.*, vol. 84, no. 3, pp. 1201–1223, 2016.
- [9] Z. Li and J. Tang, “A cosine generalized Padé approximation method and its application in solving periodic solutions of strongly nonlinear oscillators,” *J. Vib. Shock*, vol. 38, no. 22, pp. 159–167, 2019.
- [10] Z. Li and J. Tang, “High accurate homo-heteroclinic solutions of certain strongly nonlinear oscillators based on generalized Padé–Lindstedt–Poincaré method,” *J. Vib. Eng. Technol.*, vol. 10, no. 4, pp. 1291–1308, 2022.